

Main panel bottle

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS



Amgrow
Chemspray
DISEASE CONTROL

COPPER OXYCHLORIDE
Fungicide

ACTIVE CONSTITUENT:
500 g/kg COPPER (Cu) present as COPPER OXYCHLORIDE

For the control of certain fungal diseases of fruit trees, vegetables and
ornamentals in the home garden.

Before using product, read directions on outer pack.
Sale of this bottle except in outer pack is illegal.

NET CONTENTS 200 g

Ancillary panel bottle

STORAGE & DISPOSAL: Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS: Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126.

NOTE: this product is sold by weight not volume. The contents may settle during transit.

APVMA Approval No. 30306/52939

Amgrow Pty Ltd

A.B.N. 81 100 684 786

82 Christensen Road, Stapylton Qld 4207

Customer Service Number

1800 063 619

www.amgrow.com.au

Batch number:

Date of Manufacture:

BAR CODE

Main panel carton

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS

Amgrow
Chemspray
DISEASE CONTROL

COPPER OXYCHLORIDE
Fungicide

ACTIVE CONSTITUENT:
500 g/kg COPPER (Cu) present as COPPER OXYCHLORIDE

**For the control of certain fungal diseases of fruit trees, vegetables and
ornamentals in the home garden.**

**The bottle must not be sold separately.
Do not destroy box while product still remains.**

NET CONTENTS 200 g

Ancillary panel 1 carton

HOW TO USE:

Crop	Disease	State	Rate	How To Apply
Apples Pears	Black Spot (Scab)	All States	5g per 2 litres of water + 5mL white oil	Apply at green tip.
Avocados	Anthrachnose	Qld, NSW, NT, SA, Vic only	5g per 2 litres of water	Apply from flowering to harvest at 28-day intervals.
Beans (French)	Common Blight	SA, Tas, Vic, WA only	5g per 2 litres of water	Apply every 10-14 days when disease first appears.
	Bacterial Brown Spot	All States		Apply the first spray within 3 weeks of emergence and repeat every 10-14 days.
Brassicas (cabbage, broccoli etc)	Black Spot, Peppery leaf spot, Ring Spot, Downy Mildew	All States	5g per 2 litres of water	Apply every 14 days.
Capsicums	Bacterial Spot	All States	5g per 2 litres of water	Apply every 10-14 days.
Carrots	Leaf Spot	All States	5g per 2 litres of water	Apply at 7-14 day intervals.
Celery	Septoria Spot	All States	5g per 2 litres of water	Apply at 7-14 day intervals.
	Early Blight	NSW, Vic, SA, WA, NT only		
Citrus	Black Spot Melanose	NSW, SA, Vic, WA only	7g per 4 litres of water + 6mL of white oil	Apply at 50-80% petal fall and again after 6 –12 weeks if necessary.
Cucurbits (Cucumbers, Melons etc)	Angular Leaf Spot, Bacterial Leaf Spot	All States	5g per 2 litres of water	Apply at 10-14 day intervals.
Grape vines	Downy Mildew	NSW, Qld, SA, Tas, Vic, WA only	5g per 2 litres of water	Commence when shoots are approx. 20cm long and apply at 14-21 day intervals. Do not apply during flowering.
Lettuce	Downy Mildew	All States	5g per 2 litres of water	Spray when disease first appears or as a preventative and repeat every 10 days if necessary.
Onions	Downy Mildew	All States	5g per 2 litres of water	Apply at 10 day intervals.
Ornamentals (Flowers and Shrubs)	Fungal Leaf Spots, Downy Mildew	All States	3g per litre of water	Apply when disease first appears. Repeat at 7-14 day intervals.
Passionfruit	Brown Spot, Septoria Spot	Qld only	2g per litre of water	Apply every 7-14 days.
		NSW, NT, Vic, SA, Tas, WA only	3g to 4g per litre of water	
Peas	Ascochyta Blight, Bacterial Blight	All States	5g per 2 litres of water	Apply at 7-14 day intervals.
Potatoes	Target Spot (Early Blight), Late (or Irish) Blight	All States	5g per 2 litres of water	Apply at 7-14 day intervals.
Silver Beet Spinach	Downy Mildew	All States	5g per 2 litres of water	Apply at 10-14 day intervals.

Stone Fruit (eg peaches, nectarines)	Blossom Blight, Freckle, Rust	NSW, Vic, Tas, SA, WA only	4g per litre of water	Apply at budswell.
	Leaf Curl, Shot Hole			Apply at budswell. For shot hole apply again at leaf fall..
Tomatoes	Bacterial Spot	All States	5g per 2 litres of water	Apply at 7-14 day intervals.
	Bacterial Speck			
	Target Spot/Early Blight			
	Septoria Leaf Spot, Irish & Late Blight			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.
DO NOT PICK FRUIT AND VEGETABLES FOR 1 DAY AFTER SPRAYING.**

HOW TO PREPARE

Add the required quantity of product to water and stir thoroughly. Mix required oil with equal volume of water prior to adding to spray mixture. Carefully follow recommended rates and timing for application.

CAUTION: DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

STORAGE & DISPOSAL: Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS: Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126.

NOTE: this product is sold by weight not volume. The contents may settle during transit. APVMA Approval No. 30306/52939

Amgrow Pty Ltd

A.B.N. 81 100 684 786

82 Christensen Road, Stapylton Qld 4207

Customer Service Number

1800 063 619

www.amgrow.com.au

Batch number:

Date of Manufacture:

BAR CODE