

Final label following the reconsideration of Omethoate



Company Name: Arysta Lifescience North America LLC
Product Name: Folimat 800 Insecticide Spray
APVMA Approval Number: 33055/RV1216_02

Label name:	Folimat 800 Insecticide Spray
Signal headings:	DANGEROUS POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent statements:	ACTIVE CONSTITUENT: 800 g/L OMETHOATE (an anti-cholinesterase compound)
Mode of action	GROUP 1B INSECTICIDE
Statement of Claims:	For the control of a certain pests on ornamentals as specified in the DIRECTIONS FOR USE table. IMPORTANT: Read the attached booklet before use
Net Contents:	CONTENTS: 5L, 20L
Restrains:	DO NOT apply by airblast equipment. DO NOT apply with boomspray equipment unless operators are protected by enclosed cabs. DO NOT use open mixing/loading systems for boom spray and aerial equipment. DO NOT apply by spray equipment carried on the back of the user.
Directions for Use:	Attached below table.
Other limitations:	DO NOT use this product in the Home Garden
Withholding Periods:	WITHHOLDING PERIOD: Not applicable if used according to label instructions

Trade Advice:	-
----------------------	---

General instructions:	<p>Mixing Add the required quantity of Folimat to water in the spray vat while stirring or with agitators in motion.</p> <p>Note on ornamentals Folimat 800 Insecticide Spray has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.</p> <p>Compatibility Folimat is compatible with Antracol ®, white oil, misting oil, water miscible oil. DO NOT mix concentrates together but add each to the spray tank separately. As formulations of other manufacturer's products are beyond the control of Arysta Lifescience all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Arysta Lifescience cannot be responsible for the behaviour of such mixtures.</p>
------------------------------	---

Resistance warning:	<p>Insecticide Resistance Warning GROUP 1B INSECTICIDE</p> <p>For insecticide resistance management Folimat is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to Folimat and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Folimat or other Group 1B insecticides are used repeatedly. The effectiveness of Folimat on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Arysta Lifescience accepts no liability for any losses that may result from the failure of Folimat to control resistant insects. Folimat may be subject to specific resistance management strategies. For further information contact your local supplier, Arysta Lifescience representative or local agricultural department agronomist.</p>
----------------------------	---

Precautions:	<p>RE-ENTRY INTERVALS:</p> <p>Cut flowers hand harvesting DO NOT allow hand harvesting of cut flowers for 29 days. If prior entry is required for hand harvesting, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.</p> <p>Floriculture, nursery and greenhouse crops (ornamentals non-bearing) (except hand harvesting of cut flowers) DO NOT allow entry into treated areas of nursery and greenhouse crops for all activities (other than hand harvesting of cut flowers) until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.</p> <p>Tree crops (forestry) DO NOT allow entry into treated areas for 17 days for tree crops. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.</p>
---------------------	---

Protections:	<p>PROTECTION OF LIVESTOCK Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
Storage & disposal:	<p>STORAGE AND DISPOSAL Store in the closed, original container in a cool, dry and well ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging below 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.</p>
Safety Directions:	<p>SAFETY DIRECTIONS</p> <p>Very dangerous, particularly the concentrate. Product and spray are poisonous if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin and clothing. DO NOT inhale spray mist.</p> <p>When opening the container and preparing spray for aerial spraying equipment wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, water resistant footwear and full face-piece respirator with gas/dust cartridges.</p> <p>When opening the container and preparing spray for boom spray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, water resistant footwear and face shield or goggles.</p> <p>When opening the container and preparing spray for high pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, water resistant footwear and face shield or goggles.</p> <p>If applying by boom spray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.</p> <p>If applying by high pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrists (or equivalent clothing) and a washable hat, elbow-length chemical resistant gloves and full face-piece respirator with gas/dust cartridges.</p> <p>If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.</p>
First Aid:	<p>FIRST AID</p> <p>If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131 126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be</p>

	advised. Give atropine if instructed.
--	---------------------------------------

First Aid Warnings:	
----------------------------	--

DIRECTIONS FOR USE

Non-tree/Vine crops

CROP	PEST CONTROLL ED	STATE	RATE	WHP Days	CRITICAL COMMENTS
Carnations, Chrysanthemums, Pelargoniums, Roses, Callistemons, Eucalyptus spp., Grevillea spp., Paperbarks, Wattles	Aphids, lace bugs, mealybugs, mites, thrips, whiteflies	All States	75 mL/100 L	-	Spray thoroughly as a full cover spray. Repeat at beginning of renewed pest activity. See note on ornamentals.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION