

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ACCENSI

# Bromacil 800WP Herbicide

**ACTIVE CONSTITUENT: 800 g/kg BROMACIL**



**GROUP C HERBICIDE**

**For the control of certain annual and perennial broadleaved weeds and grasses in citrus, pineapple, asparagus and in commercial and industrial areas, rights of way and around agricultural buildings.**

**Label:**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING**

**Leaflet:**

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING**

**Net Contents: 2kg, 10kg  
The bag must not be sold separately.  
DO NOT destroy box while product still remains.**

**ACCENSI Pty Ltd ABN 94 079 875 184  
60-76 Potassium Street, Narangba Qld 4504  
Ph: (07) 3897 2000**

**APVMA Approval No.: 45401 / 59867**



Batch Number

Date of Manufacture

RELEVANT LABEL PARTICULARS

**DIRECTIONS FOR USE**

**Restrains:** DO NOT allow spray to contact foliage or fruit.

CROP & SITUATION	WEEDS CONTROLLED	STATE	RATE/ha		CRITICAL COMMENTS
			Light soils	Heavy soils	
Asparagus	Most annual weeds	NSW, Vic, Tas, SA, WA, NT only	1.1 kg	2.5 kg	Apply after cultivation to established plantings only. Apply 3-4 weeks prior to spear emergence in Spring or before the early cutting period. If cultivation occurs during cutting period apply immediately after last cultivation. If needed apply a second spray after harvest to keep fields weed-free. Irrigate before and as soon as possible after spraying.
	Hard-to-kill perennials such as Johnson Grass, Couch Grass, Kikuyu Grass and Nut Grass		4.5 kg	6.5 kg	Apply preferably in a split application 2 months apart. May take 1-2 years to effectively bring these weeds under control.
			1 kg/500 L	1 kg/500 L	Spot Spray – wet plant thoroughly to point of run-off.
Citrus – established groves only	Most annual weeds	All States	2.2 kg	4.5 kg	Apply to weed-free ground in Autumn and Spring. If weeds are present add a wetting agent at the use rate on its label. After 1-2 years of application, once per year may be adequate. DO NOT apply more than 9 kg/ha/year.
	Hard-to-kill perennials such as Couch Grass, Nut Grass, Johnson Grass, Kikuyu, Skeleton Weed		4.5 kg	6.5 kg	Apply just prior to period of most active weed growth. Irrigate with 25-75 mm as soon as possible after application. Rates may be reduced when weeds are brought under control.
			2 kg/500 L	2 kg/500 L	Spot treatment – spray to point of run-off. If weeds are present add a wetting agent at the use rate on its label.
Pineapple	Most annual weeds	Qld only	2.2 kg	4.5 kg	Apply as a blanket spray prior to weed emergence immediately after planting or before planting material begins to grow.
			2.2 kg	2.2 kg	If needed apply as a directed inter-row spray prior to flower differentiation. If weeds are present add a wetting agent at the use rate on its label. DO NOT spray ovetop of plants. DO NOT apply more than 9 kg/ha/year.
Industrial Weed Control Commercial and Industrial areas, rights of way, around agricultural buildings	ANNUAL WEEDS Barley Grass, Bathurst Burr, Brome Grass, Caltrop, Capeweed, Cobbler's Pegs, Three-cornered Jack, Double Gee (Spiny Emex), Fat Hen, Khaki Weed (seedling), Noogoora Burr, Paterson's Curse, Prairie Grass, Annual Ragweed, Annual Ryegrass, Saffron Thistle, Summer Grass, Thornapple, Wild Mustard, Wild Oats, Wild Radish, Wild Turnip, Winter Grass	All States	3.5 kg	6.5 kg	The lowest rate will provide control in low rainfall areas (250 mm or less). If weeds are present add a wetting agent at the use rate on its label.
	PERENNIAL WEEDS Bahia Grass, Bluetop, Guinea Grass, Horehound, Para Grass, Purpletop, Plaintain, St. John's Wort, Soursob		8 kg	13 kg	
	HARD-TO-KILL PERENNIALS Bindweed (Convolvulus), Couch Grass, Dock, Johnson Grass, Kikuyu, Nutgrass, Skeleton weed, Sorrel, Water Couch		16.5 kg	22 kg	
	RETREATMENT All the above weeds		2 kg	6.5 kg	
	SMALL AREAS Annual and Perennial weeds			20 g/10 m <sup>2</sup> (20 kg/ha)	20 g/10 m <sup>2</sup> (20 kg/ha)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## RELEVANT LABEL PARTICULARS

### GENERAL INSTRUCTIONS

For best results apply shortly before weed growth occurs to moist well prepared trash and clod free soil. Well-established weeds should be removed before applying. DO NOT cultivate soil surface after application as efficacy may be reduced and/or crop injury may result. For best results sufficient moisture, either by rainfall or irrigation is required within 3-4 days of treatment to carry the product into the soil. With irrigation thoroughly wet the entire area. Where rainfall is limited during the growth period, control of deep rooted hard to kill perennial weeds will not be satisfactory.

### HERBICIDE RESISTANCE WARNING

ACCENSI Bromacil 800WP Herbicide is a member of the uracils group of herbicides. ACCENSI Bromacil 800WP has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management ACCENSI Bromacil 800WP is a Group C herbicide. Some naturally occurring weed biotypes resistant to ACCENSI Bromacil 800WP and other inhibitors of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ACCENSI Bromacil 800WP or other inhibitors of photosynthesis at photosystem II herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, ACCENSI Pty Ltd accepts no liability for any losses that may result from the failure of ACCENSI Bromacil 800WP to control resistant weeds.

**GROUP C HERBICIDE**

### EQUIPMENT AND APPLICATION

Mix proper amount of the product into a minimum of 20L water per kg of product.

*Large Areas:* Use a fixed boom sprayer or handgun. To prevent nozzle blockage, strainer and nozzle screens should be 50 mesh or coarser. Always agitate in spray tank to keep material in suspension.

a) Fixed boom sprayer: Use 1000-2000 L/ha water for industrial weed control.

For selective weed control use 250-450 L/ha at 200-300 kPa pressure.

b) Hand gun: Use 1000-2000 L/ha and 600-800 kPa pressure.

*Small Areas:* Use a knapsack or pressurised hand sprayer or watering can. Shake or stir frequently.

### CROP ROTATIONS

DO NOT replant asparagus and citrus treated areas to any crop within 3 years after application. For pineapple crops only pineapples can be replanted into the same area within the 3 year period.

### CROP SAFETY

**Asparagus:** DO NOT apply to newly seeded asparagus, or to young plants during the first growing season after setting or to plants whose roots are exposed.

**Citrus:** Use with caution on light sandy soils.

### PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

**KEEP OUT OF REACH OF CHILDREN.** Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. On-site disposal of undiluted chemicals is unacceptable. Single rinse before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Puncture and bury empty bags and boxes in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

### SAFETY DIRECTIONS

May irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE:** The use of ACCENSI Bromacil 800WP Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by ACCENSI Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and ACCENSI Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.