

FRONT (MAIN) PANEL

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



AgrEVO

Acaban[®]

MITICIDE

Active Constituent: 50 g/L FENPYROXIMATE
Pyrazole Miticide

Infopest verifies that this label is consistent with the
NRA Approved Text Label

1 L

Controls Two Spotted Mite and European Red Mite in Apples and Pears.
IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.

FRONT (MAIN) PANEL

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



AgrEVO

Acaban[®]

MITICIDE

Active Constituent: 50 g/L FENPYROXIMATE
Pyrazole Miticide

2.5 L

Controls Two Spotted Mite and European Red Mite
in Apples and Pears.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.

FRONT (MAIN) PANEL

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Agrevo

Acaban[®]

MITICIDE

Active Constituent: 50 g/L FENPYROXIMATE
Pyrazole Miticide

5 L


Controls Two Spotted Mite and European Red Mite in Apples and Pears.
IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.

ANCILLARY PANEL (5 L pack only)

ACABAN MITICIDE

DIRECTIONS FOR USE

**Full directions are in the leaflet enclosed in the attached pocket.
This must be read for detailed directions for use.**



NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

REAR PANEL**ACABAN MITICIDE****Protection of Wildlife, Fish, Crustaceans and Environment**

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with product or used containers.

Storage and Disposal

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes. Avoid contact with eyes and skin. Avoid inhaling spray mist. When preparing spray and using the prepared spray wear elbow-length PVC gloves, goggles and half facepiece respirator. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and respirator (if rubber, wash with detergent and warm water).

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF.

Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet available from AgrEvo Pty. Ltd.

Exclusion of Liability

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

NRA Approval No.: 45466/0898

Acaban[®] is a Registered Trademark of Novartis AG, Basel, Switzerland
Acaban is produced by Nihon Nohyaku Co. Ltd.

REAR PANEL (cont.)

AgrEvo Pty. Ltd. A.C.N. 062 299 469
1731 - 1733 Malvern Road, Glen Iris, Vic. 3146
Telephone: (03) 9248 6666

THIS PRODUCT IS NOT CLASSIFIED AS A DANGEROUS
GOOD UNDER THE AUSTRALIAN CODE FOR THE
TRANSPORT OF DANGEROUS GOODS BY ROAD AND
RAIL.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE
FOR SPECIALIST ADVICE, CONTACT
UNITED TRANSPORT SERVICES EMERGENCY
RESPONSE COORDINATOR:
1800 024 973 (24 HOURS)

BAR CODE

Batch Number:
Date of Manufacture:

(1 L) 998362/0998
(2.5 L) 998234/0998
(5 L) 997765/0998

LABEL LEAFLET

Acaban[®]

MITICIDE

Active Constituent: 50 g/L FENPYROXIMATE
Pyrazole Miticide

DIRECTIONS FOR USE LEAFLET

To be read in conjunction with label affixed to container.

AgrEvo Pty. Ltd. A.C.N. 062 299 469
1731-1733 Malvern Road, Glen Iris, Vic. 3146
Telephone: (03) 9248 6666

998226/0998
NRA Approval No.: 45466/ 0898

LABEL LEAFLET (cont.)**DIRECTIONS FOR USE****Restraint**

DO NOT use in orchards where products likely to cause mite flaring have been used, or are planned to be used.

CROP	PEST	SITUATION	RATE	CRITICAL COMMENTS
			Dilute (high volume) per 100 L	
Apples and Pears	Two spotted mite (<i>Tetranychus urticae</i>) (Qld, NSW, Vic, SA, Tas, WA) European red mite (<i>Panonychus ulmi</i>) (Qld, NSW, Vic, SA, Tas)	IPM (where mite predators are present in significant numbers)	50 mL	Thorough coverage is essential. The minimum water volume recommended is 1,500 L/ha. See Application section in General Instructions for details. Use only one (1) application per season. See also Mite Resistance Guidelines in General Instructions.
		non IPM (where no mite predators or very low levels are present)	100 mL	Apples Apply before mite numbers exceed 8 to 10 mites/leaf or 70% of leaves are infested. Pears Apply when there are 1 to 2 mites/leaf or 25% of leaves are infested (whichever comes first).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

Apples, Pears - **DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

DO NOT GRAZE ANY TREATED AREA.

DO NOT FEED PRODUCE HARVESTED FROM TREATED AREAS TO LIVESTOCK, INCLUDING POULTRY.

LABEL LEAFLET (cont.)**GENERAL INSTRUCTIONS****Export of Treated Produce**

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Acaban Miticide. If you are growing produce for export, please check with AgrEvo Pty. Ltd. for the latest information on MRLs and import tolerances BEFORE using Acaban Miticide.

Application

Apply by high volume (dilute) sprayer to the point of run-off to ensure thorough coverage of foliage and fruit. The following spray volumes are offered as a guide to proper spraying of average sized trees at conventional planting intervals.

Stage	Dilute volume per ha (average trees)	Rate of Acaban/ha	
		at 50 mL/100 L	at 100 mL/100 L
Petal fall to first cover spray	1,500 to 2,500 L	750 to 1,250 mL	1,500 to 2,500 mL
Full foliage	2,500 to 3,500 L	1,250 to 1,750 mL	2,500 to 3,500 mL

Mixing

Shake product before opening as settling of contents may occur during storage. Partly fill tank with water and add the required amount of product to the spray tank while agitating. Continue agitation while spraying.

Compatibility

Acaban can be mixed with Insegar[®], Topas[®], Dipel[®], Systhane[®], Wuxal[®] Calcium, and some formulations of azinphos methyl, calcium chloride, calcium nitrate, chlorpyrifos, mancozeb and parathion. For compatible mixes please contact your reseller or AgrEvo representative.

Mite Resistance Guidelines

Acaban is a pyrazole miticide. To prevent the onset of resistance do not apply more than one spray of a pyrazole miticide per season. If more sprays are necessary use a miticide from a different chemical group.

Experience has shown that strains of mite resistant to pyrazole miticides have developed and that further strains may develop, resulting in reduced efficacy of these products. Since the occurrence of resistance cannot be foreseen, AgrEvo Pty. Ltd. accept no responsibility for any loss or damage resulting from the failure of Acaban to control resistant strains. In the event that Acaban is found by the buyer to be ineffective in controlling mites, or if the buyer has a reasonable suspicion that a tolerant or resistant strain of mite exists, then Acaban should not be used. At the first sign or appearance of any mite resistance to Acaban, the buyer should immediately contact a representative of AgrEvo Pty. Ltd. and should seek advice as to suitable alternative products.

Protection of Livestock

Low hazard to bees. No special precautions are required.

Protection of Wildlife, Fish, Crustaceans and Environment

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with product or used containers.

LABEL LEAFLET (cont.)**Drift Warning**

DO NOT apply in weather conditions, or from spraying equipment, which could be expected to cause spray to drift onto adjacent crops, crop lands, pasture, livestock, wetlands, waterbodies or watercourses.

Integrated Pest Management

Acaban is compatible with Integrated Pest Management orchards where mite predators such as *Typhlodromus pyri*, *T. occidentalis* or *Stethorus* are established. Depending on rate, levels of these mite predators may be suppressed for 1 to 3 weeks after application.

Storage and Disposal

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes. Avoid contact with eyes and skin. Avoid inhaling spray mist. When preparing spray and using the prepared spray wear elbow-length PVC gloves, goggles and half facepiece respirator. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and respirator (if rubber, wash with detergent and warm water).

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF.

Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet available from AgrEvo Pty. Ltd.

Re-entry Period

Do not allow entry into treated areas until the spray deposits have dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and elbow-length PVC gloves. Clothing must be laundered after each day's use.

Exclusion of Liability

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

LABEL LEAFLET (cont.)

For further information contact:

AgrEvo Pty. Ltd. A.C.N. 062 299 469

1731 - 1733 Malvern Road, Glen Iris, Vic. 3146

Telephone: (03) 9248 6666

Facsimile: (03) 9248 6650

or your local reseller or AgrEvo representative.

Acaban is produced by Nihon Nohyaku Co. Ltd.

Wuxal[®] is a Registered Trademark of Hoechst Schering AgrEvo GmbH

Acaban[®], Insegar[®] and Topas[®] are Registered Trademarks of Novartis AG, Basel, Switzerland

Dipel[®] is a Registered Trademark of Abbott Laboratories, North Chicago, Illinois, USA

Sythane[®] is a Registered Trademark of Rohm & Haas