

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

IMIDAN

INSECTICIDE / MITICIDE



RLP

Approved

ACTIVE CONSTITUENT: 150 g/L PHOSMET (An anti-cholinesterase compound)

SOLVENT: 568 g/L LIQUID HYDROCARBONS

GROUP

1B

INSECTICIDE

For the control of blue oat mite, red legged earth mite and lucerne flea as per the directions for use.

CONTENTS: 20 LITRES
: 110 LITRES

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie Qld 4172

Directions for use

Crop	Pest Controlled	State	Rate	Withholding Period	Critical Comments
Cereals	Red legged earth mite	All states	250 to 350 mL/ha	7 days (grazing)	Apply 3-5 weeks after autumn rains commence and when low temperatures favour pest development. Use the higher rate only when severe infestations are present. Further treatment may be necessary later in the season if re-infestation occurs.
Lucerne Pasture Seed crops	Red legged earth mite			2 days (grazing)	
	Lucerne flea	Vic., SA, WA, & Tas. only			
	Blue oat mite	Qld, NSW & WA only			Apply in autumn when populations indicate.
Pasture	Red legged earth mite	All states		2 days (grazing)	Apply 3-5 weeks after autumn rains commence and when low temperatures favour pest development. Use the higher rate only when severe infestations are present. Further treatment may be necessary later in the season if re-infestation occurs.
	Lucerne flea	N.S.W., Vic., S.A., W.A. & Tas. Only			Apply in autumn when populations indicate.
	Blue oat mite	Qld & WA only			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIOD

PASTURES - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 2 DAYS AFTER APPLICATION

CEREALS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Insecticide Resistance Warning

GROUP	1B	INSECTICIDE
-------	----	-------------

For insecticide resistance management Imidan Insecticide/Miticide is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to Imidan Insecticide/Miticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Imidan Insecticide/Miticide or other Group 1B insecticides are used repeatedly. The effectiveness of Imidan Insecticide/Miticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of Imidan Insecticide/Miticide to control resistant insects. Imidan Insecticide/Miticide may be subject to specific resistant management strategies. For further information contact your local supplier, Crop care representative or local agricultural department agronomist.

Application

Boomspray - apply in 50-100 L water/ha

Misting machine - apply in 20-40 L water/ha

Aerial application - apply in a minimum of 11 L water/ha

In low volume applications ensure water volume is sufficient to provide adequate coverage for control of target pests.

COMPATIBILITY

This product may be applied in combination with one of the following: 'Spray Seed', 'Gramoxone', 'Fusilade', 500 g/kg copper as copper oxychloride, non-ionic wetting agents, Urea (low biuret contents). A mixture with more than any one of the above products may be ineffective or cause serious damage. The use of such a mixture is therefore not recommended and entirely at the user's risk.

COMPATIBILITY with Selective Herbicides

This product may be used at up to 350 mL (55g active ingredient) per 80 litres water minimum per hectare in combination with: 500 g/L 2,4-D as amine-soft and hard water (1000 ppm); 800 g/L 2,4-D as ester; 500 g/L 2,4-D as ester; 400 g/L 2,4-DB as sodium-soft water only; 500 g/L 2,4-DB as amine; 500 g/L MCPB as amine; 242 g/L MCPA as sodium soft water only; 500 g/L MCPA as amine.

PROTECTION OF LIVESTOCK

Dangerous to bees. Do NOT spray any plants in flower while bees are foraging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container. Any spillage of concentrate should be immediately well covered with soil and placed in a waste dump approved for pesticide disposal.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Avoid spray drift from target area.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhale spray mist.

When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre Phone eg Australia 131 126; New Zealand 0800 764 766 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

Additional Information is listed in the Material Safety Data Sheet.

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatory imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA-WIDE
--	--

Gramoxone, Fusilade and Spray.Seed are Registered Trade Marks of Syngenta Limited.
Imidan is used under licence from the Trade Mark proprietor.



Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
Murarrie Qld 4172
Tel: (07) 3909 2000

Insert barcode,
label code and
drumMUSTER
logo

APVMA Approval No.: 46192/57588

BN :
DOM :

