



Product Name: OMITE 300W WETTABLE POWDER MITICIDE
APVMA Approval No: 48086/136569

Label Name:	OMITE 300W WETTABLE POWDER MITICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 300 g/kg PROPARGITE
Mode of Action:	GROUP 12C INSECTICIDE
Statement of Claims:	For the control of mites in crops as per the Directions of Use table
Net Contents:	1 kg, 8 kg, 10 kg, 100 g, 500 g, 1 kg (10 x 100 g), 1 kg (2 x 500 g), 5 kg (10 x 500 g), 6 kg (12 x 500 g), 1 kg, 1.3 kg, 10 kg (10 x 1 kg), 10.4 kg (8 x 1.3 kg)
Restrains:	RESTRAINTS DO NOT add wetting agents or crop oils to spray solution.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	WITHHOLDING PERIODS STRAWBERRIES: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION. APPLES, BANANAS, HOPS, PASSIONFRUIT, PEARS, STONEFRUIT AND VEGETABLES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Trade Advice:	
---------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS DO NOT apply using spraying equipment carried on the back of the user.</p> <p>MIXING Slowly add the required amount of product to the water under agitation in the spray vat.</p> <p>Once the water soluble bag(s) is added to the spray tank, ensure that thorough agitation of the spray mixture occurs. As a recommendation, the spray mixture should be agitated for 5 minutes prior to application, or until the water soluble bag is completely dissolved and the product is fully dispersed in the water.</p> <p>APPLICATION This product is not an ovicide. For fruit trees, two sprays 10 - 14 days apart are desirable for long term results. If lower volume sprays are applied, the 2 spray program is essential. Early applications are recommended. These should be made before the mite population is sufficient to cause serious damage. This product is non-systemic in action and high volume sprays and complete coverage are recommended.</p> <p>COMPATIBILITY A mixture of this product with any other product or more than one product may be ineffective or may cause serious damage. The use of such a mixture is, therefore, entirely at the user's risk. DO NOT add wetting agents or crop oils to spray solution as phytotoxicity may occur.</p>
-----------------------	---

Resistance Warning:	<p>INSECTICIDE RESISTANCE WARNING GROUP 12C INSECTICIDE</p> <p>For insecticide resistance management Omite® 300W is a Group 12C insecticide. Some naturally occurring insect biotypes resistant to Omite® 300W and other Group 12C insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Omite® 300W or other Group 12C insecticides are used repeatedly. The effectiveness of Omite® 300W on resistant individuals can be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, UPL Australia Pty Ltd accepts no liability for any losses that may result from the failure of Omite® 300W to control resistant insects. Omite® 300W may be subject to specific resistance management strategies. For further information contact your local supplier, UPL Australia Pty Ltd representative or local agricultural department agronomist.</p>
---------------------	---

Precautions:	<p>PRECAUTION RE-ENTRY: DO NOT allow entry into treated areas for low exposure activities such as irrigation, scouting or weeding for 21 days unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p> <p>DO NOT allow entry into treated areas for high exposure activities such as harvesting, pruning, training or tying for 32 days unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p>
--------------	--

Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the product or used containers.</p>
--------------	---

Storage and Disposal:	<p>STORAGE AND DISPOSAL</p> <p>Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.</p> <p>Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.</p>
-----------------------	---

Safety Directions:	<p>SAFETY DIRECTIONS</p> <p>Product is harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist.</p> <p>When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield, goggles and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	<p>FIRST AID</p> <p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting.</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

CROP	PEST CONTROLLED	STATES	RATE		WITHHOLDING Period	CRITICAL COMMENTS
			per 100L	per ha		
Stone fruit	Two spotted mite European red mite	All States Qld & WA only	200 g	5.5 kg	7 days	Apply as a high volume spray as soon as mites appear. Further applications may be necessary later in the season. Whatever volume is applied a minimum of 5.5 kg/ha is essential.
Apples	Two spotted mite European red mite	All States Vic, NSW, ACT, SA, WA & Qld only	200 g	5.5 kg 8.5 kg		
		Tas only	-	5.5 kg		
Apples and Stonefruit	Biological control of mites	All States	100 g	-		Apply where biological control is being practised. Complete spray coverage is essential for adequate control.
Pears	Two spotted mite	All States	200 g	-		Post-harvest spray only: Apply as high volume spray to achieve a minimum of 5.5kg/ha. This product may cause leaf spoiling and possible defoliation, this effect is more likely in hot conditions.
Strawberries	Two spotted mite (<i>Tetranychus urticae</i>) Biological control of mites	Vic, Tas, NSW, ACT, WA & Qld only	100 g	-		3 days

Table continued on next page

DIRECTIONS FOR USE *Continued*

CROP	PEST CONTROLLED	STATES	RATE		WITHHOLDING Period	CRITICAL COMMENTS
			per 100L	per ha		
Bananas	Two spotted mite (red spider)	Qld, NSW, ACT & WA only	100 g	1.25 kg	7 days	Spray when bunches covered and when pests appear and repeat as necessary. Avoid spraying under hot-humid-slow drying conditions when young bananas may be susceptible to insecticide damage.
Passionfruit	Passionvine mite (<i>Brevipalpus phoenicis</i>)	Qld & WA only	100 g	-		Apply as pest populations indicate. Do NOT apply more than twice per season.
Beans, Tomatoes, other vegetables	Two spotted mite	All States	100 g	-		Apply when pests appear and repeat as necessary. Good spray cover is necessary.
	Spider mites (<i>Tetranychus</i> spp.)	Qld & WA only				
Hops	Two spotted mite	Vic, WA & Tas only	200 g	-		Apply as a high volume spray on first appearance of mites. e.g. when growth is 2-3 metres high. Thoroughly spray all foliage with special attention to base of plants, repeat as necessary.
Ornamental plants	Two spotted mite European red mite	All States	100 g	-	-	Spray when mites appear and repeat when necessary.
	False spider mites (<i>Brevipalpus</i> spp)	Qld & WA only				Note: (1) This product may be quite phytotoxic to some ornamentals. (2) May be used in integrated mite control with the Chilean predatory mite <i>Phytoseiulus persimilis</i> for control of two spotted mite.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.