

STOLLER'S LIME SULPHUR

GENERAL INFORMATION:

STOLLER'S LIME SULPHUR Fungicide Insecticide is readily miscible in water and should be used on the same day as it is mixed.

MIXING & APPLICATION:

Shake contents well before using. Add the required amount of STOLLER'S LIME SULPHUR to the water in the spray tank with stirring or agitation. Agitate while spraying. Do not mix the STOLLER'S LIME SULPHUR with acidic water of pH below 4. Spray in the cool of the day.

COMPATIBILITY:

STOLLER'S LIME SULPHUR may be mixed with winter oil for use on Pome and Stone fruit during the dormant to semi-dormant stages only. For details of this mixture consult the label of winter oil. STOLLER'S LIME SULPHUR has limited compatibility with other pesticides and the use of any mixture is entirely at the user's risk.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:

Do not contaminate streams, rivers, drains or waterways with the chemical or used containers.

STORAGE AND DISPOSAL:

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Otherwise break, crush, puncture and bury empty containers in a local authority landfill. If not available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers should not be burnt.

SAFETY DIRECTIONS:

Avoid contact with eyes and skin. When preparing the spray wear elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

First Aid:

If poisoning occurs, contact a doctor or Poisons Information Centre.

If in eyes hold eyes open and flush with water continuously for 15 minutes. Seek medical attention. If on skin, wash thoroughly with soap and water.

MATERIAL SAFETY DATA SHEET:

For additional information, refer to MSDS No. S/LS200597

CONDITIONS OF SALE:

Seller warrants that this product consists of ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. Because the time, place, rate of application and other conditions of use are beyond seller's control, seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

APVMA Approval No: 49990/204/0707

Batch No:

Date of Manufacture:

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Stoller's

LIME SULPHUR



FUNGICIDE
INSECTICIDE

Active Constituent: 200g/L SULPHUR (S) as Polysulphide Sulphur

For the control of certain diseases, insects & mites on Citrus, Grapevines, Ornamentals, Pome & Stone Fruits & Vegetables.

As specified in the Directions for Use Table.

20 LITRES NET



S/LS 20-0597

Stoller Australia Pty. Ltd. A.C.N. 065 320 747

3 Todd Street Port Adelaide, South Australia 5015 Phone (08) 8440 2420

P.O. Box 420, Henley Beach, SA, 5022



490W220D3P

DIRECTIONS FOR USE:**RESTRAINTS:**

Do not apply during the growing period to Apricots and Peaches or other sulphur sensitive plants.
Do not apply when freezing weather is predicted, or the air temperature is over 32°C.
Do not apply within 2 weeks of an oil spray except as indicated under the compatibility heading.

| CROP | PEST / DISEASE | STATE | RATE per 100L of water | CRITICAL COMMENT |
|--|--|------------------------------------|------------------------------|---|
| CITRUS Grapefruit Oranges Lemons Mandarins | White Louse Scale (<i>Unaspis citri</i>) | NSW, Vic, SA & Tas only | 10L | Winter Spray: Apply in July and thoroughly wet all parts of the tree. |
| | | Qld only | 3.0-6.5L | |
| | Maori Mite (<i>Phyllocoptuuta oleivroa</i>) | NSW, Vic, WA, SA & Tas only | 2.5L | Summer Spray: Spray to thoroughly wet leaves. |
| GRAPEVINES | Bud Mite Erinose Mite (<i>Eriophyes vitis</i>) Rust Mite (<i>Calepitrimerus vitis</i>) Vine Bunch Mite (<i>Brevipalpus lewisi</i>) | Vic, WA, SA & Tas only | 10L | Apply as near as possible to budburst. NOTE: Spread of mites is checked by sulphur treatment used for powdery mildew. |
| | | Qld & NSW only | 5L | |
| | Post harvest treatment for mites | SA only | 2L | As soon as possible after harvest. |
| POME FRUIT Apples | Black Spot (Scab) (<i>Venturia inaequalis</i>) Bryobia Mite (<i>Bryobia rubrioculus</i>) Pearleaf Blister Mite (<i>Phytoptus pyri</i>) Powdery Mildew (<i>Podosphaera leucotricha</i>) | All States | 5L | Apply during dormant to late bud swell. NOTE: Lime Sulphur used after late green tip will restrict growth and depress yield. DO NOT use on Delicious or Cox's Orange Pinnin apples. |
| | San Jose Scale (<i>Comstockaspis perniciosus</i>) | Qld, NSW, SA, Vic & WA only | | |
| STONE FRUIT Almonds Apricots Peaches Plums Prunes Nectarines | Byrobia Mite (<i>Bryobia rubrioculus</i>) Freckle (<i>Venturia carpophila</i>) Leaf Curl (<i>Taphrina deformans</i>) Rust (<i>Tranzschelia discolor</i>) | All States | 5L | Spray while trees are dormant to the bud swell stage. |
| | Shot Hole (<i>Stigmina carpophila</i>) Brown Rot (Blossom Blight) (<i>Monilinia laxa</i>) San Jose Scale (<i>Comstockaspis perniciosus</i>) | Qld, NSW, SA, Vic & Tas only | | |
| | Frosted Scale (<i>Eulecanium prunosum</i>) | Qld, Vic, SA, WA & Tas only | | |
| | | | | |
| ORNAMENTALS Hedges Roses | Rust Two Spotted Mite (<i>Tetranychus urticae</i>) | All States | 1L | Apply as a Summer Spray. Spray to wet leaves and repeat as necessary. |
| | VEGETABLES Tomatoes | | | |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

In an Emergency
DIAL 000
Police or
Fire Brigade

**SPECIALIST ADVICE IN EMER-
GENCY ONLY**
Stoller (Australia) Pty Ltd
(08) 8440 2420

This product is not classified as
Dangerous Goods under the Australian
Code for the Transportation of
Dangerous Goods by road or rail

STOLLER'S LIME SULPHUR

GENERAL INFORMATION:

STOLLER'S LIME SULPHUR Fungicide Insecticide is readily miscible in water and should be used on the same day as it is mixed.

MIXING & APPLICATION:

Shake contents well before using. Add the required amount of STOLLER'S LIME SULPHUR to the water in the spray tank with stirring or agitation. Agitate while spraying. Do not mix the STOLLER'S LIME SULPHUR with acidic water of pH below 4. Spray in the cool of the day.

COMPATIBILITY:

STOLLER'S LIME SULPHUR may be mixed with winter oil for use on Pome and Stone fruit during the dormant to semi-dormant stages only. For details of this mixture consult the label of winter oil. STOLLER'S LIME SULPHUR has limited compatibility with other pesticides and the use of any mixture is entirely at the user's risk.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:

Do not contaminate streams, rivers, drains or waterways with the chemical or used containers.

STORAGE AND DISPOSAL:

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Otherwise break, crush, puncture and bury empty containers in a local authority landfill. If not available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers should not be burnt.

SAFETY DIRECTIONS:

Avoid contact with eyes and skin. When preparing the spray wear elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

First Aid:

If poisoning occurs, contact a doctor or Poisons Information Centre.

If in eyes hold eyes open and flush with water continuously for 15 minutes. Seek medical attention. If on skin, wash thoroughly with soap and water.

MATERIAL SAFETY DATA SHEET:

For additional information, refer to MSDS No. S/LS200597

CONDITIONS OF SALE:

Seller warrants that this product consists of ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. Because the time, place, rate of application and other conditions of use are beyond seller's control, seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

APVMA Approval No: 49990/10001/0707

Batch No:

Date of Manufacture:

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Infopest

Verified

Stoller's

LIME SULPHUR



FUNGICIDE INSECTICIDE

Active Constituent: 200g/L SULPHUR (S) as Polysulphide Sulphur

**For the control of certain diseases, insects & mites on Citrus,
Grapevines, Ornamentals, Pome & Stone Fruits & Vegetables.**

As specified in the Directions for Use Table.

1000 LITRES NET

Stoller Australia Pty. Ltd. A.C.N. 065 320 747
3 Todd Street Port Adelaide, South Australia 5015 Phone (08) 8440 2420
P.O. Box 420, Henley Beach, SA, 5022



S/LS 20-0597



490W220D3P

DIRECTIONS FOR USE:**RESTRAINTS:**

Do not apply during the growing period to Apricots and Peaches or other sulphur sensitive plants.

Do not apply when freezing weather is predicted, or the air temperature is over 32°C.

Do not apply within 2 weeks of an oil spray except as indicated under the compatibility heading.

| CROP | PEST / DISEASE | STATE | RATE per 100L of water | CRITICAL COMMENT |
|--|--|------------------------------------|------------------------------|---|
| CITRUS Grapefruit Oranges Lemons Mandarins | White Louse Scale (<i>Unaspis citri</i>) | NSW, Vic, SA & Tas only | 10L | Winter Spray: Apply in July and thoroughly wet all parts of the tree. |
| | | Qld only | 3.0-6.5L | |
| | Maori Mite (<i>Phyllocoptuuta oleivora</i>) | NSW, Vic, WA, SA & Tas only | 2.5L | Summer Spray: Spray to thoroughly wet leaves. |
| GRAPEVINES | Bud Mite | Vic, WA, SA & Tas only | 10L | Apply as near as possible to budburst. NOTE: Spread of mites is checked by sulphur treatment used for powdery mildew. |
| | Eriose Mite (<i>Eriophyes vitis</i>) Rust Mite (<i>Calepitrimerus vitis</i>) Vine Bunch Mite (<i>Brevipalpus lewisii</i>) | Qld & NSW only | 5L | |
| | Post harvest treatment for mites | SA only | 2L | As soon as possible after harvest. |
| POME FRUIT Apples | Black Spot (Scab) (<i>Venturia inaequalis</i>) Bryobia Mite (<i>Bryobia rubrioculua</i>) Pearleaf Blister Mite (<i>Phytoptus pyri</i>) Powdery Mildew (<i>Podosphaera leucotricha</i>) | All States | 5L | Apply during dormant to late bud swell. NOTE: Lime Sulphur used after late green tip will restrict growth and depress yield. DO NOT use on Delicious or Cox's Orange Pinnin apples. |
| | San Jose Scale (<i>Comstockaspis perniciosus</i>) | Qld, NSW, SA, Vic & WA only | | |
| STONE FRUIT Almonds Apricots Peaches Plums Prunes Nectarines | Byrobia Mite (<i>Bryobia rubrioculua</i>) Freckle (<i>Venturia carpophila</i>) Leaf Curl (<i>Taphrina deformans</i>) Rust (<i>Tranzschella discolor</i>) | All States | 5L | Spray while trees are dormant to the bud swell stage. |
| | Shot Hole (<i>Stigmia carpophila</i>) Brown Rot (Blossom Blight) (<i>Monilinia laxa</i>) San Jose Scale (<i>Comstockaspis perniciosus</i>) | Qld, NSW, SA, Vic & Tas only | | |
| | Frosted Scale (<i>Eulecanium prunosum</i>) | Qld, Vic, SA, WA & Tas only | | |
| ORNAMENTALS Hedges Roses | Rust Two Spotted Mite (<i>Tetranychus urticae</i>) | All States | 1L | Apply as a Summer Spray. Spray to wet leaves and repeat as necessary. |
| VEGETABLES Tomatoes | Powdery Mildew Tomato Mite (<i>Aculops lycopersici</i>) | | | |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

In an Emergency
DIAL 000
Police or
Fire Brigade

**SPECIALIST ADVICE IN EMER-
GENCY ONLY**
Stoller (Australia) Pty Ltd
(08) 8440 2420

This product is not classified as
Dangerous Goods under the Australian
Code for the Transportation of
Dangerous Goods by road or rail