

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SIPCAM PACIFIC

AMESIP 500 FLOWABLE HERBICIDE

ACTIVE CONSTITUENT: 500 g/L AMETRIN

GROUP C HERBICIDE

For the control of summer grasses and broadleaf weeds in pineapples and sugar cane and for use in industrial situations as per the directions for use table.

50161/0897

N.R.A.

APPROVED LABEL

1, 5, + 20L

Date

Max Dorell

SIPCAM PACIFIC AUSTRALIA PTY LIMITED

40 WENTWORTH AVENUE

SYDNEY NSW 2000

ACN 073 176 888

EMERGENCY CONTACT NUMBER 0299602875

*Page 1 of 7
21/8/97*

DIRECTIONS FOR USE:

Restrains: DO NOT use in high winds
 DO NOT allow spray to drift
 DO NOT spray if conditions are very dry.
DO NOT re-plant pineapples or sugar cane for 8 months after last application

*Page 2 of 7
21/8/12*

Crop/ Situation	Weeds Controlled	State	Rate				WHP days	Critical Comments
			Boom Spray per ha	Hand Spray per 100L water	Knapsack per 15L water	Misting per L		
Pineapples	Crowsfoot grass, pretty boy, Sedges, Summer Grass, Thickhead	Qld,NSW 4 only	to 8L in 2000 L water	-			98	Apply when soil moisture is good and when weeds are actively growing. Will provide both knockdown and residual action. Use the lowest rate for broadleaf weeds up to 15cm high and for short residual control. Use the higher rate for broadleaf weeds up to 30cm high and seedling grasses and where moderate residual control is required. Rates of more than 4L/ha should not be applied at less than 4 month intervals. Do not apply more than a total of 20L/ha per plant crop.
Sugar Cane	Barnyard grass, Bell vine, Billy-goat Weed, Chinese burr, Crowsfoot Grass. Flannel Weed, Gambia Pea, Rattlepod, Summer Grass	Qld, NSW, WA only	4.5L in 2000L water + 250mL 100% non-ionic surfactant	400mL + 125mL 100% non-ionic surfactant (Apply at least 1100 mixture/ha)	60mL + 20mL 100% non-ionic surfactant (Apply at least 1100 mixture/ha)	40mL + 10mL 100% non-ionic surfactant (Apply at least 1100 mixture ha)	-	Ensure thorough coverage. Apply to broadleaf weeds up to 38cm high and grasses up to 30 cm high. Avoid spraying onto leaves of sugar cane plant. NOTE: Slight decreases in crop vigour from post-emergence application may result and will last for 4 weeks.

Page 3 of 7
21/8/17

<p>Sugar cane Plant and Ratoon)</p>	<p>African stargrass, Awnless Barnyard grass, Bellvine, Billy-goat weed, Chinese burr, Crowsfoot grass, Cupids Flower, Flannel weed, Gambia Pea, Green summer grass, Mossman River grass, Rattlepod, summer grass</p>	<p>Qld only</p>	<p>4L + 6L Atrazine 500 Flowable in sufficient water for thorough coverage.</p>	<p>-</p>				<p>Apply pre-emergent or early post emergent to both crop and weeds. If weeds are present they should be actively growing and not beyond the 3-4 leaf stage. Best results are achieved if application is made to moist soil and when rain is likely to occur within 10 days. Apply as either a band spray or as a blanket spray. If the cane has emerged, apply as a directed spray to minimise contact with cane plants. NOTE: Some phytotoxicity may occur on variety Q90.</p>
<p>Rights of way, Roadsides, Drains, Fallow croplands, Footpaths, Industrial Areas, Railway Lines, Firebreaks, Headlands etc</p>	<p>Bathurst Burr</p>	<p>All States</p>	<p>-</p>	<p>550mL + 140mL 100% non-ionic surfactant</p>	<p>75mL + 20mL 100% non-ionic surfactant</p>	<p>-</p>		<p>Apply to young actively growing plants. Do not spray water stressed plants or plants that are more than 60 cm high. Ensure complete coverage of leaves and stems to ground level. Spray plants to point of run off.</p>
	<p>California Burr, Noogoora Burr</p>		<p>-</p>	<p>720mL + 140mL 100% non-ionic surfactant</p>	<p>100mL + 20mL 100% non-ionic surfactant</p>			
	<p>Caltrop (Puncture vine)</p>			<p>360mL + 140mL 100% non-ionic surfactant</p>	<p>50mL + 20mL 100% non-ionic surfactant</p>			

Page 4 of 7
2/18/97

Weeds and Grasses such as Annual Love grasses, Barnyard grass, Billy-goat weed, Cobbler's Peg, Couch Grass, Crowsfoot Grass, Flannel Weed, Fleabane, Gambia Pea, Guinea Grass, Paspalum, Plantain, Rattlepod, Rhodes Grass, Sida retusa, Stinking Roger, summer grass, Tridax Daisy				As an additive to recommended rates of Weedazol' Total or Weedazol' Industrial Total. Add Sipcam Amesip 500 Flowable at the following rates:		-	As an additive to knockdown herbicides, improved grass and weed control is obtained.
				270mL + 140mL 100% non-ionic surfactant (mm 2200L spray/ha)	100mL + 20 mL 100% non-ionic surfactant (min 80L spray/ 1000 square metres)	20mL + 1 mL 100% non-ionic surfactant (min 10L spray per 1000 square metres)	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: PINEAPPLES - DO NOT APPLY WITHIN 98 DAYS OF HARVEST

GENERAL INSTRUCTIONS

Sipcam Amesip 500 Flowable Herbicide is a liquid pre and post emergent selective herbicide which provides knockdown and residual control of annual grasses and broadleaf weeds. It is a systemic herbicide which is absorbed by the leaves and roots of target weeds.

Mixing

As settling may occur during storage, stir the product or invert the container prior to use. Pour the product into the spray vat through a sieve to remove any dry particles or flakes which may occur under hot storage conditions.

Partially fill the spray tank with water and maintain agitation while adding the required quantity of **Sipcam Amesip 500 Flowable** Herbicide. Agitation should continue after filling and during spraying.

If spray is allowed to stand, agitate vigorously from the bottom before resuming spraying.

Application

Application may be made in two ways:

1) Post-emergence to crops and weeds:

Seedling grass and broadleaf weeds should be at the 2 to 3 leaf stage and actively growing. Always add, a non-ionic (100%) surfactant at the rate of **200mL** per 1 OOL mixture.

2) Post-emergence to Crops and Pre-emergence to weeds:

The product may be applied before the weeds emerge but it is preferable to wait until weeds have emerged before spraying.

Method of Application

The following minimum spray volumes per hectare are suggested:

Method of Application	Water Volume
High volume boom	2,000L
Medium volume boom	1,100L
Low volume boom	100 to 300L
Knapsack	1,100L
Misting machine	110L
Aerial Spray	30 to 50L

Weed Resistance Warning

Sipcam Amesip Flowable Herbicide is a member of the **triazines** group of herbicides. **Sipcam Amesip Flowable** Herbicide has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management **Sipcam Amesip Flowable** Herbicide is a Group C Herbicide. Some **naturally occurring** weed biotypes resistant to **Sipcam Amesip 500 Flowable** Herbicide and other triazine herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used

Page 5 of 7
21/8/97

repeatedly. These resistant weeds will not be controlled by **Sipcam** Amesip 500 Flowable Herbicide or any other **triazine** herbicide.

Since the occurrence of resistant weeds is difficult to detect prior to use, **Sipcam** Pacific Australia Pty Limited accepts no liability for any losses that may result **from** the failure of **Sipcam** Amesip 500 Flowable Herbicide to control resistant weeds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Do not contaminate streams, rivers or waterways with the chemical or used containers.
Dangerous to fish.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do not apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

STORAGE AND DISPOSAL

Store in the closed original container, in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to **recycler** or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
For **further** information refer to Material Safety Data Sheet.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between **Sipcam** Pacific Pty Limited and the buyer but not limited to the following.

To the extent permitted by law **all** conditions and other warranties or other rights of action which buyer or any other user may have against **Sipcam** Pacific Pty Limited (**Sipcam**) or Seller are hereby excluded. **Sipcam** hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising **from** reliance on product information or advice provided by **Sipcam** or on its behalf unless it can be established that such information

or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice **Sipcam** will not be held liable for any loss, injury or damage indirect or consequential **arising** from the sale, supply, use or application of this product. **Sipcam's** liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

NRA Approval Number: 50161/0897

Date of Manufacture:

Batch No:

¹Registered Trade Mark of Union Carbide

Page 787
21/8/97