

KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Kendon

N.A.A. STOP DROP

Active Constituent: 20g/L N.A.A.

FRUIT THINNING AND SETTING COMPOUND

To assist in the prevention of pre-harvest drop of Apples and Pears. May also be used as a thinning spray for Apples, and for striking hard and softwood cuttings.

BATCH NO:

NET: 1L, 5L & 20L

KENDON CHEMICAL & MANUFACTURING CO. PTY. LTD.

71 McCLURE STREET, THORNBURY, VIC. 3071

NRA APPROVAL NO: 50357/095

Directions for Use:

CROP	ACTION	RATE	CRITICAL COMMENTS
Apples and Pears	Assist prevention of pre-harvest fruit drop	1L/2000L of water (10 ppm) 500 ml/2000L of water (5-ppm)	Apply as a high volume spray 12-14 days before harvest to thoroughly wet fruit and foliage. A second, half strength spray (5 p.p.m.) may be necessary if conditions favour excessive fruit drop.
Apples	Fruit thinning	250-1000mL per 2000L of water. (2:5 - 10 ppm)	Apply as a high volume spray 10-14 days after full bloom to the top two thirds of tree for uniform thinning. For specific varieties, consult local Dept. of Agriculture, in Tasmania, apply, particularly to Golden Delicious, from balloon blossom to 7 days after full bloom. Application rates for mature trees should be a minimum of 4000 L/ha.
Hard or Soft-wood cuttings	Root striking	12.5mL to 50mL per 10L of water.	Dip base of cutting to depth of 5mm into N.A.A. solution for approx. 5 seconds. Allow to dry and insert half the length of the cutting into striking medium. Use the stronger solution for hardwood cuttings, weaker solution for softwood cuttings.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION WITHHOLDING PERIOD: Do not apply later than 1 day before harvest.

General Instructions: When spraying for the prevention of pre-harvest fruit drop, do not allow fruit to remain on trees beyond normal harvest date otherwise fruit may become over-mature.

Use a suitable weter, such as Tween 20, at 125mL/100L of spray volume.

N.A.A. must be absorbed into the tree to be effective. Therefore slow drying conditions after spraying, high volume spraying and the use of a wetting agent will enhance effectiveness.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available,

bury the containers below 500mm in a disposal pit, specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre, phone 131126.

If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Conditions of Sale: This product as supplied is of high grade and believed to be suitable for any purpose for which it is expressly supplied and must be used in accordance with the directions for use given on this label. No responsibility is accepted in respect of this product, save those non-excludable conditions implied by the Trade Practices Act or any applicable State Legislation.

N.R.A. 
 APPROVED LABEL

Date: 8/11/98