

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 1 of 11

**7 & 15 kg label**

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**VITAVAX<sup>®</sup> 750C**  
**SEED TREATMENT**



**ACTIVE CONSTITUENTS:**  
**750 g/kg CARBOXIN**  
**6.4 g/kg CYPERMETHRIN** (isomer ratio 45:55)

GROUP	<b>G</b>	FUNGICIDE
-------	----------	-----------

For the treatment of wheat, oats, barley and triticale  
as specified in the Directions for Use Table

Net Contents 7 & 15 kg

(Crompton logo)

Crompton Specialties Pty Ltd  
3 Opala Street Regency Park SA 5010  
Tel: (08) 8268 5700 Fax: (08) 8268 7555

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 2 of 11

### 7 & 15 kg label

#### **DIRECTIONS FOR USE:**

#### **RESTRAINTS:**

Only treat good quality sound seed as treatment of mechanically damaged or poor quality seed may result in reduced germination and/or reduced seedling vigour and emergence.

CROP	DISEASE / PEST CONTROLLED	STATES	RATE /100 kg SEED	CRITICAL COMMENTS
Wheat	Loose smut ( <i>Ustilago tritici</i> )	All States	125 g	Mix thoroughly and ensure that the recommended rate is applied.
	Bunt ( <i>Tilletia laevis</i> , <i>T. tritici</i> )		70 g	
	Seedborne flag smut ( <i>Urocystis agropyri</i> )	WA only	125 g	
Barley	Soilborne flag smut ( <i>Urocystis agropyri</i> )		125 g	
	Loose smut ( <i>Ustilago tritici</i> )	All States	125 g	
Oats	Covered smut ( <i>Ustilago segetum</i> var <i>hordei</i> )		70 g	
	Smut ( <i>Ustilago segetum</i> var <i>hordei</i> )	All States	70 g	
Triticale	Smut ( <i>Ustilago avenae</i> )		70 g	
	Loose smut ( <i>Ustilago tritici</i> )	Vic, NSW, ACT, WA, Qld & TAS only	125 g	
	Seedborne flag smut ( <i>Urocystis agropyri</i> )	Vic, NSW, ACT, WA, Qld & TAS only	70 g	
Wheat Barley Oats Triticale	Protection against insect pests of stored grains: Granary weevil ( <i>Sitophilus granarius</i> ) Indian meal moth ( <i>Plodia interpunctella</i> ) Lesser grain borer ( <i>Rhyzopertha dominica</i> ) Rice weevil ( <i>Sitophilus oryzae</i> ) Rust-red flour beetle ( <i>Tribolium castaneum</i> ) Sawtoothed grain beetle ( <i>Oryzaephilus surinamensis</i> ) Tropical warehouse moth ( <i>Ephesia cautella</i> )	All States	70 g or 125 g	Both rates give protection against insect pests. Use rate appropriate to disease control required.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD:**

**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION.**

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 3 of 11

**7 & 15 kg label**

**CONDITIONS OF SALE:**

Crompton Specialties Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crompton Specialties Pty Ltd is limited to the replacement of the goods or (at the option of Crompton Specialties Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crompton Specialties Pty Ltd within thirty days of delivery.

® Vitavax is a registered trademark of Crompton Corporation, USA.

In an Emergency Dial <b>000</b> Police or Fire Brigade	<b>SPECIALIST ADVICE IN EMERGENCY  ONLY</b> Crompton Specialties Pty Ltd <b>1800 033 111</b> ALL HOURS - AUSTRALIA WIDE
---	--

BN  
DOM

Printer Code

NRA Approval No. 51415/0703

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 4 of 11

### **7 & 15 kg label**

## **GENERAL INSTRUCTIONS:**

Vitavax 750C Seed Treatment is a dry seed dressing for treatment of grain by standard methods. This product contains Cypermethrin for protection against insect pests of stored grain.

**Seed Quality:** Seed treatment should not be used on seed with more than 12% moisture content, or on sprung, damaged or severely pinched seed, or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard.

**Storage of Treated Seed:** If seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place. Although Vitavax 750C Seed Treatment has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for performance of stored treated seed.

### **Fungicide Resistance Warning:**

GROUP	<b>G</b>	FUNGICIDE
-------	----------	-----------

Vitavax 750C Seed Treatment is a member of the Oxathiin group of fungicides. For fungicide resistance management, Vitavax 750C Seed Treatment is a Group G fungicide.

Some naturally occurring individual fungi resistant to Vitavax 750C Seed Treatment and other Group G fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Vitavax 750C Seed Treatment and other Group G fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crompton Specialties Pty Ltd accepts no liability for any losses that may result from the failure of Vitavax 750C Seed Treatment to control resistant fungi.

## **PRECAUTIONS:**

Do not use treated seed for animal or human consumption. Do not allow treated seed to contaminate grain or other seed intended for animal or human consumption. Do not feed treated seed, or otherwise expose, to wild or domestic birds.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:**

Treated seed should not be held over from one sowing season for sowing in subsequent seasons.

## **PROTECTION OF LIVESTOCK:**

DO NOT feed treated seed to animals, including poultry. It is important that any treated seed not used for sowing is destroyed. Bags which have held treated seed should not be used for any other purpose. When treated seed is stored, it should be kept apart from other grain and the bags (or other containers) should be clearly marked to indicate the contents have been treated with Vitavax 750C Seed Treatment.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers or with bags which have held treated seed.

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 5 of 11

### **7 & 15 kg label**

#### **STORAGE AND DISPOSAL:**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight and away from foodstuffs, drink containers and livestock. Do not use container for any other purpose. Shake empty bag into spray tank. Do not dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### **SAFETY DIRECTIONS:**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling dust. When using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield and disposable face mask. Repeated exposure may cause allergic disorders. Facial skin contact may cause temporary facial numbness. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

#### **FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131 126.*

#### **MATERIAL SAFETY DATA SHEET:**

For additional information, refer to the Material Safety Data Sheet.

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 6 of 11

Measure pack 7 kg water soluble label

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **VITAVAX<sup>®</sup> 750C**

## **SEED TREATMENT**

**ACTIVE CONSTITUENTS:**  
**750 g/kg CARBOXIN**  
**6.4 g/kg CYPERMETHRIN** (isomer ratio 45:55)

**WATER SOLUBLE PACKAGING. KEEP DRY.**  
**BEFORE USE READ ALL DIRECTIONS ON THE OUTER PACK.**  
**USE CONTENTS IMMEDIATELY AFTER OPENING PLASTIC BAG.**

**Not to be sold separately.**

**Net Contents 700 g**

(Crompton logo)

Crompton Specialties Pty Ltd  
3 Opala Street Regency Park SA 5010  
Tel: (08) 8268 5700 Fax: (08) 8268 7555

NRA Approval No. 51415/0703

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 7 of 11

*Outer Pack 7 kg water soluble label*

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **VITAVAX<sup>®</sup> 750C**

## **SEED TREATMENT**

**ACTIVE CONSTITUENTS:**  
**750 g/kg CARBOXIN**  
**6.4 g/kg CYPERMETHRIN** (isomer ratio 45:55)

GROUP	<b>G</b>	FUNGICIDE
-------	----------	-----------

For the treatment of wheat, oats, barley and triticales  
as specified in the Directions for Use Table

Net Contents 7 kg

**Contains 10 x 700 g water soluble packs which it is illegal to sell separately.**

**DO NOT DESTROY BOX WHILE PRODUCT STILL REMAINS.**

(Crompton logo)

Crompton Specialties Pty Ltd  
3 Opala Street Regency Park SA 5010  
Tel: (08) 8268 5700 Fax: (08) 8268 7555

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 8 of 11

Outer Pack 7 kg water soluble label

**DIRECTIONS FOR USE:**

**RESTRAINTS:**

Only treat good quality sound seed as treatment of mechanically damaged or poor quality seed may result in reduced germination and/or reduced seedling vigour and emergence.

CROP	DISEASE / PEST CONTROLLED	STATES	RATE /100 kg SEED	CRITICAL COMMENTS
Wheat	Loose smut ( <i>Ustilago tritici</i> )	All States	125 g	Mix thoroughly and ensure that the recommended rate is applied.
	Bunt ( <i>Tilletia laevis</i> , <i>T. tritici</i> )		70 g	
	Seedborne flag smut ( <i>Urocystis agropyri</i> )	WA only	125 g	
Barley	Soilborne flag smut ( <i>Urocystis agropyri</i> )		125 g	
	Loose smut ( <i>Ustilago tritici</i> )	All States	70 g	
Oats	Covered smut ( <i>Ustilago segetum</i> var <i>hordei</i> )		70 g	
	Smut ( <i>Ustilago segetum</i> var <i>hordei</i> )	All States	70 g	
Triticale	Smut ( <i>Ustilago avenae</i> )		70 g	
	Loose smut ( <i>Ustilago tritici</i> )	Vic, NSW, ACT, WA, Qld & TAS only	125 g	
	Seedborne flag smut ( <i>Urocystis agropyri</i> )	Vic, NSW, ACT, WA, Qld & TAS only	70 g	
Wheat Barley Oats Triticale	Protection against insect pests of stored grains: Granary weevil ( <i>Sitophilus granarius</i> ) Indian meal moth ( <i>Plodia interpunctella</i> ) Lesser grain borer ( <i>Rhyzopertha dominica</i> ) Rice weevil ( <i>Sitophilus oryzae</i> ) Rust-red flour beetle ( <i>Tribolium castaneum</i> ) Sawtoothed grain beetle ( <i>Oryzaephilus surinamensis</i> ) Tropical warehouse moth ( <i>Ephestia cautella</i> )	All States	70 g or 125 g	Both rates give protection against insect pests. Use rate appropriate to disease control required.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD:**

**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION.**



Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 9 of 11

Outer Pack 7 kg water soluble label

**CONDITIONS OF SALE:**

Crompton Specialties Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crompton Specialties Pty Ltd is limited to the replacement of the goods or (at the option of Crompton Specialties Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crompton Specialties Pty Ltd within thirty days of delivery.

® Vitavax is the Registered Trademark of Crompton Corporation, USA

In an Emergency Dial <b>000</b> Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY Crompton Specialties Pty Ltd <b>1800 033 111</b> ALL HOURS - AUSTRALIA WIDE
---	---

BN  
DOM

Printer Code

NRA Approval No. 51415/0703

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 10 of 11

### Outer Pack 7 kg water soluble label

#### **GENERAL INSTRUCTIONS:**

Vitavax 750C Seed Treatment is a dry seed dressing for treatment of grain by standard methods. This product contains Cypermethrin for protection against insect pests of stored grain.

**Mixing & Application:** Add Vitavax in water soluble bags to agitating tank containing sufficient water to form a slurry. Use only through equipment designed to apply slurry seed treatments. Ensure the packs are completely dissolved prior to treating of seed. Ensure that the recommended rate is uniformly applied to the seed.

**Seed Quality:** Seed treatment should not be used on seed with more than 12% moisture content, or on sprung, damaged or severely pinched seed, or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard.

**Storage of Treated Seed:** If seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place. Although Vitavax 750C Seed Treatment has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for performance of stored treated seed.

#### **Fungicide Resistance Warning:**

GROUP	<b>G</b>	FUNGICIDE
-------	----------	-----------

Vitavax 750C Seed Treatment is a member of the Oxathiin group of fungicides. For fungicide resistance management, Vitavax 750C Seed Treatment is a Group G fungicide.

Some naturally occurring individual fungi resistant to Vitavax 750C Seed Treatment and other Group G fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Vitavax 750C Seed Treatment and other Group G fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crompton Specialties Pty Ltd accepts no liability for any losses that may result from the failure of Vitavax 750C Seed Treatment to control resistant fungi.

#### **PRECAUTIONS:**

Do not use treated seed for animal or human consumption. Do not allow treated seed to contaminate grain or other seed intended for animal or human consumption. Do not feed treated seed, or otherwise expose, to wild or domestic birds.

#### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:**

Treated seed should not be held over from one sowing season for sowing in subsequent seasons.

#### **PROTECTION OF LIVESTOCK:**

DO NOT feed treated seed to animals, including poultry. It is important that any treated seed not used for sowing is destroyed. Bags which have held treated seed should not be used for any other purpose. When treated seed is stored, it should be kept apart from other grain and the bags (or other containers) should be clearly marked to indicate the contents have been treated with Vitavax 750C Seed Treatment.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers or with bags which have held treated seed.

Vitavax 750C Seed Treatment	Approved 0703
Company name change, 30-6-03	Page 11 of 11

Outer Pack 7 kg water soluble label

**STORAGE AND DISPOSAL:**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight and away from foodstuffs, drink containers and livestock. DO NOT repack or use container for any other purpose.

Shake empty bag into spray tank. Do not dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling dust. When using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield and disposable face mask. Repeated exposure may cause allergic disorders. Facial skin contact may cause temporary facial numbness. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

**FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.*

**MATERIAL SAFETY DATA SHEET:**

For additional information, refer to the Material Safety Data Sheet.