

Product Name: TERRAZOLE 350 WP SOIL FUNGICIDE
APVMA Approval No.: 53332/136568



Label Name:	TERRAZOLE 350 WP SOIL FUNGICIDE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 350 g/kg ETRIDIAZOLE
Mode of Action:	GROUP 14 FUNGICIDE
Statement of Claims:	Controls Damping Off, Root & Stem Rot of Ornamentals & Turf caused by Phytophthora & Pythium Fungi as specified in the Directions For Use table
Net Contents:	10kg pail 10kg carton 2kg carton 2kg pail
Restrains:	
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

Trade Advice:	
---------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS</p> <p>MIXING Add the required quantity to the water in the spray tank with agitation. Spray solutions need to be continuously agitated after initial mixing. Water or irrigate after application to improve penetration of the fungicide into the soil.</p> <p>APPLICATION Use only at recommended rates, as overdosage may be harmful to sensitive plants. A test for compatibility with other chemicals is advisable before use.</p>
-----------------------	--

Resistance Warning:	<p>FUNGICIDE RESISTANCE WARNING GROUP 14 FUNGICIDE For fungicide resistance management, Terrazole 350 WP Soil Fungicide is a Group 14 fungicide. Some naturally occurring individual fungi resistant to Terrazole 350 WP Soil Fungicide and other Group 14 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal populations if these fungicides are used repeatedly. These resistant fungi will not be controlled by Terrazole 350 WP Soil Fungicide or other Group 14 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, UPL Australia Pty Ltd accepts no liability for any losses that may result from the failure of Terrazole 350 WP Soil Fungicide to control resistant fungi.</p>
---------------------	--

Precautions:	
--------------	--

Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT plant food crops in rotation on treated soils.</p> <p>PROTECTION OF LIVESTOCK DO NOT allow stock to graze in any treated area. DO NOT feed turf/lawn clippings from any treated area to poultry or livestock.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
--------------	--

Storage and Disposal:	<p>STORAGE AND DISPOSAL Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding</p>
-----------------------	--

it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

Safety Directions:

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

First Aid Instructions:

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 13 11 26, New Zealand 0800 764 766.

First Aid Warnings:

DIRECTIONS FOR USE

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
BEDDING PLANTS (ornamental seedlings)	Root Rot, Stem Rot (<i>Phytophthora</i> spp) Damping Off (<i>Pythium</i> spp)	ALL STATES	60-100 g in 250 L water to 40 m ²	Apply as a drench. Repeat applications may be made at 4 - 6 week intervals if required. Use higher rate on soils high in organic matter.
HERBACEOUS AND WOODY ORNAMENTAL PLANTS			60-100 g in 100 L water to 10 m ²	Repeat applications at 2-3 month intervals to maintain disease control.
BEDS POTS			25-75 g in 100 L water	Apply 300 mL of solution per 15 cm pot or sufficient solution to saturate the soil in the pot. Use higher rate for soils with high organic matter or peat.
TURF ESTABLISHED	Damping Off Grease Spot (<i>Pythium</i> spp)	NSW, ACT, Vic, SA and WA	100-250 g in 25 L water to 100 m ²	Apply uniformly as a spray. Repeat treatment at 5-10 days to maintain disease control.
TURF NEWLY SEEDED			150-200 g in 25-50 L water to 100 m ²	Apply as a spray immediately after seeding. Repeat application at 5-7 day intervals to maintain disease control.
DRY SOIL MIX (seed box and potting mixes)	Damping Off (<i>Pythium</i> spp)	ALL STATES	45-90 g per m ³ of soil	Apply as dust and mix into one cubic metre of soil during soil fertilizer blending operation.
ROOT DIP	Root infections (<i>Phytophthora</i> spp)	Qld, NSW, ACT, WA, Tas and SA	60-70 g in 50 L water	Mix well and stir frequently to maintain adequate suspension during dipping operation.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.