

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FLUOCAMTM 500 HERBICIDE

ACTIVE CONSTITUENT: 500g/L FLUOMETURON

GROUP	C	HERBICIDE
--------------	----------	------------------

For control of many broadleaf weeds and annual grasses in cotton.

20, 200 LITRES

**SIPCAM PACIFIC AUSTRALIA PTY LTD
Suite 11, 23-31 Gheringhap Street
GEELONG VIC 3220
ABN 94 073 176 888
TEL: 03 5223 33746**

TM Trademark of Sipcam Pacific Australia Pty Ltd

DIRECTIONS FOR USE – NSW AND QLD ONLY

RESTRAINTS- for use in cotton only for the control of those weeds listed in the General Instructions
 Severe plant injury may result if heavy rain occurs between sowing and emergence.
 DO NOT make more than one post-emergent application per season.

Method of Application	Rate per hectare	Critical Comments
Soil incorporated pre-planting treatment	2.8 to 5.6 litres	Spray as an overall treatment. Pre-irrigate then apply just prior to incorporation (higher rates for heavy high organic soils, lower rates for lighter low organic soils). Incorporation: Lillistone cultivators are ideal for incorporation and should be utilised wherever available. Regardless of machinery used, it is essential that incorporation depth of seed placement be closely governed and not allowed to exceed the depth of seed placement (approximately 5 cm). Scrape off the top of the hill at planting time. Apply a band immediately after planting, to replace Fluocam which has been scraped off into the furrow. Use lower rate for lighter soils, the higher for heavier soils.
At seeding	1.8 to 3.6 litres	
Pre-emergence Post-sowing treatment	4.5 to 7.2 litres per sprayed hectare	Apply immediately after sowing as an overall or band treatment. Use higher rates for heavier organic soils, lower rates for lighter low organic contents soils. If band treatments are used, band width should be a minimum of 40 cm. Adjustment in application rate is necessary.
Post-emergence		
Early spraying	1.3 to 2.8 litres mixed in 220 to 440 litres of water	Fluocam may be applied as a post-emergent application to control broadleaf weeds which escaped the pre-emergent application or to control broadleaf weeds in situations where pre-emergent applications have not been previously made. Apply as a directed spray when cotton is 15 cm high. The weeds should be actively growing and not be more than 5 cm high. Add a non-ionic surfactant.
Lay-by spraying	2.8 to 5.6 litres mixed in 220 to 440 litres of water	Apply as a directed spray after the last cultivation. The weeds should be actively growing and not more than 8 cm high. Add a non-ionic surfactant.

**NOT TO BE USED FOR ANY PURPOSE, OR ANY MANNER, CONTRARY TO THIS LABEL
 UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:
 DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION**

GENERAL INSTRUCTIONS

Fluocam 500 Herbicide may be applied in several ways and growers should select the method most useful in their particular management programme.

RESISTANT WEEDS WARNING

Fluocam 500 Herbicide is a member of the urea group of herbicides and has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management Fluocam 500 Herbicide is a Group C herbicide.

Some naturally-occurring weed biotypes resistant to Fluocam 500 Herbicide and other urea herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Fluocam 500 Herbicide or other urea herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Fluocam 500 Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Sipcam representative.

Mixing

Settling may occur after storage for some weeks. Stir product or invert container several times before opening. Pour the flowable into the spray vat through a strainer to remove any dry particles or flakes, which can occasionally occur under hot storage conditions. Add the full quantity of the product to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing. Continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand.

Reseal part used container immediately.

Application

Fluocam 500 Herbicide should be applied to bare soil before weeds emerge. In irrigated areas irrigation should precede or immediately follow spraying. The product should be applied in a minimum of 220 litres of water per hectare sprayed.

Weeds Controlled

Fluocam 500 Herbicide controls weeds and grasses including Barley Grass, Barnyard Grass, Bathurst Burr, Bellvine, Bladder Kemia, Boggabri Weed, Capeweed, Common Sowthistle, Common Thornapple, Fat-hen, Liverseed Grass, Mossman River Grass, Nightshade, Noogoora Burr, Paddy Melon, Pigweed, Paterson's Curse, Prince-of-Wales Feather, Ryegrass, Sesbania Pea (Qld only), Summer Grass, Variegated Thistle, Winter Grass, Yellow Vine (Caltrop).

Re-entry Period

DO NOT enter treated areas until spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants or crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DANGEROUS TO FISH. DO NOT contaminate dams, rivers or streams with the product or used containers. Avoid use in channels and drains where roots or desirable plants may extend and in other situations where desirable trees grow close to treated areas.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Keep out of reach of children.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If refilling, replace cap and return to point of supply for refill or storage. If recycling, replace cap and return clean containers to a recycler or

designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. If product gets on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (phone 131 126).

Material Safety Data Sheet

If additional hazard information is required, refer to the Material Safety Data Sheet.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following. To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

NRA Approval No: 53761/0601

Batch No:

Date of Manufacture: