

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



PLANTVAX[®] 750 WP
SYSTEMIC FUNGICIDE

ACTIVE CONSTITUENT: 750 g/kg OXYCARBOXIN

GROUP	G	FUNGICIDE
-------	----------	-----------

For control of rust in
green beans,
geraniums, roses, carnations, irises,
chrysanthemums and snapdragons

Net Contents: 500 g, 1, 2 & 5 kg

(Crompton Logo)

Crompton Specialties Pty Ltd
3 Opala Street Regency Park SA 5010
Tel: (08) 8268 5700 Fax: (08) 8268 7555

DIRECTIONS FOR USE:

Foliar Spray:

Crop	Disease Controlled	States	Rate	Critical Comments
Carnations Irises Geraniums Roses Chrysanthemums Snapdragons	Rust	All States	130 g per 100 L water Small areas: 20 g per 15 L knapsack	Spray to run-off at 14 day intervals.
Boronia Fuchsia		TAS & WA		
Chrysanthemums	White Rust Puccinia (<i>Puccinia horiana</i>)	QLD & WA	130 g per 100 L	Apply at 5 to 7 day intervals. Rotation with other recommended fungicides may be necessary to avoid risk of build up of resistance.
Ornamentals	Rust	QLD & WA	130 g per 100 L	Apply on a 14 day schedule. Rotation with other recommended fungicides may be necessary to avoid risk of build up of resistance.
Green Beans	Rust	All States	1 kg/ha	High volume equipment: use 500 - 1,000 L water per ha. Low volume equipment: use 50 - 200 L water per ha. Apply at first sign of disease and again 14 days later.

Soil Treatment: All States

Crop & Condition	Disease Controlled	Rate of Irrigation per m ² of bed	Rate of Plantvax per 1,000 L of irrigation water	Critical Comments
Carnations Moderate incidence	Rust in the glasshouse	20 L	7 g	Apply this product by root drench in irrigation water at 7 day intervals. The rate may be varied as shown if a different volume per m ² of bed is desired or under severe rust conditions.
		10 L	14 g	
		5 L	28 g	
High incidence	Rust in the glasshouse	20 L	14 g	
		10 L	28 g	
		5 L	56 g	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

GREEN BEANS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Plantvax 750W Systemic Fungicide	Approved 0703
Company name change 30-6-03	Page 3 of 4

GENERAL INSTRUCTIONS:

This product is a wettable powder for use as a spray or in irrigation water.

Mixing:

Make a thin paste with portion of the water and the required quantity of this product. Dilute to the required volume under agitation.

Compatibility:

This product may be combined in the spray vat with any one of the following products: Agral[®] 600, Ambush[®] 500 EC, Endosan[®], Omite[®] and Pirimor[®]. A mixture of this product with any other product or with more than one of these products may be ineffective or may cause serious damage. The use of such a mixture is, therefore, entirely at the user's risk.

Fungicide Resistance Warning:

GROUP G FUNGICIDE

Plantvax 750W Systemic Fungicide is a member of the Oxathiin group of fungicides. For fungicide resistance management, Plantvax 750W Systemic Fungicide is a Group G fungicide.

Some naturally occurring individual fungi resistant to Plantvax 750W Systemic Fungicide and other Group G fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal populations if these fungicides are used repeatedly. These resistant fungi will not be controlled by Plantvax 750W Systemic Fungicide or other Group G fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crompton Specialties Pty Ltd accepts no liability for any losses that may result from the failure of Plantvax 750W Systemic Fungicide to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Avoid contact with eyes and skin. Do not inhale spray mist. Wash hands after use.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone *Australia* 131 126.

MATERIAL SAFETY DATA SHEET:

For additional information, refer to the Material Safety Data Sheet.

CONDITIONS OF SALE:

Crompton Specialties Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crompton Specialties Pty Ltd is limited to the replacement of the goods or (at the option of Crompton Specialties Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crompton Specialties Pty Ltd within thirty days of delivery.

- ® Plantvax & Omite are Registered Trade Marks of Crompton Corporation, USA.
- ® Agral, Ambush, Pirimor & Endosan are registered trademarks of Syngenta Crop Protection Pty Ltd.

In an Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY Crompton Specialties Pty Ltd 1800 033 111 ALL HOURS - AUSTRALIA WIDE
---	---

BN
DOM

Printer Code

NRA Approval No. 56010/0703