


EASY OPENING

PULL ON ONE OF THE ENDS OF THE RED THREAD

NC

405-106712-03-2003

EMB 64410 B



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

NOVOFIX®
FUNGICIDE



ACTIVE CONSTITUENTS:
300 g/kg MANCOZEB
120 g /kg COPPER PRESENT AS THE TRIBASIC SULPHATE

NOVOFIX®
DISPERSS

NOVOFIX®
DISPERSS

GROUP Y FUNGICIDE

For the control of Downy mildew and phomopsis in grape vines,
Anthracnose in mango and Target spot in capsicums, potatoes
and tomatoes as shown in the DIRECTIONS FOR USE table.

Net Contents:
20 kg



Manufacturer:
Cerexagri s.a.
1 rue des Frères Lumière B.P.9,
78373 PLAISIR Cedex - France.

Marketed by:
Crop Care Australasia Pty Ltd
Unit 15/16 Metroplex Avenue, Mararrie Qld 4172
Tel: (07) 3909 2000

Emergency Advice
Phone 1800 033 498
And ask for shift supervisor.

NOVOFIX®
DISPERSS



NOVOFIX® DISPERSS® FUNGICIDE

DIRECTIONS FOR USE

RECOMMENDATIONS
Do not apply from a windy day before the spray is dry.
Do not apply from a windy day after the spray is dry.
Do not apply from a windy day when the conditions are such as to cause the spray to drift or to be blown away.
Do not apply from a windy day when the conditions are such as to cause the spray to drift or to be blown away.

CROP	DISEASE	DATE	APPLICATOR	APPLICATOR COMMENTS
Grapevine	Downy mildew	15/10/2003	1000 L/ha	Apply to the underside of the leaves of the grapevine. Do not apply to the top of the leaves. Do not apply to the top of the leaves. Do not apply to the top of the leaves.
Mango	Anthracnose	15/10/2003	1000 L/ha	Apply to the underside of the leaves of the mango. Do not apply to the top of the leaves. Do not apply to the top of the leaves. Do not apply to the top of the leaves.
Potato	Target spot	15/10/2003	1000 L/ha	Apply to the underside of the leaves of the potato. Do not apply to the top of the leaves. Do not apply to the top of the leaves. Do not apply to the top of the leaves.
Tomato	Phomopsis	15/10/2003	1000 L/ha	Apply to the underside of the leaves of the tomato. Do not apply to the top of the leaves. Do not apply to the top of the leaves. Do not apply to the top of the leaves.

DO NOT BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

WARNING: KEEP OUT OF REACH OF CHILDREN. READ SAFETY DIRECTIONS BEFORE OPENING OR USING.

GENERAL INSTRUCTIONS:
NOVOFIX DISPERSS FUNGICIDE is a product for the control of certain fungi in grape vines, mangoes, capsicums, potatoes and tomatoes.

REMARKS:
Add the required quantity of NOVOFIX DISPERSS FUNGICIDE to the water in the spray tank and mix thoroughly. Mix can application during morning and application.

APPLICATION:
Downy Mildew
Apply to the underside of the leaves of the grapevine. Do not apply to the top of the leaves. Do not apply to the top of the leaves. Do not apply to the top of the leaves.

ENVIRONMENTAL HAZARDOUS SUBSTANCE, SOLID, N.O.S (MANCOZEB), UN 3077

HAZCHEM: 2X

EMERGENCY ADVICE
Please refer to the MSDS and ask for shift supervisor.
This product is not considered to be a Dangerous Good under the Australian Code for the Transport of Dangerous Goods by Road or Rail.

APMA Approved No. 56902/2003/0600
Batch No. see bottom of the bag
Date of Manufacture see bottom of the bag
Expiry Date: 2 years after the Date of Manufacture.

REGISTERED TRADEMARK OF CEREXAGRI S.A.

Distributed by:
Arkema Pty Ltd
600 Victoria St.
Richmond, VIC 3121
ABN 000 330 772

Cerexagri s.a.
1 rue des Frères Lumière
B.P.9 - 78373 PLAISIR Cedex - France
TEL: 33 1 30 81 73 00 FAX: 33 1 30 81 72 50
email: contact@cerexagri.com
R.C.S. Versailles 569 904 902



U N 5M2 / Z 21 / S / 2003 / F / BVT 199226 / CEREXAGRI

001-106712-03-2003 (Rev.01)

160mm(w)x360mm(h) 48%

NOVOFIX® ISPERSS® FUNGICIDE

DIRECTIONS FOR USE

RESTRAINTS

Do NOT apply if rain is likely before the spray is dry
 Do NOT apply to wet foliage
 Do NOT apply this product when hot conditions (>35 C) or frosts occur as damage may result
 Do NOT apply this product to copper sensitive crops or cultivars

CROP	DISEASE	RATE	WITH-HOLDING PERIOD	CRITICAL COMMENTS
Grape vines	Downy mildew (<i>Plasmopara viticola</i>)	Dilute spraying 300 to 350 g/100L water Concentrate spraying Refer to Mixing/Application section	14 days	If Downy mildew is expected begin spray application at 10-15 cm stage of shoot growth. Continue a programme of application every 10 to 14 days until the risk of infection diminishes. Use the higher rate when infection levels are particularly high. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
	Phomopsis Cane and Leaf spot			Spray at bud burst and repeat 7-10 days later. Use the higher rate when infection levels are particularly high.
Mangoes	Anthraxnose (<i>Colletotrichum gloeosporioides</i>)	Dilute spraying 300 g/100L water Concentrate spraying Refer to Mixing/Application section	1 day	Spray at weekly intervals during flowering then at two to three week intervals until harvest. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Potatoes	Target spot (<i>Alternaria solani</i>)	Dilute spraying 300 to 350 g/100L water Concentrate spraying Refer to Mixing/Application section	1 day	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use higher rate and shorter intervals during weather conditions favourable to disease development.
Tomatoes	Target spot (<i>Alternaria solani</i>)	Dilute spraying 300 to 350g/100 L water Concentrate spraying Refer to Mixing/Application section	7 days	Spray at first sign of disease. Repeat at 7-10 day intervals. Do not apply to tomato seedlings before transplanting. Use higher rate and shorter intervals during weather conditions favourable to disease development.
Capsicums			14 days	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS (WHP)

CAPSICUMS, GRAPE VINES: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

TOMATOES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

MANGOES, POTATOES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

GENERAL INSTRUCTIONS:

NOVOFIX ISPERSS FUNGICIDE is a protectant fungicide for the control of certain fungus diseases in grape vines, mangoes, capsicums, potatoes and tomatoes.

MIXING

Add the required quantity of NOVOFIX ISPERSS FUNGICIDE to water in the spray tank and mix thoroughly. Maintain agitation during mixing and application.

APPLICATION

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can be then calculated in the following way:

EXAMPLE ONLY

- Dilute spray volume as determined above: For example 1000 L/ha
 - Your chosen concentrate spray volume: For example 250 L/ha
 - The concentration factor in this example is: 4 X (i.e. 1000 L/250 L = 4)
 - If the dilute label rate is 300 g/100 L, then the concentrate rate becomes 4 x 300, that is 1.2 kg/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake competency training and follow industry Best Practices.

Aerial application

This product may be applied by agricultural aircraft. Use at least 30 to 50 litres spray mixture per hectare. Reference should be made to the Directions for Use table and instructions for preparing concentrate sprays for the correct rate of application in specific crop situations.

Compatibility:

This product is compatible with formulations of most commonly used fungicides, insecticides and miticides. Check compatibility before using by preparing a test solution of the required proportions and observing stability over several minutes.

FUNGICIDE RESISTANCE WARNING

GROUP Y FUNGICIDE

For fungicide resistance management NOVOFIX ISPERSS FUNGICIDE is a Group Y fungicide. Some naturally occurring individual fungi resistant to NOVOFIX ISPERSS FUNGICIDE may exist through normal genetic variability in any fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by NOVOFIX ISPERSS FUNGICIDE and other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure NOVOFIX ISPERSS FUNGICIDE to control resistant fungi.

PRECAUTIONS

Do not use human flaggers/markers unless they are protected by engineering controls such as enclosed cabs.

Re-Entry: Do not allow entry into treated areas until spray has dried. When prior entry is necessary, wear personal protective equipment as specified in the Safety Directions. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, that may cause spray drift or run-off onto sensitive areas including (but not limited to) streams, rivers or waterways. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

DO NOT feed treated crops and or trash to livestock.

PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic organisms. Do NOT allow spray drift or run-off onto sensitive areas including (but not limited to) streams, rivers or waterways. Do not contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container, in a well-ventilated area, building or room, as cool as possible, away from children, animals or fodder. Keep away from foodstuffs. Do NOT store for prolonged periods in direct sunlight.

Shake empty bag into spray tank. Do not dispose of undiluted chemicals on site. Puncture or shred or bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Will damage eyes. Will irritate nose, throat and skin. Avoid contact with eyes and skin. Do not inhale dust or spray mist. If product in eyes, wash it out immediately with water.

When opening the container and preparing the spray, wear elbow-length nitrile or PVC gloves, goggles and disposable dust mask.

When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length nitrile or PVC gloves or goggles. Wash hands after use. After each days use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 13 11 26.

Avoid giving alcohol. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

CONDITIONS OF SALE:

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.


Emergency Advice
 Phone 1800 033 498 and ask for shift supervisor
 This product is not considered to be a Dangerous Good under the Australian Code for the Transport of Dangerous Goods by Road or Rail.

APVMA Approval No: 56997/20kg/0806
 Batch No: see bottom of the bag
 Date of Manufacture: see bottom of the bag
 Expiry Date: 2 years after the Date of Manufacture.

® Registered trademark of Cerexagri s.a.

Distributed by :
 Arkema Pty Ltd
 600 Victoria St,
 Richmond, VIC, 3121
 ABN 000 330 772

Cerexagri s.a
 1 rue des Frères Lumière -
 B.P.9 - 78373 PLAISIR Cedex - France
 TEL. : 33 1 30 81 73 00 FAX : 33 1 30 81 72 50
 e-mail : contact@cerexagri.com
 R.C.S. Versailles 569 804 982



**ENVIRONMENTALLY HAZARDOUS
 SUBSTANCE, SOLID, N.O.S
 (MANCOZEB), UN 3077
 HAZCHEM: 2X**

501-108712-03-2003 (text ref.)



5M2 / Z 21 / S / 2003 / F / BVT 199226 / CEREXAGRI