

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Conquest Drover Liquid Seed Dressing Fungicide

ACTIVE CONSTITUENT: 250g/L IPRADIONE
SOLVENT: 316 g/L LIQUID HYDROCARBONS

GROUP	B	FUNGICIDE
-------	----------	-----------

For the control of Brown Spot and the suppression of Hypocotyl Rot, in the first four weeks after emergence, in Lupins and for the control of Hypocotyl Rot in potatoes as specified in the Directions for Use Table.

Contents: 20 Litres

(5L, 60L, 100L, 110L, 200L and 1000L)

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932
Suite 10, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9204 2831

DIRECTIONS FOR USE

Crop	Disease	State	Rate	Critical Comments
Lupins	Brown Leaf Spot (<i>Pleiochaeta setosa</i>) Rhizoctonia hypocotyl Rot (<i>Rhizoctonia solani</i>) (Suppression only)	WA only	Pre- sowing treatment (Feb-May) 100-400mL /100kg seed Post- harvest treatment (Dec-Jan) 400mL /100kg seed	Mix with water and apply in a total volume of 600mL mixture/100kg seed using suitable seed dressing application equipment. Ensure thorough coverage of seed. Use in conjunction with other disease management practices. See General Instructions for details. Use high rate in areas likely to be exposed to high disease pressure. (Eg. Central and Eastern Wheatbelt areas with close rotations and a history of Brown Leaf Spot or where other disease management practices are not followed). Application of Conquest Drover Liquid Seed Dressing Fungicide will provide suppression of Hypocotyl Rot and can reduce Pleiochaeta Root Rot in some situations.
	Brown Leaf Spot (<i>Pleiochaeta setosa</i>)	NSW, ACT, Vic, SA only	100 mL - 500 mL /100kg seed in 500 mL water	Apply using suitable seed dressing application equipment. Ensure thorough coverage of seed. Application can reduce Pleiochaeta Root Rot in some situations. Use high rate in areas likely to be exposed to high disease pressure. (Eg. areas with close rotations and a history of Brown Leaf Spot or where other disease management practices are not followed).
Potatoes	Hypocotyl Rot (Black Scurf) (<i>Rhizoctonia solani</i>)	All States	800 mL /tonne seed material	Conquest Drover Liquid Seed Dressing Fungicide will protect emerging shoots from Hypocotyl Rot, improving overall germination. Conquest Drover Liquid Seed Dressing Fungicide may also reduce occurrence of Black Scurf on the harvested potatoes. Ensure good coverage of seed material and planting furrow. Do not plant into waterlogged soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD (Harvest and Grazing)

Lupins, Potatoes: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

Lupins only: This product will reduce inoculum levels on inoculated seed and should only be used on seed to be sown in ground where existing soil inoculum levels will provide adequate nodulation. Spray inoculation is not recommended.

This product should be used as part of an integrated disease management strategy, particularly in areas with high disease pressure. Where possible, the following disease management practices should be used in conjunction with Conquest Drover Liquid Seed Dressing Fungicide:

- ◆ Stubble Retention
- ◆ Direct Drilling
- ◆ Correct Seeding Depth
- ◆ Correct Rotations
- ◆ Clean Seed Source
- ◆ Correct Seeding Rate
- ◆ Avoid Late sowing

Fungicide Resistance Warning

GROUP	B	FUNGICIDE
-------	----------	-----------

Conquest Drover Liquid Seed Dressing Fungicide is a member of the Dicarboximide group of fungicides. For fungicide resistance management the product is a Group B fungicide.

Some naturally occurring individual fungi resistant to Conquest Drover Liquid Seed Dressing Fungicide and other Group B fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled Conquest Drover Liquid Seed Dressing Fungicide or other Group B fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Conquest Agrochemicals Pty Ltd accepts no liability for any losses that may result from the failure of Conquest Drover Liquid Seed Dressing Fungicide to control resistant fungi.

Mixing

Shake the container well before using. Add the required volume of Conquest Drover Liquid Seed Dressing Fungicide to the water while agitating.

Application

Lupins only: Slowly add the mixture to the seed and mix thoroughly ensuring thorough coverage of seed.

Potatoes only: Ensure good coverage of seed material and planting furrow. This can be achieved by applying Conquest Drover Liquid Seed Dressing Fungicide as a fine spray to the seed at the time of planting using spray equipment mounted on the planter, and nozzles located at three points on each planter row to ensure uniform coating of the seed. A minimum water volume of 80L/tonne should be used.

Further information can be obtained from a Conquest Agrochemicals Pty Ltd. representative.

PRECAUTION

When treated seed is stored, it should be kept apart from other grains or seeds, and the containers or bags should be clearly marked to show that the contents have been treated with Conquest Drover Liquid Seed Dressing Fungicide. DO NOT use treated seed for human consumption. DO NOT allow treated seed to contaminate food for human consumption. Bags that have held treated seed should not be used for any other purpose. Any seed not used for sowing should be destroyed.

PROTECTION OF LIVESTOCK

DO NOT feed treated seed to animals, including poultry. DO NOT allow treated seed to contaminate grain intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose wildlife or domestic birds. DO NOT contaminate streams, rivers or waterways with chemical or bags which have held treated seed.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Avoid contact with eyes and skin and avoid inhalation of vapour. Wear suitable protective clothing, gloves and goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking and smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

For specialist advice in an emergency dial

1 800 033 111

24 hours Australia wide.

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932

Suite 10, 41 Walters Drive

Herdsmen Business Park

Osborne Park WA 6017

Telephone (08) 9244 1268

Facsimile: (08) 9204 2831

Batch No:

DOM:

NRA Approval No: 57016/0703