



CAUTION
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST
STINGRAY
 430 FUNGICIDE

ACTIVE CONSTITUENT: 430g/L TEBUCONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, papaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

CONQUEST AGROCHEMICALS PTY. LTD
 ABN 84 098 814 932
 Unit 1, 46 Hasler Road, Osborne Park, WA 6017
 Telephone: 08 9347 0500, Fax 08 9347 0551

CONTENTS

1L

5L

STORAGE AND DISPOSAL
 Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.
 Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
 If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.
For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS
 Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID
 If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET
 Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE
 Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
---	--

Batch No: _____

DOM: _____

APVMA APPROVAL No: 57810/1L/0409

57810/5L/0409

CONQUEST AGROCHEMICALS PTY. LTD
 ABN 84 098 814 932
 Unit 1, 46 Hasler Road, Osborne Park, WA 6017
 Telephone: 08 9347 0500, Fax 08 9347 0551

Prepared For: Conquest Agrochemicals Pty Limited Unit 1, 46 Hasler Rd, PERTH, WA		Regulatory Author: Conquest Agrochemicals Pty Limited		APVMA Approval No: 57810/1L/0409		
Prepared By: 	Detail: Name: Conquest Stingray 430 Fungicide 1L Label Last Revision: 25 May 2009 Size: 260mm x 145mm	Colour Key: C M Y BLACK 	% Proof Sizing of actual article A3 sheet 100%			Copyright © 2008 Conquest Agrochemicals Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Agrochemicals Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST
STINGRAY
430 FUNGICIDE

ACTIVE CONSTITUENT: 430g/L TEBUCONAZOLE

GROUP 3 FUNGICIDE



APVMA
APPROVED LABEL

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, papaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE



CONQUEST
Agrochemicals

CONQUEST AGROCHEMICALS PTY. LTD
 ABN 84 098 814 932
 Unit 1, 46 Hasler Road, Osborne Park, WA 6017
 Telephone: 08 9347 0500, Fax 08 9347 0551

CONTENTS

1L

5L

STORAGE AND DISPOSAL
 Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.
 Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
 If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.
For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS
 Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID
 If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET
 Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE
 Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
---	--

Batch No: _____

DOM: _____


APVMA APPROVAL No: 57810/1L/0409

57810/5L/0409



CONQUEST
Agrochemicals

CONQUEST AGROCHEMICALS PTY. LTD
 ABN 84 098 814 932
 Unit 1, 46 Hasler Road, Osborne Park, WA 6017
 Telephone: 08 9347 0500, Fax 08 9347 0551

Prepared For: Conquest Agrochemicals Pty Limited Unit 1, 46 Hasler Rd, PERTH, WA	Regulatory Author: Conquest Agrochemicals Pty Limited	APVMA Approval No: 57810/5L/0409										
Prepared By: 	Detail: Name: Conquest Stingray 430 Fungicide 5L Label Last Revision: 25 May 2009 Size: 260mm x 145mm	Colour Key <table style="width: 100%; text-align: center; font-size: x-small;"> <tr> <td>C</td> <td>M</td> <td>Y</td> <td>BLACK</td> </tr> <tr> <td style="width: 20px; height: 15px; background-color: cyan;"></td> <td style="width: 20px; height: 15px; background-color: magenta;"></td> <td style="width: 20px; height: 15px; background-color: yellow;"></td> <td style="width: 20px; height: 15px; background-color: black;"></td> </tr> </table>	C	M	Y	BLACK					% Proof Sizing of actual article A3 sheet <h1 style="text-align: center; margin: 0;">100%</h1>	Copyright © 2008 Conquest Agrochemicals Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Agrochemicals Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.
C	M	Y	BLACK									

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST
STINGRAY
430 FUNGICIDE

ACTIVE CONSTITUENT: 430g/L Tebuconazole

GROUP 3 FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, papaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

CONQUEST Agrochemicals
CONQUEST AGROCHEMICALS PTY. LTD
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park, WA 6017
Telephone: 08 9347 0500, Fax 08 9347 0551

CONTENTS
10L
20L

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.
For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
---	--



Batch No:

DOM:

APVMA APPROVAL No: 57810/10L/0409
57810/20L/0409

CONQUEST AGROCHEMICALS PTY. LTD
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park, WA 6017
Telephone: 08 9347 0500, Fax 08 9347 0551


Prepared For: **Conquest Agrochemicals Pty Limited** Unit 1, 46 Hasler Rd, PERTH, WA

Regulatory Author: **Conquest Agrochemicals Pty Limited**

APVMA Approval No: **57810/10L/0409**



Detail:
Name: Conquest Stingray 430 Fungicide 10L Label
Last Revision: 25 May 2009
Size: 350mm x 150mm



% Proof Sizing of actual article A3 sheet
100%

Copyright © 2008 Conquest Agrochemicals Pty Ltd
All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Agrochemicals Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST
STINGRAY
430 FUNGICIDE

ACTIVE CONSTITUENT: 430g/L Tebuconazole
GROUP 3 FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, papaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE



APVMA
APPROVED
LABEL

CONQUEST AGROCHEMICALS PTY. LTD
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park, WA 6017
Telephone: 08 9347 0500, Fax 08 9347 0551

CONTENTS
10L
20L

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.
For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
---	--



Batch No:

DOM:

APVMA APPROVAL No: 57810/10L/0409
57810/20L/0409

CONQUEST AGROCHEMICALS PTY. LTD
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park, WA 6017
Telephone: 08 9347 0500, Fax 08 9347 0551

Prepared For: **Conquest Agrochemicals Pty Limited** Unit 1, 46 Hasler Rd, PERTH, WA

Regulatory Author: **Conquest Agrochemicals Pty Limited**

APVMA Approval No: **57810/20L/0409**



Detail:
Name: Conquest Stingray 430 Fungicide 20L Label
Last Revision: 25 May 2009
Size: 350mm x 150mm



% Proof Sizing of actual article A3 sheet
100%

Copyright © 2008 Conquest Agrochemicals Pty Ltd
All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Agrochemicals Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.

In a Transport Emergency

**Dial
000**

Police or Fire Brigade

FOR SPECIALIST ADVICE
IN AN EMERGENCY DIAL

1800 033 111

ALL HOURS AUSTRALIA WIDE



Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932

Unit 1, 46 Hasler Road
Osborne Park, WA 6017

Telephone (08) 9347 0500

Facsimile: (08) 9347 0551

APVMA Approval No: 57810/0409

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

STINGRAY

430 FUNGICIDE

ACTIVE CONSTITUENT: 430g/L Tebuconazole

GROUP 3 FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, papaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: READ THIS BOOKLET BEFORE USE



CONQUEST AGROCHEMICALS PTY. LTD

ABN 84 098 814 932

Unit 1, 46 Hasler Road, Osborne Park, WA 6017

Telephone: 08 9347 0500, Fax 08 9347 0551

CROPS	DISEASE	APPLICATION	RATE L/ha	WHP (days)	CRITICAL COMMENTS
Bananas (Qld, NSW, WA, NT only)	Leaf spot (yellow sigatoka) Leaf speckle Black sigatoka	Ground Application Apply by misting machine or airstblast sprayer in a convenient volume of water Aerial Application Apply at least 20 L of spray mixture per hectare	230 mL/ha Add 3-6 L water miscible oil/ha	1 day	Maintain good deleafing practices to reduce disease inoculum. Very old leaves and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of Stingray. Tropical areas (e.g. North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a minimum of 2 and a maximum of 3 consecutive Stingray sprays at 14-day intervals. Do not apply more than 6 Stingray sprays in any 12-month period. Do not apply any Stingray sprays in the months of July, August and September. Sub-tropical areas (e.g. South Qld, NSW): Commence spraying with Stingray at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a minimum of 2 consecutive Stingray sprays. Do not apply more than 5 Stingray sprays in any 12-month period. Stingray is approved for use in banana plantations interplanted with avocados.
Peanuts (South QLD and NSW only)	Early leaf spot Late leaf spot Rust Net blotch	Low disease Ground Application Apply in at least 100 L of water/ha Aerial Application Apply at least 30 L of spray mixture per hectare	175 mL/ha +Agridex 1 L/ha	H 3 weeks G 3 weeks	Regularly check high-risk areas in the crop for disease (eg lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found then either; - spray <u>immediately</u> after the last disease conducive weather (e.g. rain or heavy dews) OR -spray before the next disease conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat before or just after the next disease conducive weather. For resistance strategy, see General Instructions.
		High disease OR Wet weather	290 mL/ha +Agridex 1 L/ha		
		High disease AND Wet weather	440 mL/ha +Agridex 1 L/ha		

CROPS	DISEASE	APPLICATION	RATE L/ha	WHP (days)	CRITICAL COMMENTS
Peanuts (North QLD, WA, NT only)	Early leaf spot Late leaf spot Rust	Low disease	230 mL/ha +Agridex 1 L/ha	H 3 weeks G 3 weeks	Begin spraying at 3-4 weeks after planting. If band spraying, apply half the appropriate rate as a 45 cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14-day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.
		Moderate disease	290 mL/ha +Agridex 1 L/ha		
		Severe disease	440 mL/ha +Agridex 1 L/ha		
	Net blotch	Low disease	290 mL/ha +Agridex 1 L/ha		Apply at 14-day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.
		Moderate – severe disease	440 mL/ha +Agridex 1 L/ha		
	Green beans	Rust	Can be applied by aircraft or ground rig		350 mL/ha +Agridex 1 L/ha
Peas	Powdery mildew	Ground Application Apply in at least 50 L of water/ha Aerial Application Apply in at least 10 L of water/ha	145 mL/ha	Apply at flowering or at first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions.	

CROPS	DISEASE	APPLICATION	RATE L/ha	WHP (days)	CRITICAL COMMENTS
Wheat	Leaf rust, Stripe rust Septoria nodorum blotch, Yellow leaf spot	Ground Application Apply in at least 50 L of water/ha Aerial Application Apply in at least 10 L of water/ha	145 mL/ha or 290 mL/ha	H 5 weeks G 14 Days	Use higher rate when longer disease control is required. Stripe rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil (e.g. D-C Trate or equivalent) at 1%, may improve performance of Stingray on wheat, oats and barley.
	Septoria tritici blotch		290 mL/ha		
Oats	Crown rust	145 or 290 mL/ha			
Barley	Scald	145 mL/ha			
	Powdery mildew	145 or 290 mL/ha			
Wheat, Oats	Stem rust				
Onions (TAS only)	White root rot	Before sowing, apply Stingray onto lime super. Ensure good coverage of all lime super particles.	1.45 mL/100m of row mixed with 145-218g lime super/ 100m of row	-	Apply Stingray treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a bandwidth of 2 cm. Ensure that the correct rate of Stingray is used otherwise some delay in emergence and reduced stands of seedlings may occur.
Papaw	Black spot	Ensure thorough coverage of leaves and fruit	290 mL/ha	3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of Stingray. Apply Stingray at 14-day intervals. Alternate Stingray with sprays of a protectant fungicide (eg Mancozeb). Do not apply more than 6 sprays of Stingray (or any DMI fungicide) on any block in any 12-month period.

CROPS	DISEASE	APPLICATION	RATE L/ha	WHP (days)	CRITICAL COMMENTS
Pyrethrum	<i>Sclerotinia sclerotiorum</i>		350 mL/ha	-	Apply twice, in rotation with other control measures, at 7-10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf rust Stem rust	Apply in at least 100 L of water/ha	290 mL/ha	-	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or re-infection may require a further application 3 - 4 weeks later. Ensure thorough coverage and use higher water volumes in dense or advanced crops.
Lettuce (ACT, NSW, QLD, SA, TAS, NT, WA)	Sclerotinia Rot (<i>Sclerotinia sclerotiorum</i>)		350 mL/ha	H 5 weeks	Apply by boom spray. Apply only during the early stages of plant development (note the 5 week WHP). Apply a maximum of 2 applications at 7-10 day intervals in rotation with other chemicals registered for this use (i.e procymidone, iprodione and benomyl). Apply only to field crops. DO NOT apply in greenhouse (i.e protected cropping) situations or hydroponically grown lettuce crops. Control of Sclerotinia in lettuce should not be solely reliant on fungicides. Other control measures such as crop rotations, resistant varieties and planting techniques should be employed.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

BANANAS, AVOCADOS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

BEANS, PEAS: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 DAYS AFTER APPLICATION.

CEREALS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

ONIONS: NOT REQUIRED WHEN USED AS DIRECTED

PAPAW: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

PEANUTS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION

LETTUCE: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION

H = Harvest, G = Grazing

Fungicide Resistance Warning**GROUP 3 FUNGICIDE**

Conquest Stingray 430 Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Conquest Stingray 430 Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Conquest Stingray 430 Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Conquest Agrochemicals Pty Limited accepts no liability for any losses that may result from the failure of Stingray 430 Fungicide to control resistant fungi.

Resistance management recommendation – peanuts

Apply no more than 3 consecutive sprays of DMI fungicide (eg Stingray) before switching to a non-DMI fungicide. Apply no more than 5 DMI sprays per season.

Mixing:

Prior to pouring, shake container vigorously, then add the required quantity of Stingray 430 Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex® (peanuts, beans) or water miscible oil (bananas) and mix thoroughly.

Application

Aircraft should fly as low as possible under the prevailing conditions to minimise drift.

Special Warning – bananas

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking. ***Never include adjuvants*** other than water miscible oils with Stingray sprays. The inclusion of wetting agents is known to cause phototoxicity to young fruit.

Foliar diseases on cereal crops

Do not apply to cereal crops more than once in a season. Treatment will give approximately three weeks disease suppression. Economic responses may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application. Yield potential: crops with potential yield under 2t/ha are unlikely to give economic responses to a fungicide spray except under conditions of very severe disease. Economic responses are most likely with crops with potential yield of over 3t/ha.

Spray timings for stripe rust control

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties. In South Australia consult Plant Protection Note PPN 21. In other states the following spray program is suggested:

Seedling infections: When approximately 20 out of 100 leaves show first signs of infection during tillering to jointing, apply a spray within one week.

Adult infections – susceptible varieties: When approximately 10 out of 100 leaves show first sign of infection, apply a spray within one week.

Adult infections – moderately susceptible varieties: When approximately 15 to 20 leaves out of 100 leaves show first sign of infection, apply a spray within one week. Do not delay.

Adult infections – moderately resistant and resistant varieties: Monitor carefully. If rust appears and spreads, spray within one week.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

A spray drift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.