

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP

APVMA
✓
APPROVED LABEL

CONQUEST

EXPLODE 250


HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.

CONQUEST
CROP PROTECTION

CONQUEST CROP PROTECTION PTY LTD.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

CONTENTS

5L	<input checked="" type="checkbox"/>	100L	<input type="checkbox"/>
200L	<input type="checkbox"/>	220L	<input type="checkbox"/>



FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE.
THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN

STORAGE AND DISPOSAL (5, 20 and 200L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

(100, 110L and 220L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of storage for refill or storage.

SAFETY DIRECTIONS
Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION
Do not work in spray mist.
Do not continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gaitlength gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator, which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID
If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26).
If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

NOTE TO PHYSICIAN
For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET
Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE
Conquest Crop Protection Pty Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

UN NO. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police, or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X














Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551




Batch No: _____

DOM: _____

APVMA Approval No: 57817/5L/0510 ☒ 57817/100L/0510 ☐
57817/200L/0510 ☐ 57817/220L/0510 ☐

Prepared For: Conquest Crop Protection Pty Limited 46 Hasler Rd, PERTH, WA 6017		Regulatory Author: Conquest Crop Protection Pty Limited		APVMA Approval No: 57817/5L/0510													
Prepared By: 		Detail: Name: Conquest Explode 250 Herbicide 5L Last Revision: 10 June 2010 Size: 460mm x 165mm		Colour Key <table border="1" data-bbox="991 1879 1406 1902"> <tr> <td>C</td> <td>M</td> <td>Y</td> <td>K</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </table>		C	M	Y	K					% Proof Sizing of actual article A3 sheet <div style="font-size: 48px; text-align: center;">80%</div>		Copyright © 2008 Conquest Crop Protection Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Crop Protection Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.	
C	M	Y	K														
																	

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP

APVMA

APPROVED LABEL

CONQUEST

EXPLODE 250

HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.

CONQUEST

CROP PROTECTION

CONQUEST CROP PROTECTION PTY LTD.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

CONTENTS

20L☒ 110L☐

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE.
THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN

STORAGE AND DISPOSAL (5, 20 and 200L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure ring containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. (If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

(100, 110L and 220L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of storage for refill or storage.

SAFETY DIRECTIONS
Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION
Do not work in spray mist.
Do not continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gait length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator, which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID
If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.


NOTE TO PHYSICIAN
For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET
Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.


CONDITIONS OF SALE
Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

UN NO. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police, or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551



TOXIC
6




CONQUEST
CROP PROTECTION

Batch No: ☐ 57817/20L/0510 ☒ 57817/110L/0510 ☐

DOM: ☐

APVMA Approval No: 57817/20L/0510 ☒ 57817/110L/0510 ☐

Prepared For: Conquest Crop Protection Pty Limited 46 Hasler Rd, PERTH, WA 6017		Regulatory Author: Conquest Crop Protection Pty Limited		APVMA Approval No: 57817/20L/0510							
Prepared By: 	Detail:	Colour Key	% Proof Sizing of actual article A3 sheet		Copyright © 2008 Conquest Crop Protection Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Crop Protection Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.						
	Name: Conquest Explode 250 Herbicide 20L Last Revision: 10 June 2010 Size: 460mm x 165mm	<table><tr><td>C</td><td>M</td><td>Y</td><td>K</td></tr><tr><td><div></div></td><td><div></div></td><td><div></div></td><td><div></div></td></tr></table>	C	M		Y	K	<div></div>	<div></div>	<div></div>	<div></div>
C	M	Y	K								
<div></div>	<div></div>	<div></div>	<div></div>								

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP

APVMA

APPROVED LABEL

CONQUEST

EXPLODE 250

HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.

CONQUEST

CROP PROTECTION

CONQUEST CROP PROTECTION PTY LTD.

ABN 84 098 814 932

Unit 1, 46 Hasler Road, Osborne Park WA 6017

Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

CONTENTS

5L

100L

200L

☐

☒

☐

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE.
THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN

STORAGE AND DISPOSAL (5, 20 and 200L only)
Store in the closed, original container in a cool, well-ventilated area.
Do NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

(100, 110L and 220L only)
Store in the closed, original container in a cool, well-ventilated area.
Do NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of storage for refill or storage.

SAFETY DIRECTIONS
Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. Do NOT inhale spray mist. After use before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION
Do not work in spray mist.
Do not continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gaitlength gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator, which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID
If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26).
If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

NOTE TO PHYSICIAN
For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET
Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE
Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.














UN NO. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police, or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

Batch No:

DOM:

APVMA Approval No:	57817/5L/0510	<input type="checkbox"/>	57817/100L/0510	<input checked="" type="checkbox"/>
	57817/200L/0510	<input type="checkbox"/>	57817/220L/0510	<input type="checkbox"/>

Prepared For: Conquest Crop Protection Pty Limited 46 Hasler Rd, PERTH, WA 6017		Regulatory Author: Conquest Crop Protection Pty Limited		APVMA Approval No: 57817/100L/0510											
Prepared By: 		Detail: Name: Conquest Explode 250 Herbicide 100L Last Revision: 10 June 2010 Size: 460mm x 165mm		Colour Key: <table border="1" data-bbox="991 1873 1421 1902"> <tr> <td>C</td> <td>M</td> <td>Y</td> <td>K</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </table>		C	M	Y	K					% Proof Sizing of actual article A3 sheet <div style="font-size: 48px; text-align: center;">80%</div>	Copyright © 2008 Conquest Crop Protection Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Crop Protection Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.
C	M	Y	K												
															

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP

APVMA

APPROVED LABEL

CONQUEST

EXPLODE 250

HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.

CONQUEST
CROP PROTECTION

CONQUEST CROP PROTECTION PTY LTD.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

CONTENTS
20L ☐ 110L ☒

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE

THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN

STORAGE AND DISPOSAL (5, 20 and 200L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

(100, 110L and 220L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of storage for refill or storage.

SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION

Do not work in spray mist.

Do not continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gaitlength gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator, which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.

Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID

If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

NOTE TO PHYSICIAN

For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

UN NO. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police, or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932

Unit 1, 46 Hasler Road, Osborne Park WA 6017



Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

Batch No:

DOM:

APVMA Approval No: 57817/20L/0510 ☐

57817/110L/0510 ☒

Prepared For: Conquest Crop Protection Pty Limited 46 Hasler Rd, PERTH, WA 6017		Regulatory Author: Conquest Crop Protection Pty Limited		APVMA Approval No: 57817/110L/0510	
Prepared By: 	Detail:		Colour Key		% Proof Sizing of actual article A3 sheet
	Name: Conquest Explode 250 Herbicide 110L Last Revision: 10 June 2010 Size: 460mm x 165mm				80%
Copyright © 2008 Conquest Crop Protection Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Crop Protection Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.					

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CAN KILL IF SWALLOWED

DO NOT PUT IN DRINK BOTTLES

KEEP LOCKED UP

APVMA

APPROVED LABEL

CONQUEST

EXPLODE 250

HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.

CONQUEST

CROP PROTECTION

CONQUEST CROP PROTECTION PTY LTD.

ABN 84 098 814 932

Unit 1, 46 Hasler Road, Osborne Park WA 6017

Telephone (08) 9357 0500, Facsimile: (08) 9347 0551

CONTENTS

5L

200L

100L

220L

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE.
THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN

STORAGE AND DISPOSAL (5, 20 and 200L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

(100, 110L and 220L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of storage for refill or storage.

SAFETY DIRECTIONS
Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION
Do not work in spray mist.
Do not continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator, which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID
If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.


NOTE TO PHYSICIAN
For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET
Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE
Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

UN NO. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police, or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551
















CONQUEST
CROP PROTECTION

Batch No: _____

DOM: _____

APVMA Approval No: 57817/5L/0510 ☐ 57817/100L/0510 ☐
57817/200L/0510 ☒ 57817/220L/0510 ☐

Prepared For: Conquest Crop Protection Pty Limited 46 Hasler Rd, PERTH, WA 6017		Regulatory Author: Conquest Crop Protection Pty Limited		APVMA Approval No: 57817/200L/0510													
Prepared By: 		Detail: Name: Conquest Explode 250 Herbicide 200L Last Revision: 10 June 2010 Size: 460mm x 165mm		Colour Key: <table border="1" data-bbox="994 1896 1439 1915"> <tr> <td>C</td> <td>M</td> <td>Y</td> <td>K</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </table>		C	M	Y	K					% Proof Sizing of actual article A3 sheet <div style="font-size: 48px; text-align: center;">80%</div>		Copyright © 2008 Conquest Crop Protection Pty Ltd <small>All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Crop Protection Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.</small>	
C	M	Y	K														
																	

DANGEROUS POISON
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING
 CAN KILL IF SWALLOWED
 DO NOT PUT IN DRINK BOTTLES
 KEEP LOCKED UP

CONQUEST

EXPLODE 250


HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.



FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE.
THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN

STORAGE AND DISPOSAL (5, 20 and 200L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose care of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

(100, 110L and 220L only)
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of storage for refill or storage.

SAFETY DIRECTIONS
Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION
Do not work in spray mist.
Do not continue to use if you get skin irritation or nosebleeds. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator, which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID
If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26).
If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.


NOTE TO PHYSICIAN
For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET
Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.


CONDITIONS OF SALE
Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

UN NO. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police, or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Unit 1, 46 Hasler Road, Osborne Park WA 6017
Telephone (08) 9357 0500, Facsimile: (08) 9347 0551






TOXIC
6



CONQUEST
CROP PROTECTION

Batch No: ☐ 57817/5L0510 ☐ 57817/100L0510 ☐
DOM: ☐ 57817/200L0510 ☐ 57817/220L0510 ☒

Prepared For: Conquest Crop Protection Pty Limited 46 Hasler Rd, PERTH, WA 6017		Regulatory Author: Conquest Crop Protection Pty Limited		APVMA Approval No: 57817/220L/0510	
Prepared By: 	Detail:	Colour Key	% Proof Sizing of actual article A3 sheet		Copyright © 2008 Conquest Crop Protection Pty Ltd All rights Reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or in any information storage or retrieval system, without prior permission in writing from Conquest Crop Protection Pty Ltd. All translation rights are reserved by the publisher. Respective Trademarks and copyrights are held by the relevant owners.
	Name: Conquest Explode 250 Herbicide 220L Last Revision: 10 June 2010 Size: 460mm x 165mm				

UN No. 3016	BIPYRIDILIUM PESTICIDE, LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency Dial 000 Police or Fire Brigade	FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
PG III	HAZCHEM 2 X



Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932
Unit 1, 46 Hasler Road
Osborne Park, WA 6017

Telephone (08) 9347 0500
Facsimile: (08) 9347 0551

APVMA Approval No: 57817/0510

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP

CONQUEST

EXPLODE 250

HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT
present as PARAQUAT DICHLORIDE

GROUP L HERBICIDE

Conquest Explode 250 is for the control of a wide range of grasses
and broadleaf weeds as per Directions For Use.

IMPORTANT: READ THIS BOOKLET BEFORE USE



CONQUEST CROP PROTECTION PTY. LTD
ABN 84 098 814 932
46 Hasler Road, Osborne Park, WA 6017
Telephone: 08 9347 0500, Fax 08 9347 0551

RESTRAINTS

DO NOT add wetter unless spraying at high volume. Where Explode 250 is mixed with water at less than 400 mL / 100L of water, add 100 mL Conquest Wetter 1000 per 100L of spray.

DO NO spray plants which are waterlogged, under stress of any kind or covered with soil or dust.

DO NO spray plants covered with heavy dew, but rain following spraying will not affect results.

DO NO sow or cultivate for 1 hour after spraying but operations should commence within 7 days.

For ground application only – do not use through aircraft, misting machines or hand-held ultra low volume controlled droplet applications (CDA units).

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Aid to Cultivation to minimise cultivation and prepare a clean bed for sowing.	Annual grass and broadleaf weed control. Early autumn sowing.	Qld, NSW, Vic, SA, Tas, NT, ACT only	1.2 to 1.6 L *	Where cultivation follows spraying, it may commence one hour after spraying but should be completed within 7 days. Where heavy weed growth is present at spraying a better seedbed will result if cultivation is delayed 3-5 days. Use the higher rates for dense, more mature weed stands. Wild oats must have at least two leaves. Where 'Reglone' is used the lower Conquest Explode 250 rate should be sufficient to control dense mature weeds. Pasture: Remains of old pasture should be reduced by continuous heavy grazing. Remove stock 3-5 days before spraying to allow to freshen up.
	Winter, spring and early summer sowing.		1.6 to 2.4 L	
	Wild oats at 2-5 leaf stage in autumn / winter.	Qld, Vic SA, Tas, NT only	600 mL to 800 mL	
		NSW, ACT only	600 mL	
Rice	Annual grass and broadleaf weed control.	Qld, NSW, NT only	1.6 L 800 mL	Pre-sowing. Post-sowing, pre-crop emergence.
Wild Oat control in Spring Fallows	Wild oats at 2-5 stage.	Qld, NSW, NT, ACT only	1.2 to 2 L	Use higher rate for summer growth. Avoid spraying under hot, dry conditions. Best results will be obtained when spraying is carried out in the late evening.
Kikuyu / Paspalum Pasture	To suppress growth to oversow winter seed.	Qld, NSW, ACT only	1.6 or 2.4 L	Use the high rate for February spraying and the low rate in March.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Selective Weed Control Autumn / early Winter - annual and perennial clover	Annual grass and some broadleaf weed control except Paterson's Curse, Sorrel, Dock, Shepherd's Purse and some thistles.	All States	600 mL to 1.2 L 1.2 to 1.6 L *	Use the higher rates for dense weed stands.
Late winter/ early spring - annual and Perennial clover - Cocksfoot - perennial ryegrass - Phalaris - Demeter fescue only	Control of some broadleaf weeds including Paterson's Curse, Sorrel, Dock, Shepherd's Purse and some thistles will not be achieved. Alternative methods such as the spray-graze technique with 2, 4-D or MCPA should be considered.	Qld, NSW, Vic, SA, Tas, NT, ACT only	* 1.6 to 2.4 L	Use the higher rate in winter/early spring when barley grass is present. All applications: Graze pastures continuously after the seasonal break to a height of 2-4 cm. Remove stock 2-3 days before spraying to allow weeds to freshen up. Do not apply until clover has reached the 6-leaf stage. Do not spray clovers, which are affected by insect attack, disease or moisture stress. Do not use on clover pastures growing in water repellent sands or other situations subject to moisture stress at or immediately following treatment. Poor recovery of the clover will result. Use the lower rate for cocksfoot and perennial ryegrass and the higher rate for Phalaris and Demeter fescue. The perennial grasses must be at least 12 months old at spraying. DO NOT APPLY TO MEDICS
	Yorkshire fog grass		1.2 L	Apply in early spring to reduce Yorkshire fog grass component and increase the clover and desirable grass component. Mixed pastures will be scorched initially but should show good recovery and beneficial changes in composition following spring rainfall and growth. In lower rainfall areas application in mid to late winter may be almost as effective but allow better pasture recovery. If pasture has been grazed allow for sufficient time for pasture and fog grass recovery before spraying. Apply in spray volumes of 100 to 250 L/ha, the latter for dense or tall, ungrazed pastures. Add Conquest Wetter 1000 at 120 mL/100L
Lucerne Autumn/ early winter	Annual grass and some broadleaf weeds.	Qld, Vic, SA, WA, Tas, NT only	1.2 to 1.6 L *	Use the higher rates for dense weed stands. Do not spray Lucerne stands under 12 months old. For residual weed control or if Paterson's Curse, Shepherd's Purse and some other broadleaf weeds are present add Conquest Diuron 900 WDG Herbicide at 1.1 kg or 1.9 kg.
		NSW only	1.2 L	

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Lucerne Late winter/early spring	Annual grass and some broadleaf weeds.	Qld, Vic, SA, WA, Tas, NT only	1.6 to 2.4 L *	If mintweed is present is present use Atragest 900 WDG Herbicide at 600g/ha.
		NSW, ACT only	1.2 L *	WARNING - In certain areas, an uncommon species of barley grass (<i>h. glaucum</i> - common species of barley grass is <i>H. leporinum</i>) resistant to paraquat based products has become established. It may regrow after an initial scorch by Explode 250. Where this problem is suspected use 'Fusilade' for grass weed control. If Explode 250 has been applied use 'Fusilade' at 1L/ha after regrowth but before heading.
Perennial Grass Seed Crops Cocksfoot, perennial ryegrass, Phalaris and Demeter Fescue only	Annual grass and some broadleaf weeds.	All States	600 mL to 1.2 L *	Use the low rate for Cocksfoot and perennial ryegrass and the higher rate for Phalaris and Demeter Fescue. Spray about 4 weeks after a full weed germination following the autumn break. The perennial grasses must be at least 12 months old at spraying.
Spray topping to reduce seed set Chickpeas, Faba beans, Field peas, Lentils, Lupins, Vetch	Annual ryegrass	All States	400 mL or 800 mL	As an aid in managing annual ryegrass resistance. For use on escapes from a previous herbicide application in the current crop. Spray the crop when the ryegrass is at the optimum stage, that is when the last ryegrass seed heads at the bottom of the plant have emerged and the majority are at or just past flowering (with anthers present or glumes open) but before haying off is evident - usually October to November. Use of the higher rate in these crops is usually more reliable and gives a greater reduction in seed set. Reduction in crop yield may occur especially if the crop is less advanced relative to the ryegrass that is if crops have a majority of green immature pods. The higher rate may also increase any yield reduction. In practice crop losses in excess of 25% may occur. Apply by ground boom only in 50-100L/ha. Spray with a calibrated boom spray raised to give double overlap at the level of the ryegrass seed heads. Pressures of 250-350 kPa and use of 110015 or 02 nozzles or equivalent will aid coverage.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Spray topping to reduce seed set Pastures	Grasses generally (particularly annual ryegrass).	All States	400 mL	Heavily graze paddocks during spring flush to encourage even head development. Remove stock 2-3 weeks before the anticipated maturity date of the target species. However, if this is not feasible through lack of stock it is preferable to allow the pasture to mature ungrazed. Delay spraying until the last seed-heads at the bottom of the plant have emerged and initial signs of haying off appear. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Barley grass			Manage paddocks as above. Spray after head emergence but when all seed heads are green and there is no sign of haying off. Inspect paddocks before returning stock. Provided spraying was carried out before hardening seeds are present harrow to knock seed from the heads. Do not introduce lambs into paddock until safe from risk of seed injury. If seasonal conditions favour regeneration, stock should be returned to selectively graze new shoots. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Saffron thistle	NSW, SA, ACT only		Spray after the plant begins to run to head until flowering.
Prevention of annual ryegrass toxicity	Spray top - Graze to destroy seed heads.	WA only	400 mL	Grazing management as for spray topping above. Remove stock 3-4 weeks before anticipated maturity date. Spray must be applied within 10 days after emergence of the first ryegrass seed heads. To ensure adequate control of toxin development, <u>heavy continuous grazing is essential from 1 day after spraying</u> until the pasture has completely hayed off. The required stocking rate will vary but must be sufficient to keep all regrowth after spraying completely eaten off to prevent further growth producing new seed heads which could become toxic.
Hay Freezing	Maximum retention of protein in standing dry feed.	All States	800 mL	Graze paddocks as for spray topping above. Remove 3-4 weeks before the anticipated maturity date. Apply prior to commencement of haying off regardless of the grass species involved. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
General Weed control Hops	Annual grasses	VIC, TAS only	* 1.2 to 1.6 L plus 1.1 kg/ha Simaquest 900 WDG and/or 750 mL to 1.4L/ha Reglone	Apply as a directed inter-row spray prior to crop emergence from winter dormancy, using a minimum of 250L/ha spray volume to ensure good and even coverage of weeds.
Orchards (including bananas) Vineyards	Annual weed control	QLD, VIC, SA, WA, TAS, NT only	1.6 to 3.2 L/ sprayed ha *+ 160 to 320 mL per 100L (a) see below	<p>Spray as necessary for control of annual weeds. Avoid contacting crop foliage. In bananas apply soon after weed emergence and before weeds reach 15cm in height. Use spraying pressure less than 240kPa. Avoid chemical contact with roots and peepers near the pseudo stem. Repeat sprays as required. Explode 250 will not harm trees or vines with mature brown bark if this alone is sprayed. Use the higher rate for dense weed growth.</p> <p>If fat hen <i>Chenopodium album</i> or <i>Portulaca</i> spp. are present and Explode 250 rate is less than the ratio 800 mL/100L add 200 mL Conquest Wetter 1000 per 100L of spray mix.</p> <p>Note: Spot spray rate assumes 1000L water/ha. For lower water volumes increase dilution rate as below: Water volume 250L/ha: use 640 to 1280 mL/100L Water volume 500L/ha: use 320 to 640 mL/100L Water volume 750L/ha: use 210 to 430 mL/100L</p> <p>OR Measure how much spray is required to cover an area of 100 square metres using your normal application volume. Your dilution rate is 16 to 32 mL of Explode 250 in this volume.</p>
		NSW only	1.7L/ sprayed ha * +	

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Peanuts Post-emergence (in-crop)	<i>Datura</i> spp. (2-4 leaf) Annual ground cherry (2-3 leaf) Apple-of-Peru (2-4 leaf) Milkweed (2-3 leaf) Stagger weed (2-3 leaf) Blue heliotrope (2-3 leaf) Wandering Jew (2-3 leaf) Anoda weed (2-4 leaf) Bellvine (2-3 leaf) Common morning glory (2 leaf)	QLD, NT only	400 mL 600 mL 600 mL 600 mL 800 mL 800 mL 800 mL 800 mL 1 L 1 L	Spray peanuts up to 7-8 leaf stage but before majority of plants flowering. Foliage will be scorched following application but plants recover rapidly. Apply in 200-250L/ha for thorough coverage of weed foliage. A dense canopy of weeds may reduce weed control due to shielding. Conquest Wetter 1000/100L of spray mix. Do not spray on peanuts under extremely hot dry conditions when peanuts are very small. In environments such as Far North Queensland use the lower rates in the range.
Potatoes	General weed control (in-crop)	All States	1.2 to 1.6 L *	Spray at early crop emergence (no later than 25% emergence of potato shoots). Use the higher rate for dense weed growth.
	Pre-harvest weed control		2.8 L *	Spray about one week before digging and after tops have died down.
Row Crops, Vegetables and Market Gardens	Pre-planting and pre-crop emergence		1.2 to 1.6 L OR 200 mL/100 L * +	To control weeds in seedbeds. Treat no less than three days before sowing or before crop emergence. Use the lower rate for early autumn applications.
	Post-emergence inter-row weed control		1.2 to 1.6 L OR 200 mL/100 L * +	Apply after crop seedlings have emerged or when transplanted crops are established. Direct the spray so that it does not touch the crop. Use shielded nozzles.
	Seedling weeds		*2.4 or	Seedling weeds - use the lower rate for early autumn applications.
	Older weeds		400 mL/100 L	More mature stages of weed growth.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Sugar Cane (Plant and ratoon)	Grass and some broadleaf weeds	QLD, NSW, NT only	Up to 5 cm high	Apply as a broadcast spray over-the-top of plant cane up to the 3-4 leaf stage. Cane foliage will be scorched but new leaves will appear in 7-10 days.
	Enhancement with Conquest Diuron 900 WG Herbicide		1.2 to 1.6 L per sprayed ha + 1.2 to 1.6 L / + 275 g to 500 g Conquest Diuron 900 WG Herbicide Up to 10 cm high 1.2 to 1.6 L + 1 kg Conquest Diuron 900 WG Herbicide 1.6 L + 2.8 to 3.9 kg Conquest Diuron 900 WG Herbicide	In plant cane between the 3-4 leaf stage and the formation of the true stem use a directed interspace spray. The Irvin spray boom (or other similar equipment) is the most suitable equipment to avoid excessive drift onto cane foliage while spraying at the cane bases of plant and ratoon cane. After the formation of the true stem, which is resistant to Explode 250, the sprayer height can be raised to overlap the spray pattern to give weed control in the stool. Use the higher rate for dense, more mature weeds. Explode 250 can be mixed with Atracquest 900 WG herbicide to give residual weed control when used as a blanket or directed spray – refer to the Atracquest 900 label for specific rates. It may also be mixed with Conquest Diuron 900 WG Herbicide at 2.8 to 3.9 kg/ha for residual weed control.+ To enhance activity of Explode 250 under favourable growing and in open sunny conditions add 135g to 275g/ha Conquest Diuron 900 WG Herbicide at the rates shown for weed size. Conquest Diuron 900 WG Herbicide at rates of up to 500g/ha can be blanket sprayed. Use a directed spray for higher rates of Conquest Diuron 900 WG Herbicide. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250 L spray solution/ha, increase to 350 L/ha for weeds up to 10 cm high. Use a minimum spray volume of 400 L/ha for weeds greater than 10 cm high and for dense mature weeds. Always add Agral at 200 mL or Conquest Wetter 1000 at 120 mL per 100L of water.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Non-Agricultural situations, around sheds, roadways, paths	Annual weed control	All States	1.6 to 4 L/ha OR 200 mL/100 L * +	Spray to thoroughly wet weed growth. Explode 250 can be combined with soil residual herbicides Conquest Diuron 900 WG Herbicide, Simaquest 900 WG Herbicide or Atragest 900 WG Herbicide to give rapid knockdown and prolonged weed control. Use the higher rate for dense weed growth.
	Columbus grass	NSW only	+ Spot spraying 160 mL/ 100 L plus 1 L 'Frenock' Boom-spray 2.3 to 4.5L/ha plus 12 to 22L 'Frenock'	
Firebreaks	Knock down weed growth to eliminate fire hazard or assist firebreak burn	All States	1.6 L to 4 L	<p>Apply mid-winter to early summer.</p> <p>Use the higher rate for dense weed growth. After desiccation is complete the sprayed area may be burnt (normally 7–10 days after spraying).</p> <p>Explode 250 can be combined with soil residual herbicides Atragest 900 WG Herbicide, Conquest Diuron 900 WG Herbicide or Simaquest 900 WG Herbicide to give rapid knockdown and prolonged weed control</p>

* Capeweed or Erodium spp. present: Add 'Reglone' at 750 mL to 1.5L/ha (125 mL to 25 mL/100L for high volume spraying). Use higher rate for plants more than 10 cm diameter.
+ If Explode 250 rate is less than the ratio 400 mL/100L add 100 mL 'Agral' or 60 mL Maxiwet 1000 per 100L of spray mix.

Wetting Agent: (a) Add 170mL Agral® or 100mL BS 1000 per 100L

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE, THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR AT LEAST 1 DAY, OR GRAZE HORSES FOR 7 DAYS AFTER APPLICATION.
REMOVE STOCK FROM TREATED AREAS 3 DAYS BEFORE SLAUGHTER.
CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS AND VETCH – DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

PLEASE NOTE EXTRA WETTER REQUIREMENTS FOR HIGH VOLUME SPRAYING.

GENERAL INSTRUCTIONS**Resistant Weeds Warning****GROUP L HERBICIDE**

Conquest Explode 250 Herbicide is a member of the bipyrldyls group of herbicides. Explode 250 has the inhibitor of photosynthesis at photosystem I mode of action. For weed resistance management Explode 250 is a Group L herbicide.

Some naturally occurring weed biotypes resistant to Explode 250 and other inhibitor of photosynthesis at photosystem I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Explode 250 or other inhibitor of photosynthesis at photosystem I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of Explode 250 to control resistant weeds.

This product kills annual grasses and most annual broadleaf weeds (excluding capeweed) in specified situations and should not be used for any other purpose. Quickly kills green plant tissue on contact. Is immediately inactivated in the soil or heavy dew. The principle of selective weed control with this product is that annual weeds are killed but perennial plants and clovers recover after an initial scorch. The control of annual weeds by spraying with this product will allow the desirable perennial species to thicken up at the expense of the weeds. Moisture and fertility should not be limiting at spraying and the proportion of desirable species must be great enough for them to fill in the areas previously occupied by weeds. Long-term weed control can be obtained following the quick knockdown given by this product if it is combined with soil residual chemicals.

- 1. DO NOT** use hand-held ultra low volume controlled droplet applicators (CDA units), boomless jets or misting machines (except in banana plantations).

2. Mixing

Add the required quantity of product to water in the spray tank and agitate to give even mixing. Agitate again if left standing.

3. Wetting agent

This product contains a wetting agent and additional wetter is not required unless high volume spraying results in excessive dilution of wetter content. This will occur when product rates fall below 400 mL per 100L of spray. Under such circumstances wetter should be added at the rate of 100 mL of Conquest Wetter 1000 per 100L of spray mix.

Where Fat Hen or Portulaca are present in orchard or vineyard situations, extra wetter should be used when this product ratio is less than 800 mL per 100L. Add wetter at double the above recommendations. Do not use alkaline or anionic wetting agents.

4. Clean water

Mix this product with clean water only. Water should be clean and free from clay, silt and algae. Providing it meets this requirement, saline water, water collected from roofs, bore water, dam water and water from creeks may be used.

(i) Cereals and Broadacre Spraying

Use only through a properly calibrated boom spray which should be fitted with flat fan jets and adjusted to a height to give at least double overlap of the spray at the top of the weeds being sprayed. Spraying pressures should be in the range of 200-300 kPa. Speed of travel should be in the range of 6 – 15 km/hr. It is essential that a good marking system be used. If a disc marker is used, it must be mounted so as to turn the soil back on to the area sprayed. It is essential to obtain good leaf coverage with the spray and volumes of dilute spray must be adjusted according to density of weed growth. 100L/ha may be used for seedlings or well grazed weeds up to 2 cm high. For plant height 2-5 cm use 150L/ha and up to 6-10 cm use 200L/ha. Spray volumes may be as low as 50L/ha (30L/ha in WA) for weed growth below 5 cm high, or for spray topping and hay freezing. Equipment must be appropriate to this volume, properly calibrated and fitted with spraying tips designed to give droplets in the 200-250µm Volume Median Diameter range.

(ii) High Volume Application

Higher volumes will generally be required to give good coverage of weed growth in situations other than those specified under cereals and other broadacre crops.

(iii) Wash spray equipment with clean water immediately after use. This product is highly corrosive to metals, particularly galvanised iron and aluminium and should not be left for long periods in tanks or equipment made of these materials.**6 Compatibility**

This product combines satisfactorily with the soil active herbicides Atracquest 900 WG, Conquest Diuron 900 WG and Simaquest 900 WG where prolonged weed control is required as well as a quick knockdown. This product is compatible with Conquest Wetter 1000, 'Reglone', 'Sprayseed', 'Banvel' 200, 'Banvel' M, MCPA Amine (no more than 1L per 800 mL Conquest Explode 250 Herbicide), Conquest Chlorsulfuron 750, 'Yield', 'Avadex' and Conquest Tricon 480.

7 Spraying conditions

Avoid spraying plants under stress from waterlogging, frost, drought etc. or covered with dust and soil. Results will be better if application is made in dull weather or at the end of the day. Light rain following spraying will not affect results. Avoid drift into neighbouring crops.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto susceptible plants/crops, cropping lands or pastures. This formulation should not be applied on or near water which is used for irrigation purposes.

PROTECTION OF LIVESTOCK

Domestic pets and poultry – keep away from treated areas. This formulation should not be applied on or near water which is used for livestock watering.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container. This formulation should not be applied on or near water which is used for human consumption, livestock watering or irrigation purposes or water used for commercial or recreational fishing.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.

SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. Obtain an emergency supply of Ipecac Syrup APF. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION

Do not work in spray mist.

Do not continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

Where there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.

Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID

If poisoning occurs get to a doctor or hospital quickly (Tel: 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

NOTE TO PHYSICIANS

For additional advice on the treatment of paraquat poisoning, please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors" (available from Conquest Crop Protection Pty Ltd).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.



APVMA Approval No:	57817
Application Number:	50027
File Name:	57817_50027_Booklet_V1.pdf
Product Name:	Conquest Explode 250 Herbicide
Pack Size:	Booklet
Dimensions:	170mm x 120mm
Print Size:	A4 - 100%