

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bioclear Paraffinic Oil

ACTIVE CONSTITUENT: 823 g/L PARAFFINIC OIL
(as 1888-1976)



UNSULPHONATED RESIDUE NO LESS THAN 98.0%

**A NARROW RANGE EMULSIFIABLE PARAFFINIC OIL FOR THE
MANAGEMENT OF PESTS AND DISEASES AS PER DIRECTIONS
FOR USE**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS
PRODUCT**

CONTENTS: 20L, 205L, 1000L

Caltex Australia Petroleum Pty Ltd
ACN 000 032 128
2 Market Street
SYDNEY NSW 2000

STORAGE AND DISPOSAL

Store in the original container, tightly closed in a safe well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple rinse container for disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.

For refillable containers (1000L) - empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and may irritate the skin. Avoid contact with the eyes and skin. When opening the container and preparing the spray, wear elbow-length PVC gloves, face shield or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre
(Phone Australia: 13 11 26).

If swallowed do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) which is available from the supplier.

NOTICE TO BUYER

Caltex accepts no responsibility for any consequence whatsoever arising from the use of the product other than used in accordance with this label.

CALTEX TECHNICAL ADVICE LINE: 1300 364 169

APVMA APPROVAL NO: 58360/62425

BATCH NO:

DATE OF MANUFACTURE:

CAUTION
KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bioclear Paraffinic Oil

ACTIVE CONSTITUENT: 823 g/L PARAFFINIC OIL
(as 1888-1976)

UNSULPHONATED RESIDUE NO LESS THAN 98.0%

**A NARROW RANGE EMULSIFIABLE PARAFFINIC OIL FOR THE
MANAGEMENT OF PESTS AND DISEASES AS PER DIRECTIONS
FOR USE**

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA APPROVAL NO: 58360/62425

Caltex Australia Petroleum Pty Ltd
ACN 000 032 128
2 Market Street
SYDNEY NSW 2000

DIRECTIONS FOR USE:

RESTRAINTS

DO NOT spray when temperatures exceed 32°C, (except where noted for Citrus) or when soil is dry and trees are suffering from moisture stress.

Crop	Pest	State	Application Rate	Critical Comments
Litres of Oil in Dilute Spray per 100L Water				
Apples	Mites (Eggs), including European Red Mite, Scales (Hard & Soft)	NSW, ACT, Vic, SA, WA, TAs only	2L	Dormant or delayed dormant to half green tip
		Qld	3L	
	Mites	NSW, ACT, VIC, SA, WA, Tas only	1L	Summer (Foliar or Cover) or Post Harvest
Bananas	Cercospora Leaf Spot	NSW, ACT	4L /ha	Apply in combination with mancozeb at 2.2-4.5 kg/ha in 200 to 500L of water. Apply at 3-4 week intervals, from December until late April. In North Queensland the interval may be reduced to 10-14 days in the wet season.
		Qld	3.5L /ha	
	Cercospora Leaf Spot, Leaf Speckle	NSW, ACT, Sth Qld, WA	5L /ha	Apply in combination with Tilt* 250 EC. Commence spraying in December and apply 4 to 5 times at 21 day intervals. Later use alternative fungicides if needed. Spray both sides of leaves.
	Cercospora Leaf Spot, Leaf Speckle, Cordana Leaf Spot	Nth Qld	5L /ha	Apply in combination with Tilt 250 EC. Spray every two weeks from the start of the wet season in December and make a maximum of 6 applications. Later use alternative fungicides if needed. Spray both sides of leaves.
Cotton	Defoliation Aid	NSW, ACT, Qld, WA	2 L/ha	Apply in combination with thidiazuron (eg DROPP*) defoliant
	Carrier for certain ULV insecticides	NSW, Qld, only	3-5 L/ha	Apply in combination with ULV insecticides as specified on the insecticide label to ensure insecticide reaches it target.
Wetting agent/adjuvant/ carrier as recommended on fungicide and insecticide labels	Refer to particular pesticide label	All	500mL – 5L /ha	Bioclear acts as a carrier/adjuvant/ wetting agent which may improve the performance of certain pesticides. Always check pesticide label for correct rates.

Crop	Pest	State	Application Rate	Critical Comments
Litres of Oil in Dilute Spray per 100L Water				
Pears	European Red Mite, Eggs, Scales	NSW, ACT, Vic, SA, WA	1-2L	Dormant or delayed dormant (up to and including petal fall)
		Tas Only	2L	
		Qld	3L	
	Two Spotted Spider Mite	NSW, ACT, Vic, SA, WA, Tas only	1L	Summer (Foliar or Cover) or Post Harvest
	Brown Mite, Pear Rust Mite	NSW, ACT, Vic, SA, WA, Tas only	1-2L	
	Pear Leaf Blister Mite	NSW, ACT, Vic, SA, Tas, WA only	1.5-2L	
Apricots Cherries	San Jose Scale	NSW, ACT, Vic, SA, WA only	2L	Dormant
		Qld	3L	
	Mites and Scales	NSW, ACT, VIC, SA, Tas Only	1-1.5L	Summer (Foliar or Cover) or Post harvest. Applications should not be made over oil sensitive varieties. Do not apply to trees lacking adequate moisture.
Almonds Peaches Nectarines	Aphids (Eggs), Mites (Eggs)	NSW, ACT, Vic, SA, WA, Tas Only	3L	Dormant or delayed dormant
	San Jose Scales	NSW, ACT, Vic, SA, WA only	2L	Dormant
	Mites and Scales	NSW, ACT, Vic, SA, WA only	1-2L	Summer (Foliar or Cover) Post Harvest
Passionfruit	Red Scale, Hemispherical Scale Suppression of: Passionvine Mealy Bug	All States	500mL – 1L	Please refer to the General Instructions under "Crop Safety – Passionfruit" for full details on crop safety. Apply via a low volume rate of 1000L per hectare. A minimum interval of 14 days should elapse between sprays in the first month with additional applications applied at monthly intervals for up to 6 months. Good coverage is essential for Red Scale.

Plums Prunes	Aphids (Eggs) Scales Mites (Eggs)	NSW, ACT, SA, WA, Tas Only	1.5-2L	Dormant or delayed dormant. To half green tip.
	Mites Scales	NSW, ACT, VIC, SA, WA, Tas Only	1-1.5L	Summer (Foliar or Cover) or Post Harvest. Do not apply over fresh market fruits after bloom starts to form as the oil will remove the bloom. Applications should not be made over sensitive varieties.

Crop	Pest	State	Application Rate	Critical Comments
Litres of Oil in Dilute Spray per 100L Water				
Pecans	Aphids (Eggs) Scales Mites (Eggs)	NSW, ACT, Qld, SA, WA, Tas Only	3L	Dormant
	Aphids Mites	NSW, ACT, QLD, SA, WA, Tas only	1-1.5L	Summer (Foliar or Cover) or Post Harvest
Avocado	Scale Insects	Qld	1.2L to 1.6 L	Apply when heavy scale populations occur on stem, foliage or fruit. Do not apply if trees need watering. Application is most effective against young crawler stages.
		NSW, ACT, WA	2L	
Kiwi Fruit	Scale Insects	Qld	1.0L to 1.6L	
		NSW, ACT, WA	2L	
Custard Apple	Scale Insects	Qld	1.0 to 1.7L	
		NSW, ACT, WA	2L	
Banana	Scale Insects	Qld, NSW, ACT, WA	2L	
Mango	Scale Insects	Qld	1.25L	
		WA	2L	
Shade Trees* and Shrubs including Conifers, Deciduous Broadleaf Evergreens and Woody Ornamentals, Ornamental Trees* Shrubs along city streets, other rights of way including Deciduous and Broadleaf Evergreens	Aphids Scales (Immature) Spider Mites Whitefly (immature)	NSW, ACT, VIC, SA, WA, Tas only	2-4L (Winter) 1-3L (Summer)	Winter dormant period as needed for plants that become dormant in winter. Summer foliage (Foliar or Cover) as needed

Flowers and Foliage Plants including Roses and other Flowering Shrubs, Foliage Ornamentals and Bedding Plants			2-3L (Winter) 1-2L (Summer)	Winter Dormant as needed for plants that become dormant in Winter. Summer foliage (Foliar or Cover) as needed.
Christmas Trees	Aphids Scales (Soft & Hard) (Immature) Spider Mites		2-4L (Winter) 1-3L (Summer)	Winter Summer

CAUTION: Spray no more than 4 times during growing season with two weeks minimum application interval. Do not spray when buds are fully opened and shoot elongation is occurring. Do not spray when there is no obvious moisture in the leaf or the plant is under stress. Avoid spraying open blooms. Bleaching and spotting has been observed with open blooms of certain plants.

Oil removes the glaucous (blue) bloom from some evergreens. Always use lower dosage or test spray oil sensitive plants first – do not spray walnut foliage.

Crop	Pest	State	Application Rate	Critical Comments
Litres of Oil in Dilute Spray per 100L Water				
Blueberries	Mites Scales	All States	2L	Foliage
			3L	Dormant
Strawberries	Aphids Mites	NSW, ACT, SA, WA, Tas only	1L	Dormant and Foliage
Grapes	Mealybugs	NSW, ACT, Qld, SA, WA, Tas only	4-6 Litres with 306 to 460 Litres Water per Hectare	Dormant – A tank mix using insecticides improves control Air carrier or Air blast type sprayers strongly recommended to avoid plant injury and reduce effect on bloom of table grapes.
	Grapevine Scale	NSW, Vic, SA, WA, Qld Only	3L	Apply as a post pruning spray in mid- winter when vines are fully dormant.
		Tas Only	2L	
Asparagus Beans Beet Corn Cucurbits Peppers / Capsicums Radish Squash Tomatoes	Aphids Mites Thrips Leafhopper	All States (Except Qld for beans, tomato, corn, and peppers/ capsicums)	1-2L	As needed
Corn (Sweet and Field) Sugar Beet	Aphids Mites Whitefly Bugs (immature)	NSW, ACT, SA, Tas, WA only	2L	As needed
Chrysanthemum Differnbachia Dracaena Ferns Gardenias Jade Plant Palms Philodendron	Aphids Scales Whitefly	All States	1L-2L	Do not apply to plants in direct sunlight behind glass. Do not use on Coconut Palms and Maidenhair Ferns. Chrysanthemum blooms have shown phytotoxic symptoms at the higher rate. Applicator should conduct a test for phytotoxicity by treating a few specimens before making a large- scale application. Protect floor, floor coverings and furnishings from overspray.
			2.5mL- 5.0mL/L	

Crop	Pest	State	Application Rate	Critical Comments
Rate of Oil in Dilute Water Spray				
Azalea Begonia Camellia Chrysanthemum Crown of Thorns Diffenbachia*	Aphids Mealybugs Scale (Soft and Hard) Thrips Whitefly (Immature) Reduces number of Spider mites	All States	10 – 20mL/1L or 1L-2L/100L	Do not apply to plants in direct sunlight behind glass. Do not use on Coconut Palms and Maidenhair Ferns, Chrysanthemum blooms have shown phytotoxicity symptoms at the higher rate. Applicator should conduct a test for phytotoxicity by treating a few specimens before making a large-scale application. For greenhouse pests listed use once a week initially then as the pest is controlled decrease the frequency to every 2-3 weeks as needed
			10mL/1L OR 1L/100L	
Easter Lilly Fern Gardenia Hibiscus Foliage Jade Plant Palm Philodendron			10 – 20mL/1L or 1L-2L/100L	
Poinsettia			10mL/1L OR 1L/100L	
Reiger Begonia Zinnias			10 – 20mL/1L or 1L-2L/100L	
Hardy Plants			10mL/1L or 1L/100L	
	Leaf Polish	All States		

- Oil removes the glaucous (blue) bloom from some evergreens. Always use lower dosage or test spray oil sensitive plants first – do not spray walnut foliage.

This table shows rates for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.				CRITICAL COMMENTS
CROP	PEST/DISEASE	STATE	RATE/100 L	For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. For concentrate spraying, DO NOT use at rates greater than 5 times the dilute spraying rate.
CITRUS	Citrus Leafminer	NSW, Qld, Vic, SA, WA, Tas only	250 to 500 mL for control of citrus leafminer 500 mL to 1 L if application coincides with need to control pests such as red scale	Spray every 5 to 14 days during each flush cycle; sprays will be required more frequently during Summer and early Autumn than in mid-to-late Autumn.
	Red Scale	Vic, SA, WA, NSW only	0.5 to 1.0 L	Apply in late Nov/early Dec and again in late Feb/early Mar if required.
			0.5 to 1.0 L + 50 mL Lorsban 500EC Insecticide	Apply between Nov-Mar. A second application may be necessary for heavy infestations.
		Qld only	0.5 to 0.6 L	Apply in late Nov/early Dec and in late Jan/early Feb if required.
			0.5 L + 100 mL Lorsban 500EC Insecticide	Apply between Nov-Mar. A second application may be necessary for heavy infestations.
	White Wax Scale, Pink Wax Scale	Vic, WA, NSW only	0.5 to 1.0 L	Apply in Dec/early Jan.
		Qld only	0.5 to 0.6 L	Apply in late Nov/early Dec and in late Jan/early Feb if required.
		Vic, Tas, SA only	1.0 L + 100 mL carbaryl	Apply in accordance with registered carbaryl insecticide label.
		NSW, WA only	1.0 L + 140 mL carbaryl	
		Qld only	0.6L + 140mL carbaryl	
	Circular Black Scale	Qld only	0.5 to 0.6L	Apply in late Nov/early Dec and in late Jan/early Feb if required.
		WA only	0.5 to 1.0L	
	Black Scale	Tas, WA only		Apply mid-Dec/early Jan and in Feb/Mar.
	Soft Brown Scale	Qld only	0.5 to 0.6 L	Apply in late Nov/early Dec and again in February/March.
		WA only	0.5 L to 1.0 L	
	Broad mite	All states	250 mL + 15 or 25 mL Vertimec ² Miticide/Insecticide	Apply in accordance with the Vertimec ² Miticide/Insecticide label.
	Brown citrus rust mite		0.7 to 1.0L	Apply Nov/Dec and again in Feb/Mar if required.
	Citrus red mite	NSW only		Apply Aug/Sep and again in Mar/Apr if required.
	Citrus rust mite	All states	250 mL + 15 or 25 mL Vertimec ² Miticide/Insecticide	Apply in accordance with the Vertimec ² Miticide/Insecticide label.
		Qld Only	0.7 to 1.0L	Apply Nov/Dec and again in Feb/Mar if required.
		NSW, Vic, SA, WA only		Apply in Feb/Mar.

CROP	PEST/ DISEASE	STATE	RATE/100 L	CRITICAL COMMENTS
CITRUS	Black Spot	All States	600 mL + copper oxychloride	Apply at half to ¾ petal fall. Consult copper fungicide label for suitable local rate and intervals for repeat application. Ensure thorough coverage is achieved throughout canopy.
		Qld only	600 mL + 200 g Dithane Rainshield Fungicide or Penncozeb 750DF Fungicide	Apply in accordance with registered Dithane Rainshield Fungicide or Penncozeb 750DF Fungicide label.
		NSW, Tas, Vic & SA only	250 mL + 200 g Dithane Rainshield Fungicide or Penncozeb 750DF Fungicide	Apply in accordance with registered Dithane Rainshield Fungicide or Penncozeb 750DF Fungicide label.
	Melanose, Smoky Blotch, Scab	All States	600 mL + copper oxychloride	Apply in accordance with registered copper oxychloride label. Ensure thorough coverage is achieved throughout canopy.
	Brown Spot in Mandarins	Qld only	600 mL + 375 g copper oxychloride + 500 g zinc sulphate heptahydrate + 140 g caustic soda	Apply in accordance with registered copper oxychloride label.
			625 mL + 200 to 300 g copper oxide	Apply in accordance with registered copper oxide label.
			625 mL + 200 to 300 g copper oxide	
	Brown Rot			

Crop	Pest	State	Application Rate	Critical Comments
Wetting Agent if recommended on Herbicide label	Weeds as specified on herbicide label	All States	250-500 mL/100L 2L/ha	Bioclear Paraffinic Oil may be used as a crop oil in a spray mixture with selective herbicides to improve their wetting and penetrating properties.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

GENERAL INSTRUCTIONS:

This product controls Aphids, Lace Bugs, leaf Hoppers, Mealybugs, Mites, Plant Bugs, Scale, Whitefly, eggs of Aphids & Mites on vegetables, Fruit Trees, Nut Trees, certain field crops, Shrubs, Plantation Trees, Greenhouse Plants, Ornamental Foliage Plants and Flowers.

Bioclear Paraffinic Oil can be used in combination with certain fungicides, herbicides and insecticides as a wetting agent/adjuvant/carrier or as specified on the products label.

General

Different plant stresses can have cumulative effect. In general, follow good farming practice to minimise all other sources of plant stress. Avoid applying Bioclear when environmental conditions elevate plant stress. AVOID spraying complex tank mixtures. In particular DO NOT spray:

- when the soil is dry
- if trees are suffering from moisture stress
- if hot weather (> 35°C) is imminent or has occurred in the previous 4 days.
- if the shade temperature exceeds 32°C
- in slow drying conditions

Since Bioclear can be used as a herbicide adjuvant, any phytotoxic tendencies of other insecticides mixed with Bioclear may be enhanced. Follow the label precautions on the other insecticides carefully and if in doubt about a certain mixture then test on a small area first before widespread use.

Bioclear sprays greater than 250 mL/100L water should not be applied for at least one month after spraying with sulphur or lime sulphur, as severe injury may occur, especially if the oil is applied in hot weather.

DO NOT apply Bioclear in combination with Propargite or within 4 weeks of a Propargite application.

Adding spraying oils with TORQUE^{*1} Miticide/Insecticide can reduce the speed of action of Torque^{*1}

Miticide/Insecticide. Avoid applying Torque^{*1} with Bioclear where rapid speed of action is required.

DO NOT apply Bioclear in combination with, or following a chlorothalonil application.

DO NOT use in combination with Supracide^{*2} 400 Emulsifiable Concentrate Insecticide on mangoes after fruit set and while fruit is on the tree.

CITRUS

Application

Bioclear is an effective insecticide when used alone to control light-moderate infestations of pests as indicated in the directions for use table. When a range of Bioclear rates is indicated, use the lower rate for light pest infestations and the high rate for moderate to heavy infestations, but only if weather and tree conditions are favourable (see general instructions – precautions). Alternatively for heavy pressures, apply low rate 2-3 times at an interval of about a fortnight. DO NOT exceed maximum recommended annual dosage (see precautions section.) Time applications to target crawler stages of pests. Apply when trees are in good condition 7-10 days after rain or irrigation. Apply sprays thoroughly to ensure complete coverage is achieved on both sides of susceptible leaves throughout whole canopy. The preferred type of sprayer for large trees is the oscillating boom with outrigger. As a general guide, 4m high trees with heavy infestations of armoured scale could require spray volumes from 10 000 to 14 000 L/ha, soft scales 8 000 to 10 000 L/ha, and mites 5 000 to 8 000 L/ha. Adjust

volume proportionally for tree size, planting density, target foliage (eg flush growth or whole canopy), canopy density and degree of infestation.

When pest pressure is moderate to heavy, and/or your spray equipment cannot achieve coverage comparable to an oscillating boom with outrigger, consider using Bioclear in one of the mixtures indicated in the directions-for-use table. However, achieving thorough coverage should always be the aim no matter what mixture is used.

Precautions - Citrus

Refer to Precautions – General section above.

DO NOT use rates higher than 0.6% in Queensland. Use high rate (1%) in southern Australia ONLY when all general spraying precautions are met (see PRECAUTIONS – General) and when pest pressure warrants. Always consider the low rate as the first spray option.

DO NOT apply when high rate of natural fruit drop is expected.

DO NOT apply within 4 weeks of harvest.

DO NOT apply Lorsban 500EC Insecticide in IPM programs. In SA, do not spray Bioclear with Lorsban 500EC Insecticide if spraying in November/December.

DO NOT apply to Glen Retreat variety of mandarins.

If petroleum spray oil is applied more than once from Spring to Autumn, either alone or in combination with other materials, the total strength of oil used in commercial orchards should not exceed 3L of oil in Queensland, and 2L of oil in southern Australia.

DO NOT apply Bioclear with Supracide[®] 400 Emulsifiable Concentrate Insecticide to any citrus.

DO NOT apply Bioclear in combination with Torque[®]1 to Torque[®]1 susceptible varieties of citrus such as tangerines, tangelos and some varieties of grapefruit.

DO NOT apply copper fungicides when slow drying conditions prevail or if it is likely to rain before spray is dry.

DO NOT use this product with copper fungicides on copper shy varieties.

MIXING INSTRUCTIONS:

1. Add water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides.
 - a. If wettable powder formulation – mix water and powder thoroughly so that power is totally suspended in the water before the oil is added.
 - b. If other pesticide to be added is an emulsifiable formulation, do so after the oil and water has been thoroughly mixed.
3. Add oil under agitation when tank is $\frac{3}{4}$ full. Top off with water to form a milky solution.
4. Maintain agitation until solution is completely used.
5. In a small equipment lacking agitators, stir or shake diluted spray frequently during applications.
6. It is important for users to read and follow all instruction on the label of the proposed tank mixed products.
7. Flush fluids in sprayer hose lines back into tank reservoir if fluid is allowed to stand for more than 20 minutes.

APPLICATION INSTRUCTIONS

- The target must be completely covered in spray.
- Oil residue on the plants surface often acts as a feeding and oviposition deterrent. However, the primary target is the pest itself as Bioclear Paraffinic Oil is a contact pesticide.
- Dilute applications. (Greater than 1400L spray per hectare in horticultural crops) in most cases ensure best coverage.
- Speed of travel is extremely important, tractor speed from 1.6kph to 5.4 kph is recommended, depending on crop size and target pest.
- Do not spray when shade temperatures are or exceed 32°C

COMPATIBILITY

- Bioclear Paraffinic Oil can be used with such pesticides as NPV's, Bt's, Pyrethroids, Abamectin, Tracer II*, Mancozeb and Follicur* as stated on their respective directions for use labels.
- Do not spray oil with Dimethoate, or any other product containing sulphur.
- If possible, either keep the spray equipment used for these compounds separate from the equipment used for Bioclear Paraffinic Oil, or make sure that the sprayer is thoroughly cleaned, so that no residue from these compounds remain.
- Do not use with Carbaryl on deciduous fruit trees.

TANK MIXTURES

- This product may be mixed with other pesticides to improve the level of kill or enhance coverage. User should read and follow all instructions on the label of the proposed tank mixed product.

TIMING THE TREATMENT

- Applicator must determine the precise timing to fit local growth and climate conditions. Do not exceed maximum rates or apply more often than recommended.
- May be used up to one day before harvest.

CROP SAFETY – PASSIONFRUIT

The sensitivity of this product has not been tested on all Passionfruit varieties, therefore it is recommended that a small number of plants be treated in the first instance before treating the whole crop.

DO NOT exceed an application rate of 0.5% or 4 applications per year for oil sensitive varieties such as those from the Panama line (Passionfruit only).

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish. Do not contaminate streams, rivers, waterways with chemicals or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do not apply pesticides or other chemicals incompatible with oil, within 4 weeks before or after application of any other spray oils, as crop damage may occur.

STORAGE AND DISPOSAL

Store in the original container, tightly closed in a safe well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple rinse container for disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.

For refillable containers (1000L) - empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and may irritate the skin. Avoid contact with the eyes and skin. When opening the container and preparing the spray, wear elbow-length PVC gloves, face shield or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre
(Phone Australia: 13 11 26).

If swallowed do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) which is available from the supplier.

NOTICE TO BUYER

Caltex accepts no responsibility for any consequence whatsoever arising from the use of the product other than used in accordance with this label.

CALTEX TECHNICAL ADVICE LINE: 1-300 364 169

* Not Caltex Trademarks