

Product Name: HITMAN ORGANIC INSECTICIDE & FUNGICIDE  
APVMA Approval No: 58472/141574



Label Name:	HITMAN ORGANIC INSECTICIDE & FUNGICIDE
Signal Headings:	READ SAFETY DIRECTIONS
Constituent Statements:	ACTIVE CONSTITUENT: 285 g/L POTASSIUM SALTS OF FATTY ACIDS
Mode of Action:	
Statement of Claims:	For the control of insect pests on broadacre crops, fruit trees, vegetables, nut crops and nurseries and suppression of botrytis cinerea in grapevines as per the directions for use table.
Net Contents:	5 L - 1000 L
Restrains:	
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	GRAPEVINES: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION OTHER SITUATIONS: NOT REQUIRED WHEN USED AS DIRECTED
Trade Advice:	
General Instructions:	This section contains file attachment.

Resistance Warning:	
Precautions:	
Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
Storage and Disposal:	Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Safety Directions:	SAFETY DIRECTIONS: Wash hands after use.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.
First Aid Warnings:	

## DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
Outdoor Crops, Vegetables, Fruit trees, Pot plants and Ornamentals	Aphids Thrips Mealybug Two Spotted Mite Spider-mite Whitefly	1.5 - 3.0L in 100L of water	This product may not be suitable for delicate ferns, mosses, flowers and plants under stress. Use the lower rate in integrated control programs and on tender foliage.
Grapevines	Suppression of grey mould (Botrytis cinerea)	13.6L per ha in 500 to 1000 litres water	Apply by high volume spraying to the point of run off. Ensure thorough coverage of all parts of treated plants. Higher water rates will give better coverage and better botrytis suppression. Apply up to four applications in a season from early flowering until 14 days before harvest.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## GENERAL INSTRUCTIONS

As insecticide:

Spray when insects are noticed. Spray insects directly and repeat as necessary. A complete coverage of the target is necessary. Apply to the point of run-off. Apply morning or evening when temperatures are cooler. Re-apply 5-7 days later or as necessary. Use high pressure with fine droplet. It is important that the spray makes direct contact with pests.

As fungicide on Grapevines:

Critical times for botrytis control are during flowering, pre bunch closure, veraison and the period up to harvest when the berries are maturing. If HITMAN is applied only at these times suppression of botrytis will be obtained but the level of control will not be as high as that achieved with a regular program of application at 21 day intervals.

MIXING: Add the product at the appropriate rate to water and mix well. Use good quality, soft water of pH 6 to 8. The use of hard water may cause precipitation and the addition of a softener is recommended.

APPLICATION: HITMAN should only be applied using suitable dilute ground application equipment. Concentrate spraying is not recommended. Equipment should produce a spray that ensures penetration of the canopy and coverage of the flowers or bunches. Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed. The sprayer should be set-up and operated to achieve even coverage throughout the crop canopy to the point of run-off. Avoid excessive run-off.