

Label image proudly supplied by



POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

BAYONET

in-FURROW and FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP C FUNGICIDE

For the control of certain fungal diseases on wheat barley and canola, when mixed with fertiliser or applied as a foliar spray as per the Directions for Use table



CONQUEST AGROCHEMICALS PTY. LTD.
ABN 84 098 814 932
Unit 1, 46 Hasler Road,
Osbourne Park, WA 6017
Telephone: 08 9347 0500, Facsimile: 08 9347 0551.



CONTENTS
20L 200L
1000L

84% of active

CONQUEST

BAYONET

in-FURROW and FOLIAR FUNGICIDE

DIRECTIONS FOR USE: ALL STATES

A. BAYONET APPLIED IN-FURROW AS A FERTILISER TREATMENT

Crop	Disease	Rate per hectare	WHP	Critical Comments																								
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400 mL	4 weeks grazing	<p>Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.</p> <p>Application: Spray undiluted product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 400 mL of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as a guide.</p> <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333												
	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																										
60	667*																											
70	571																											
80	500																											
100	400																											
120	333																											
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	200 or 400mL		<p>For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.</p> <p>Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.</p> <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare kg</th> <th colspan="2">Volume of product per 100kg of fertiliser (mL)</th> </tr> <tr> <td></td> <th>200mL rate</th> <th>400mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td></tr> <tr><td>60</td><td>333</td><td>667*</td></tr> <tr><td>70</td><td>286</td><td>571</td></tr> <tr><td>80</td><td>250*</td><td>500</td></tr> <tr><td>100</td><td>200*</td><td>400</td></tr> <tr><td>120</td><td>167*</td><td>333</td></tr> </tbody> </table> <p>* Application volumes over 600mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. † Where application volume is low, product may be diluted with water to improve coverage.</p>	Fertiliser rate per hectare kg	Volume of product per 100kg of fertiliser (mL)			200mL rate	400mL rate	50	400	800*	60	333	667*	70	286	571	80	250*	500	100	200*	400	120	167*	333
	Fertiliser rate per hectare kg	Volume of product per 100kg of fertiliser (mL)																										
	200mL rate	400mL rate																										
50	400	800*																										
60	333	667*																										
70	286	571																										
80	250*	500																										
100	200*	400																										
120	167*	333																										
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	200 or 400mL																										

B. BAYONET APPLIED AS A FOLIAR SPRAY

Crop	Disease	Rate per hectare	WHP	Critical Comments
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	250mL-500mL	7 weeks Harvest or Grazing	<p>Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3 – 4 weeks apart at the low rate.</p> <p>Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3 – 4 weeks apart at the low rate.</p>
	Leaf rust (<i>P. recondita</i>) Septoria nodorum blotch (<i>Septoria nodorum</i>) Septoria tritici blotch (<i>S. tritici</i>)	<p>Ground Application Apply in 100-150L water plus 200mL Conquest Wetter 1000 per 100L of spray solution.</p> <p>Aerial Application Apply in a minimum of 15L water plus 200mL Conquest Wetter 1000 per 100L of spray solution.</p>		
Barley	Powdery mildew (<i>Erysiphe graminis</i>)	200mL Conquest Wetter 1000 per 100L of spray solution.	10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3 – 4 weeks apart at the low rate, may be required.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

APPLIED IN-FURROW AS A FERTILISER TREATMENT:

CANOLA, WHEAT AND BARLEY
 – DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

APPLIED AS A FOLIAR SPRAY

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY – 10 WEEKS
 WHEAT – 7 WEEKS
 DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:
 BARLEY – 10 WEEKS
 WHEAT – 7 WEEKS

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

GROUP C FUNGICIDE

Bayonet In-Furrow and Foliar Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Bayonet In-Furrow and Foliar Fungicide is a Group C fungicide. Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled this product and other Group C fungicides thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use Conquest Agrochemicals Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

Application

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.
Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertiliser.

Foliar Application

Check the drop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
 DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL (20 and 200L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE AND DISPOSAL (1000L)

DO NOT store or handle near heat or naked flame. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store this tightly closed minibulk container in a safe and secure place away from children, animals, food and feedstuffs. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Conquest Agrochemicals Pty Ltd should be advised immediately.

Do NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Conquest Agrochemicals Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill or judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OF FIRE BRIGADE	SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
--	--



CONQUEST AGROCHEMICALS PTY. LTD.
 ABN 84 098 814 932
 Unit 1, 46 Hasler Road,
 Osborne Park, WA 6017
 Telephone: 08 9347 0500, Facsimile: 08 9347 0551

Batch No:

DOM:

APVMA Approval No: 58688/20/0105

58688/200/0105

58688/1000/0105

