

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

VETO T



DRY SEED DRESSING

ACTIVE CONSTITUENTS: 25 g/kg TEBUCONAZOLE
4 g/kg TRIFLUMURON

GROUP C FUNGICIDE

GROUP 15A INSECTICIDE

For control of bunt, flag smut and loose smut of wheat.
For control of covered and loose smut of barley and oats.
Treated seed is also protected against insect pests of stored grain.

 **CONQUEST**
Agrochemicals

NET CONTENTS

10 kg 15 kg

VETO T DRY SEED DRESSING

Directions for use – QLD, NSW, Vic, TAS, SA & WA Only

Restraint:

Do NOT treat physically damaged seed.

Crop	Disease/Pest	Rate	Critical Comments
Wheat	Bunt (<i>Tilletia foetida</i> , <i>T. caries</i>), Soilborne Flag smut (<i>Urocystis agropyri</i>), Seedborne Flag smut (<i>Urocystis agropyri</i>) Loose smut (<i>Ustilago segetum var tritici</i>),	100 g/ 100 kg seed	Thorough mixing is essential. Ensure that the recommended rate is applied.
Barley	Covered smut (<i>Ustilago segetum var segetum</i>) Loose smut (<i>Ustilago segetum var tritici</i>)		
Oats	Covered smut (<i>Ustilago segetum var segetum</i>) Loose smut (<i>Ustilago segetum var tritici</i>)		
Wheat, Barley, Oats	Protection against insect pests of stored grains: Granary weevil (<i>Sitophilus granarius</i>), Lesser grain borer (<i>Rhyzopertha dominica</i>), Rice weevil (<i>Sitophilus oryzae</i>), Rust-red flour beetle (<i>Tribolium castaneum</i>), Saw toothed grain beetle (<i>Oryzaephilus surinamensis</i>)		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:

**FOOD PRODUCING CROPS - NOT REQUIRED WHEN USED AS DIRECTED
DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD
WITHIN 4 WEEKS OF SOWING**

General Instructions

Conquest Veto T Dry Seed Dressing may delay emergence of wheat. Crops will recover rapidly in most soils without any adverse effect on yield. Sowing Conquest Veto T Dry Seed Dressing treated seed deeper than 50 mm may reduce plant establishment in heavier soils or soils that will crust or clod at the surface after sowing. Sowing rates may need to be adjusted accordingly. Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment probably will be necessary. Treated seed prevents insects from multiplying and preserves grain condition but may not kill adult insects from other infested grain. If the seed is not used immediately after treatment it should be stored in a dry, well ventilated place. Treated seed not used for sowing in the current or following season should be destroyed. **Resistance note:** The effectiveness may be reduced or nullified as a result of pests or diseases developing a tolerance or resistance to the product. If unacceptable control occurs immediately contact your Conquest Agrochemicals Pty Ltd. This product should not be used where the user suspects that a tolerant or resistant strain is present.

FUNGICIDE RESISTANCE WARNING

GROUP C FUNGICIDE

Conquest Veto T Dry Seed Dressing is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide. Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Conquest Agrochemicals Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

INSECTICIDE RESISTANCE WARNING

GROUP 15A INSECTICIDE

For insecticide resistance management Conquest Veto T Dry Seed Dressing is a Group 15A insecticide.

Some naturally occurring insect biotypes resistant to the product and other Group 15A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if this product or other Group 15A insecticides are used repeatedly. The effectiveness of this product on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Conquest Agrochemicals Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant insects.

Conquest Veto T Dry Seed Dressing may be subject to specific resistance management strategies. For further information contact your local supplier, Conquest Agrochemicals Pty Ltd representative or local agricultural department agronomist.

MIXING

The required quantity of Conquest Veto T Dry Seed Dressing should be thoroughly mixed with the correct weight of seed in a seed dressing machine or a suitable revolving drum or barrel. Wear a face mask when handling the dust or treated seed.

SEED QUALITY

Veto T seed treatment should not be used on seed with more than 12 % moisture content, or on sprung, sprouted, damaged or severely pinched seed, or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard.

STORAGE OF TREATED SEED

Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment will be necessary. Treated seed prevents insects from multiplying and preserves grain condition but may not kill adult insects migrating from other infested grain. If the seed is not used immediately after treatment it should be stored in a dry, well-ventilated place. Treated seed not used for sowing in the current or following season should be destroyed. Although Veto T has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for the performance of stored treated seed. Refer to STORAGE AND DISPOSAL section for additional instructions.

PRECAUTION

DO NOT use treated seed for human consumption.

DO NOT allow seed treated with this product to contaminate seed intended for animal or human consumption. If possible, use a separate auger for treating seed. If using the same auger for treating seed and moving grain for human consumption, remove all Veto T residues from the auger to avoid contaminating untreated seed with seed dressing residues. If inhalation of dust is likely, wear a disposable dust mask while handling the dust and treated seed.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption or poultry feed or mixed with animal feed.

DO NOT allow seed treated with this product to contaminate seed intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds.

Harmful to fish and aquatic arthropods. DO NOT contaminate dams, streams, rivers or waterways with the chemical, or containers or bags which have held the chemical or treated seed.

STORAGE AND DISPOSAL

Treated seed when stored should be kept apart from other seed and the containers should be clearly marked to indicate the contents have been treated with this product. Store product in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in sunlight. Seed treated with this product must not be used for human consumption or be allowed to contaminate seed intended for human consumption.

Containers which have held treated seed should not be used for any other purpose. Triple or preferably pressure rinse empty containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill or judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

**For specialist advice in an emergency dial
1 800 033 111
24 hours Australia wide.**

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932

46 Hasler Road

Osborne Park WA 6017

Telephone: (08) 9347 0500

Facsimile: (08) 9347 0551

Batch No:

D.O.M.

APVMA Approval No: 59322110kg0306

59322115kg0306

CONQUEST
Agrochemicals