

# POISON

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Label image proudly supplied by



# TRICAM

FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

**GROUP C FUNGICIDE**

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as specified in the 'Directions for Use' Table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**



SIPCAM PACIFIC AUSTRALIA PTY LIMITED Suite 11, 23-31 Gheringhap Street, Geelong Vic 3220  
TEL: (03) 5223 3746 ACN 073 176 888

**CONTENTS: 1000L  
APVMA APPROVAL NO:  
59349/1000L/1104**



# TRICAM

FUNGICIDE

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferable pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable Containers empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the MSDS, which is available from the supplier.

## NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following. To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

Date of Manufacture:

Batch No:

IN TRANSPORT  
EMERGENCY DIAL **000**  
Police or Fire Brigade

FOR SPECIALIST ADVICE IN AN  
EMERGENCY DIAL **1800 033 111**  
24 hours Australia wide

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

A. Tricam Fungicide Applied In-Furrow as a fertiliser  
TREATMENT:  
CANOLA, WHEAT and Barley: DO NOT GRAZE OR CUT FOR STOCK FOR 4 WEEKS AFTER SOWING.

B. Tricam Fungicide Applied as a Foliar Spray  
DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION.  
BARLEY - 10 WEEKS  
WHEAT - 7 WEEKS  
DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION.  
BARLEY - 10 WEEKS  
WHEAT - 7 WEEKS

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.  
Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the corrected rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

Foliar Application:

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

Fungicide Resistance Warning

**GROUP C FUNGICIDE**

Tricam Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Tricam Fungicide is a Group C fungicide. Some naturally occurring individual fungi resistant to Tricam Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Tricam Fungicide and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Tricam Fungicide to control resistant fungi. Resistant should not be assumed without first reviewing the method of application, the coverage and the time of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferable pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt. For Refillable Containers - empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the MSDS, which is available from the supplier

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following. To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

IN TRANSPORT EMERGENCY DIAL 000 Police or Fire Brigade	FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 24 hours Australia wide
--	---

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**TRICAM**  
FUNGICIDE

ACTIVE CONSTITUENT: ZENOL FLUTRIACOL

**GROUP C FUNGICIDE**

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as specified in the 'Directions for Use' Table.

APVMA APPROVAL NO: 59349/1104

THIS LABEL IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

SIPCAM PACIFIC AUSTRALIA PTY LIMITED  
Suite 11, 28-31 Sherdinghap Street,  
Geelong Vic 3220  
AGN 073 176 888  
TEL: (03) 5223 3746

DIRECTIONS FOR USE

A. Tricam Fungicide as a fertiliser treatment

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments												
Canola	Blackleg ( <i>Leptosphaeria macularis</i> )	400 mL	4 Weeks grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. <b>Application:</b> Spray undiluted product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400 mL of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as guide.												
				<table border="1"> <thead> <tr> <th>Fertiliser Rate Per hectare</th> <th>Volume of Product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser Rate Per hectare	Volume of Product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333
Fertiliser Rate Per hectare	Volume of Product per 100 kg of fertiliser (mL)															
60	667*															
70	571															
80	500															
100	400															
120	333															
Wheat	Stripe Rust ( <i>Puccinia striiformis</i> )	200 or 400 mL	4 Weeks grazing	For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. <b>Application:</b> Spray product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific ratio of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as guide.												
	Septoria Tritici blotch ( <i>S. tritici</i> ), Take - all ( <i>Gaeumannomyces graminis</i> )	400 mL		<table border="1"> <thead> <tr> <th>Fertiliser Rate Per hectare</th> <th>Volume of Product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser Rate Per hectare	Volume of Product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333
Fertiliser Rate Per hectare	Volume of Product per 100 kg of fertiliser (mL)															
60	667*															
70	571															
80	500															
100	400															
120	333															
Barley	Powdery mildew ( <i>Erysiphe graminis</i> ) Scald ( <i>Rhynchosporium secalis</i> )	200 or 400 mL	4 Weeks grazing	<table border="1"> <thead> <tr> <th>Fertiliser Rate Per hectare</th> <th>Volume of Product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser Rate Per hectare	Volume of Product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333
				Fertiliser Rate Per hectare	Volume of Product per 100 kg of fertiliser (mL)											
60	667*															
70	571															
80	500															
100	400															
120	333															
				* Application volumes over 600 mL per 100 kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilizers. # Where application volume is low, product may be diluted with water to improve coverage.												

B. Tricam Fungicide Applied as a Foliar Spray

Crop	Disease	Rate Per Hectare	WHP (days)	Critical Comments
Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	250 - 500 mL <b>Ground Application:</b> Apply in 100 - 150 mL water plus 200 mL "BS 1000" per 100 L of spray solution. <b>Aerial Application:</b> Apply in a minimum of 15L water plus 200 mL "BS 1000" per 100 L of spray solution.	7 Weeks harvest or grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.
	Leaf rust ( <i>Puccinia striiformis</i> ) Septoria nodorum blotch ( <i>Septoria nodorum</i> ) Septoria Tritici blotch ( <i>S. tritici</i> )	Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe diseases persist during flowering, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.		
Barley	Powdery mildew ( <i>Erysiphe graminis</i> )		10 weeks harvest or grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.



100%