


MAIN PANEL

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**Antracol® 700 WG**  
**FUNGICIDE**



ACTIVE CONSTITUENT: 700 g/kg PROPINEB

**GROUP M3 FUNGICIDE**

For the control of certain diseases of cucurbits, onions, potatoes, and tomatoes as well as black spot and citrus rust mite of citrus as specified in the DIRECTIONS FOR USE table

**\* kg net**

(Label code)

\* 1, 10 kg

## REAR PANEL

## DIRECTIONS FOR USE

CROP	DISEASE/PEST	STATE	RATE	WHP (Days)	CRITICAL COMMENTS
Citrus	Black spot, citrus rust mite	NSW, Vic, SA, WA only	200 g/100 L plus white oil 250 mL/100 L	7	Apply first application 6 weeks after copper spray at petal fall. Coastal lowlands and highlands (NSW): apply first application 8 weeks after the petal fall copper spray. Apply no more than 3 applications per season. <b>Dilute spraying only.</b> Refer to the <b>Special instructions for citrus</b> section in <b>GENERAL INSTRUCTIONS</b> .
		Qld only	200 g/100 L plus white oil 600 mL/100 L		
Cucurbits e.g. cucumbers, rockmelons	Downy mildew	All states	200 g/100 L or 2 kg/ha	3	Apply at weekly intervals while conditions favour disease. Apply no more than 4 applications per crop. Spray both sides of leaves thoroughly.
Onions	Downy mildew	All states	200 g/100 L or 2 kg/ha	7	Apply at intervals of 7-10 days from early in the life of the crop. Apply no more than 4 applications per crop. If conditions favour disease development consider using a systemic fungicide. Add wetting agent.
Potatoes	Target spot (early blight), Irish blight (late blight)	All states	200 g/100 L or 2 kg/ha	1	Apply when conditions favour disease and repeat at 7-10 day intervals. Apply no more than 4 applications per crop.
Tomatoes	Target spot (early blight), Irish blight (late blight), grey leaf spot ( <i>Stemphylium</i> )	All states	200 g/100 L or 2 kg/ha	3	Apply when conditions favour disease and repeat at 7-10 day intervals. Apply no more than 4 applications per crop. In places where diseases are severe commence at seedbed stage.
Watermelon, Rockmelon	Anthracnose, gummy stem blight	Qld, NSW, SA, WA only	200 g/100 L or 2 kg/ha	3	Apply every 7-10 days depending on weather and disease incidence. First application may be made when vines begin to run. Apply no more than 4 applications per crop.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS (WHP)**

**Potatoes: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION**

**Cucurbits, tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION**

**Citrus, onions: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

## REAR PANEL (continued)

**GENERAL INSTRUCTIONS****Fungicide Resistance Warning**

<b>GROUP</b>	<b>M3</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

Antracol is a member of the multi-site group of fungicides. For fungicide resistance management Antracol is a Group M3 fungicide. Some naturally occurring individual fungi resistant to Antracol and other Group M3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group M3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

**Export of treated produce**

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Antracol. If you are growing edible produce for export, please check with Bayer CropScience Pty Ltd for the latest information on MRLs and import tolerances before using Antracol.

**Mixing**

Add the required amount of Antracol to the spray tank containing half the required volume of water. Mix thoroughly and make up to the required volume with water.

**Application**

Good disease control requires even, thorough coverage of the target area. Application should be made using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Equipment settings and water volume may need to vary, depending on the growth stage of the crop. Constant agitation of the spray solution is required prior to and during spraying.

**Special instructions for citrus*****Dilute spraying***

- ◆ Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off.
- ◆ The required spray volume to achieve point of run off may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice.
- ◆ Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- ◆ The required dilute spray volume to achieve point of run off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

**REAR PANEL (continued)****Compatibility**

Ensure that water is of suitable quality for mixing with agricultural pesticides. Poor quality water can cause physical incompatibility between some products which may result in crop damage in some circumstances. Antracol is physically compatible with most commonly used insecticides and fungicides and has been successfully tested in a two way mixture with a range of products including Agrimec<sup>®</sup>, Amistar<sup>®</sup> WG, Avatar<sup>®</sup>, Bayfidan<sup>®</sup> 250 EC, Bulldock<sup>®</sup> 25 EC, Chess<sup>®</sup> 500 WG, Confidor<sup>®</sup> 200 SC, Larvin<sup>®</sup>, Proclaim<sup>®</sup>, Rovral<sup>®</sup> Aquaflor, Success<sup>®</sup> and Supracide<sup>®</sup> 400 EC. The biological compatibility of these mixtures has not been specifically tested.

Antracol is not compatible with Liquicop<sup>®</sup> (copper as ammonium acetate), lime sulphur or any alkaline materials. When mixing more than one product in the spray tank, products should be added in the following order, according to formulation type:

1. Wettable powders [WP] (pre-mixed with water)
2. Suspension concentrates [SC]
3. Water dispersible granules [WG/WDG] (e.g. Antracol 700 WG Fungicide)
4. Water soluble powders or granules [SP or SG]
5. Emulsifiable concentrates or oil in water emulsions [EC or EW]
6. Soluble concentrates [SL]
7. Wetting agents or oils

Do not mix concentrates together but add each to the spray tank separately. Constant agitation of the spray solution is required for any mixtures containing Antracol, otherwise settling will occur.

As formulations of other manufacturers' products are beyond the control of Bayer CropScience, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Bayer CropScience cannot be responsible for the behaviour of such mixtures.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, dry, well-ventilated area out of direct sunlight.

Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty bags in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt. Do not re-use empty container for any other purpose.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26). Avoid giving alcohol.

**SAFETY DIRECTIONS**

Harmful if inhaled. Do not inhale dust. Avoid contact with skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves and a disposable dust mask covering mouth and nose. When using the prepared spray [apply by hand or air-assisted spray equipment], wear cotton overalls, over normal clothing, buttoned to the neck and wrist and elbow-length PVC gloves. After each day's use wash gloves and contaminated clothing.

**Re-entry** Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Do not perform high exposure activities such as thinning or hand harvesting in citrus for 13 days, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**REAR PANEL (continued)****SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.bayercropscience.com.au](http://www.bayercropscience.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 59925/37055

Antracol®, Bayfidan®, Bulldock®, Confidor® and Larvin® are Registered Trademarks of the Bayer Group.

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

BAR  
CODE

Bayer CropScience Pty Ltd  
ABN 87 000 226 022  
391-393 Tooronga Rd  
Hawthorn East, Vic. 3123  
Phone: (03) 9248 6888  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)  
Technical Enquiries: 1800 804 479



Bayer CropScience

(Label code)

Batch Number:

Date of Manufacture: