

Product Name: GRAZON EXTRA HERBICIDE  
APVMA Approval No: 60830/147022



Label Name:	GRAZON EXTRA HERBICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	300 g/L TRICLOPYR PRESENT AS THE BUTOXYETHYL ESTER 100 g/L PICLORAM PRESENT AS THE HEXYLOXYPROPYLAMINE SALT 8 g/L AMINOPYRALID PRESENT AS HEXYLOXYPROPYLAMINE SALT
Mode of Action:	GROUP 4 HERBICIDE
Statement of Claims:	For the control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use.
Net Contents:	1000L 100L 10L 1L 20L 500mL 5L
Restrains:	<ul style="list-style-type: none"><li>• DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.</li><li>• DO NOT spray if rain is likely within 1 hour or if foliage is wet from rain or dew.</li><li>• DO NOT burn off, cut or clear blackberry or other woody weeds for at least 6 months after spraying.</li><li>• DO NOT apply by aerial application in wind in excess of 15 km/hour and/or air temperatures above 35°C.</li><li>• In areas prone to flooding treatment should commence after any annual flooding, as such areas flooded within 9 months following application may have reduced results.</li></ul>

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	<p>IN TASMANIA FOR BLACKBERRY</p> <ul style="list-style-type: none"> <li>• DO NOT treat bushes carrying mature or near mature fruit.</li> </ul> <p>FOR NATIVE VEGETATION</p> <ul style="list-style-type: none"> <li>• Use of Grazon Extra on native vegetation must be done in accordance with STATE and/or LOCAL legislation.</li> </ul>
--------------------	---

Withholding Periods:	<p>Pasture: Cutting or Grazing Pastures for Stock Food: NOT REQUIRED WHEN USED AS DIRECTED.</p>
----------------------	---

Trade Advice:	<p><b>LIVESTOCK DESTINED FOR EXPORT MARKETS</b> The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.</p> <p><b>EXPORT SLAUGHTER INTERVAL (ESI) – 3 days:</b> Livestock that has been grazed on or fed treated crops within 42 days of application should be placed on clean feed for 3 days prior to slaughter.</p> <p><b>EXPORT GRAZING INTERVAL (EGI) – 42 days:</b> Livestock that has been grazing on treated crops or pasture should not be sold for export slaughter for 42 days (6 weeks) after application of the chemical product, unless the export slaughter interval has been observed.</p> <p>When Grazon Extra is used as directed and the above withholding periods and/or export intervals are observed, treated grain and livestock commodities are considered acceptable for export. However, export requirements are subject to change. Consult your exporter for updated information about specific market requirements.</p> <p><b>IMPORTANT: Read the MANAGEMENT OF RESIDUES IN COMPOST, MULCHES AND ANIMAL WASTE in the PROTECTION OF CROPS, NATIVES AND OTHER NON-TARGET PLANTS section of this label</b></p>
---------------	---

General Instructions:

**MIXING**

- Mix only with water.
- Half fill the spray unit with water and add the required amount of Grazon Extra. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

**APPLICATION**

ONLY apply Grazon Extra under atmospheric conditions that do not allow drift onto sensitive crops to occur.

**1. WOODY WEED SITUATIONS**

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed, or areas having a previous chemical treatment.

**A. High Volume Spraying**

- Thorough coverage of foliage to the point of run-off is essential; however, avoid excess spraying which is wasteful of chemical.

**Hand-Gun**

- Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1,500 kPa (400 to 500 kPa for St John's wort).
- A spray volume of 3,000 to 4,000 L per infested hectare of 1 to 2 metre high blackberry (30 to 40 L/100 m<sup>2</sup>) should be used.
- Use 2,000 L of spray mixture/ha of galenia infestation (i.e. 20 L/100 m<sup>2</sup> infested area).

**Knapsack & 12 volt Sprayer Packs**

- Only recommended for the control of herbaceous weeds, such as capeweed, fireweed and spear thistle, and woody weeds that are not regrowth, less than 60 cm high or 60 cm diameter.
- Apply the recommended spray mix to give full coverage of leaves and stems.

**B. Aerial Application**

- Apply in 200 L of water/ha using an aircraft to apply 100 L per pass on a double overlap pattern using nozzle configurations to produce coarse to very coarse droplets as defined by ASAE S572.

**C. Boom Application**

- Application in a minimum spray volume of 200 L/ha for galenia and St. John's wort and 600 L of water/ha for sickle pod. Use nozzle configurations to produce coarse spray droplets as defined by ASAE S572 definition for standard nozzles. Boom height must be set to ensure double overlap of nozzle patterns.

**2. FALLOW SITUATIONS**

**A. Boom Application**

- Application of Grazon Extra in a minimum spray volume of 70 L/ha is recommended using nozzle configurations to produce coarse spray droplets as defined by ASAE S572 definition for standard nozzles. Boom height must be set to ensure double overlap of nozzle patterns.

## B. Blanket Wiper Application

- Blanket needs to be made from durable and wettable material with a rigid backing.
- Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting).
- Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle.
- Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10–15 kph are ideal for blanket wiping application.

## COMPATIBILITY

- Follow any regional restrictions, and all directions and restrictions on the labels of any chemical mixed with Grazon Extra (e.g. 2,4,-D amine).
- Grazon Extra is compatible with the following herbicides:

Glyphosate 450, metsulfuron-methyl, Roundup CT, Starane Advanced, Statesman 720 (2,4-D DMA 720 g/L).

Grazon Extra is compatible with the following adjuvants, as per Directions for Use on this label:

Uptake Spraying Oil, BS1000 Biodegradable Surfactant, (alternatives are Chemwet 1000 and Spreadwet 1000 Wetting Agent) or Pulse Penetrant.

\* Not all surfactants or crop oils are of equal quality, Corteva Agriscience does not support the use of alternative products other than those listed in the COMPATIBILITY section.

## CLEANING SPRAY EQUIPMENT

### Rinsing

\* After using Grazon Extra, empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.

\* After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

### Decontamination

Before spraying cotton and other sensitive crops with equipment that has been used to apply Grazon Extra (see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section).

Wash the tank and rinse the system as above. Then quarter fill the tank and add a standard alkali-based laundry detergent at 500 g (or mL)/100 L water and circulate throughout the system for at least 15 minutes. If using a concentrated laundry detergent, use 250 g (or mL)/100 L water. Do not use chlorine-based cleaners.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

Resistance Warning:	<p><b>RESISTANCE WARNING</b> <b>GROUP 4 HERBICIDE</b></p> <p>Grazon Extra Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group 4 herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Corteva Agriscience representative.</p>
---------------------	--

Precautions:	
--------------	--

Protections:	<p><b>PROTECTION OF LIVESTOCK</b></p> <ul style="list-style-type: none"> <li>• Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.</li> <li>• Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.</li> </ul> <p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b></p> <ul style="list-style-type: none"> <li>• DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</li> </ul> <p><b>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</b></p> <ul style="list-style-type: none"> <li>• Crops susceptible to Grazon Extra include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and Pinus spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines.</li> <li>• Grazon Extra is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.</li> <li>• Picloram and aminopyralid, two of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.</li> <li>• DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.</li> <li>• DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops.</li> <li>• DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.</li> <li>• DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants.</li> <li>• DO NOT move soil, which may have been treated to areas where desirable plants are to be grown.</li> </ul>
--------------	---

Storage and Disposal:	<ul style="list-style-type: none"> <li>• Store in the closed, original container in a cool, well ventilated area.</li> <li>• DO NOT store for prolonged periods in direct sunlight.</li> <li>• DO NOT store near food, feedstuffs, fertilisers or seed.</li> </ul>
-----------------------	--

The method of disposal of the container depends on the container type. Read the STORAGE AND DISPOSAL instructions on the label that is attached to the container.

500 mL and 1 L containers

Do not re-use containers.

Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

5, 10 and 20 L containers

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple rinse container for disposal. Dispose of rinsate by adding to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

100 L container

Empty contents fully into application equipment. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained. Add the rinsings to the spray tank. When the drum is empty, close all valves and return to the point of supply for refill or storage. Do not remove or tamper with the dry valves or security seal. Do not contaminate the drum with water or any other foreign matter.

IBC (intermediate bulk container) - 1,000 L

Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Close all valves and arrange for collection under the relevant return programme.

Safety Directions:

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrists (or equivalent clothing), elbow-length PVC gloves and face shield or goggles. For hand-held application: When using the prepared spray, wear cotton overalls buttoned to the neck and wrists (or equivalent clothing), elbow-length PVC gloves and a face shield or goggles.

After each day's use, wash gloves, face shield or goggles and contaminated clothing. Wash hands after use.

First Aid Instructions:

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15min and see a doctor.

First Aid Warnings:

## DIRECTIONS FOR USE

### 1. WOODY WEED SITUATIONS

**Table A: High Volume Spraying**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

Note: Corteva Agriscience only recommends the use of knapsacks or 12 volt sprayers for use on low growing herbaceous weeds and woody weeds that are not regrowth and are less than 60 cm high or 60 cm diameter.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /100 L Water	CRITICAL COMMENTS
African boxthorn	Less than 2 m tall	500 mL	Apply when bushes have good leaf cover, growth and <b>no leaf fall</b> .
<i>Angophora</i> spp.	1 to 3 m tall	350 mL	
Australian blackthorn	Less than 2 m tall	500 mL	Apply from late spring to early autumn.
<i>Banksia</i> spp.	1 to 3 m tall	350 mL	
Biddy bush (Chinese shrub) (Sifton bush)	Autumn when actively growing	500 mL	Add BS1000 or an alternative (see COMPATIBILITY section) at 125 mL/100 L of water for best results.
Blackberry in association with: Docks Ragwort Smartweed Thistles	Late spring to autumn	350 mL or 500 mL	Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry. Where herbicides other than Group 4 herbicides have been used, allow 2 seasons regrowth to occur before respraying with Grazon Extra.
Blue heliotrope	Flowering	500 mL	Apply in a minimum spray volume of 1250 L/ha.
Brooms: Cape English Flax leaf Montpellier	Spring to mid-summer prior to pod formation	250 mL	Apply as a thorough foliage spray.
	Autumn to winter	350 mL	
Camphor Laurel	Less than 2 m tall		
	Above 2 m tall	500 mL	
Capeweed	Flowering	150 mL	
<i>Casuarina</i> spp.	1 to 3 m tall	350 mL	

**Table A: High Volume Spraying - *continued***

**See GENERAL INSTRUCTIONS - APPLICATION section for application method details.**

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>RATE /100 L Water</b>	<b>CRITICAL COMMENTS</b>
Chinee apple	Less than 2 m tall	350 mL	Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Cockspur thorn	Spring to autumn		Apply as a thorough foliage spray.
Common sensitive plant	Any time when actively growing	200 mL	To avoid leaves closing during application, spray plants while moving forward. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Crofton weed	Spring to autumn	350 mL	Apply as a thorough foliage spray.
Eastern cotton bush ( <i>Maireana microphylla</i> )		500 mL	Add Uptake Spraying Oil at 500 mL/100 L of water. Some bushes may require a follow-up spray to control regrowth.
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3 m tall	350 mL or 500 mL	<b>Apply the high rate where difficult to control species of <i>Eucalyptus</i> regrowth is present.</b> Addition of an adjuvant may improve results. Contact Corteva Agriscience for details.
Fireweed	Flowering	350 mL	Apply as a thorough foliage spray.
Galenia	Fresh growth in spring to summer	500 mL	Use 2000 L of spray mixture/ha.
Giant bramble	Spring to autumn		Penetration of thick clumps may be difficult and respraying may be necessary. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Gorse	1 to 1.5 m tall	250 mL	Spring and summer treatment only. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
	Over 1.5 m tall or autumn treatment	350 mL	Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
	Winter treatment	500 mL	Brownout may not be complete until summer. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Groundsel bush ( <i>Baccharis halimifolia</i> )	1 to 1.5 m tall in spring to summer	250 mL	Apply as a thorough foliar spray.
	Over 1.5 m tall or autumn treatment	350 mL	



**Table A: High Volume Spraying - *continued***

**See GENERAL INSTRUCTIONS - APPLICATION section for application method details.**

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>RATE /100 L Water</b>	<b>CRITICAL COMMENTS</b>
Green cestrum	Late spring to early autumn	500 mL	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
Hawthorn	Less than 2 m tall		Apply from late spring to early autumn.
Horehound	Pre-flowering	350 mL	Apply as a thorough foliar spray.
Japanese sunflower			
Lantana ( <i>Lantana camara</i> ) ( <i>Lantana montevidensis</i> )	Up to 1 m tall in summer to autumn		Add one of the following adjuvants, when using 350 mL rate: Uptake Spraying Oil @ 0.5% v/v. Pulse Penetrant @ 0.1% v/v.
	1 to 2 m tall in summer to autumn	500 or 750 mL	Thoroughly wet foliage stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
Lion's tail ( <i>Leonatis nepetifolia</i> )	Pre-flowering	200 mL	Apply as a thorough foliar spray. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Limebush	Any time of year with good leaf cover and soil moisture	350 mL	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results. Contact Corteva Agriscience for details.
Manuka	At flowering	500 mL	For optimum results, add Pulse Penetrant at 200 mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite ( <i>Prosopis</i> spp.)	Seedling, full leaf and flowering before podding	350 mL	<b>DO NOT</b> spray plants bearing pods. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Prosopis velutina		670 mL	
Milfoil (Yarrow)	Flowering	350-500 mL	Use low rate when in close proximity to highly sensitive vegetation.
Mistflower	Spring to Autumn	350 mL	Apply as a thorough foliar spray.
Mother-of-millions	Flowering	500 mL	Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water.
Paddy's lucerne	Active growth		Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2 m tall	350 mL	Add Uptake Spraying Oil at 500 mL/100 L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Paterson's curse	Flowering	250 mL	
Prickly pear (common) Smooth tree pear	Active phyllode growth	500 mL	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.

**Table A: High Volume Spraying - *continued***

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>RATE /100 L Water</b>	<b>CRITICAL COMMENTS</b>
Rubber vine (Not infected with rust)	Up to 1.5 m tall at flowering	350 mL	Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands, regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5 m tall at flowering	500 mL	
Siam weed	Active growth	350 mL	Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Sicklepod	Up to flowering	200 mL	<b>DO NOT</b> apply to podding plants. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Spear thistle	Rosette to flowering	150 mL	
St John's wort	From flowering to early seed set	500 mL	Late spring to early summer.
Sweet briar	Up to 1.5 m tall	350 mL	Add metsulfuron-methyl at 10 g/100 L water to obtain more reliable results with the lower rate of Grazon Extra.
		500 mL	Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Tobacco weed	Actively growing plants	300 mL	Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water for best results.
Tropical soda apple	Flowering up to 1 m tall	350 mL	Add Uptake Spraying Oil at 500 mL/100 L water.
Wattle ( <i>Acacia</i> spp.) (except corkwood wattle)	1 to 3 m tall		
Wild Rosemary ( <i>Cassinia laevis</i> )	Active growth 0.5 to 1 m tall	350 to 500 mL	Use lower rate on seedlings 0.5 m tall. Apply as a thorough foliar spray.
Wild tobacco tree	Spring to autumn up to 2 m tall	350 mL	

**Table B: Aerial Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry	Summer to autumn	ACT, NSW, NT, QLD, SA, Vic, WA only	10 L	Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with Gazon Extra. <b>WARNING:</b> <i>Eucalyptus</i> species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 to 20 m tall may be partially defoliated but are likely to recover.
Gorse		Tas only		Helicopter application only.
Cockspur thorn Crofton weed Lantana Mistflower	Late autumn	ACT, NSW, NT, QLD only (helicopter only)	1.5 L + 5.2 L Statesman 720	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha. <b>Follow-up respraying will be required.</b>
Lantana			10 L	Helicopter application only.
Rubber vine (Not infected with rust)	When flowering	NT, QLD only (helicopter only)	3 L to 5 L	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands, however, complete coverage and penetration may be difficult. <b>Follow-up respraying will be required.</b> Any regrowth should be sprayed with a suitable basal bark herbicide.
St John's wort	Flowering to early seed set (Nov-Jan)	ACT, NSW only	4 L	Helicopter application only. Follow-up spraying will be required in the following season.
AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Parkinsonia	Seedlings 1-2 m tall, or 12 - 24 months old	QLD and NT only (helicopter only)	3 L	Add Uptake Spraying Oil at 1 L/ha.

**Table C: Boom Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
Galenia	Fresh growth during spring to summer	3 to 5 L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage. Use the low rate for seedling weeds not >-50 cm across with excellent fresh active growth in spring right after significant rain of 25 mm or more. When using the lower rate add Pulse Penetrant at 100 mL/100 L for best control.
Sicklepod	Up to flowering	3 L	<b>DO NOT</b> apply to podding plants. Add BS1000 or an alternative (see COMPATIBILITY section) at 100 mL/100 L of water.
St John's wort	Flowering to early seed set (Nov-Jan)	2 to 4 L	Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

**Table D: Boom Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
Flax-leaf fleabane ( <i>Conyza bonariensis</i> )	Up to 8 leaf or up to 10 cm diameter	2 L + 2.4 L glyphosate (450 g/L)	Add 100% non-ionic surfactant (e.g. BS1000) at 0.2 % v/v

## 2. FALLOW SITUATIONS

**Table A: Boom Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry nightshade - <b>suppression only</b>	10 to 25 cm tall, prior to flowering	ACT, NSW, NT, QLD only	200 to 400 mL + 1.2 L glyphosate (450 g/L + adjuvant)	<b>For use by ground equipment only.</b> Plants must be actively growing.

Camel melon Prickly paddy melon Cucumber melon ( <i>Cucumis melo</i> )	From 2 leaf to 50cm diameter			Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for <i>Polymeria pusilla</i> ).
Common sowthistle	From 8 leaf to flowering			Refer to glyphosate (450 g/L) label for use of adjuvant.
Cow vine	From 2 to 5 leaf up to 15cm diameter, prior to flowering			For Northern New South Wales and Queensland, <b>DO NOT</b> plant susceptible crops for up to 9 months after application, as specified in GENERAL INSTRUCTIONS - MINIMUM RECROPPING PERIODS – Table A.
Lucerne (established)	Active growth, 15 to 25 cm high, during spring	300 to 500 mL + 1.2 L glyphosate (450 g/L) + adjuvant		Dry conditions after application will increase the re-cropping interval.
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20cm diameter, prior to flowering	200 to 400 mL + 1.2 L glyphosate (450 g/L) + adjuvant		For Southern New South Wales, please refer to MINIMUM RECROPPING PERIODS – Table C.

**Table B: Blanket Wiper Application**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Bitterbark ( <i>Alstonia constricta</i> )	From summer to end of autumn	NT, Qld only	1:4 (1 part Grazon Extra to 4 parts water)	<p>For use with blanket wipers only.</p> <p>For best results apply in autumn to tall (&gt; 60 cm) plants using two opposite directional passes (up and back).</p> <p>Follow up “missed” plants with a spot spray application. These will be obvious after 6 weeks.</p> <p>Blanket wiper applications can be made in summer when plants are smaller but follow up spot spraying may be necessary.</p> <p>Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).</p>

			2 % solution for spot spray (eg. 100 mL Grazon Extra in 5 L water)	<b>Spot spraying “missed” plants:</b> thoroughly wet all stems and leaves without producing any solution run- off. Avoid any spray reaching the soil surface.
--	--	--	---	---

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### IN TASMANIA FOR BLACKBERRY

DO NOT treat bushes carrying mature or near mature fruit.

### FOR NATIVE VEGETATION

Use of Grazon Extra on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

### MINIMUM RECROPPING PERIODS

Aminopyralid and picloram remain active in the soil for extended periods depending on the rate of application, soil type (clay content), rainfall, temperature, humidity, soil moisture and soil organic matter. The following tables show plant-back periods to particular crops following application of Grazon Extra in different areas/situations of Australia.

### GENERAL INSTRUCTIONS

#### Table A: Northern New South Wales & Queensland

Plant-back periods for rotational crops following application of Grazon Extra for rates up to 600 mL/ha on black cracking clay soils. These plant-back periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plant-back period may be significantly longer.

Plant-back periods for crops following the application of Grazon Extra for rates up to 600 mL/ha.				
RATE mL/ha	200	300	400	600
CROP	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

#### Table B: Blanket Wiper Application

Plant-back periods for crops following blanket wiper application	
CROP	Months
Broadleaf crops	18
Lucerne	6
This will allow any potential soil residues to dissipate, if any, and allow effective control.	

#### Table C: Southern New South Wales

Plant-back periods for rotational crops following application of Grazon Extra for rates up to 500 mL/ha.	
Crops	Plant-back Period (months)
Barley, Canola, Wheat	9
Chickpea, Faba bean, Field pea, Lucerne, Lupin, Medic, Subclover	24

**Note:** Before using Grazon Extra in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, i.e. the product with the longest plant-back period, will determine the time between spraying and planting.

### **MANAGEMENT OF RESIDUES IN COMPOST, MULCHES AND ANIMAL WASTE.**

DO NOT cut pasture for hay or silage production within 6 months of application, where it is intended for use off-farm.

DO NOT use treated plant material to make, compost, mulches or mushroom substrate.

DO NOT send straw from treated pastures off-farm for these purposes or for animal bedding.

DO NOT send animal manure, dairy shed and feed pad effluent that has been collected from animals grazing or fed pastures treated within the previous 6 months off-farm. Spreading/irrigating this manure/effluent may cause damage to clover and other susceptible plants.

DO NOT send compost made from animal waste that has been collected from animals grazing or fed pastures treated within the previous 6 months off-farm. Such compost may cause damage to clover and other susceptible plants.

DO NOT apply animal waste (e.g. manure, slurry) collected from animals grazing or fed pastures treated within the previous 6 months to susceptible plants or land to be used to grow susceptible plants.

DO NOT grow susceptible plants within the relevant plant back period in fields treated with manure/effluent from farms where animals have grazed or been fed treated pasture until a field bioassay shows there are no residues in the soil at levels injurious to the susceptible plants (see the MINIMUM RECROPPING PERIODS section).

To promote herbicide decomposition, manure should be evenly incorporated in the surface soil. Breakdown of residues in decomposing plants or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.

#### **SOIL BIOASSAY**

A simple bioassay can be conducted by collecting at least 10 spade spits of soil to a depth of 200 mm from around the paddock and thoroughly mixing the soil together. Place some of this soil in a shallow container to a depth of 3-5 cm and sow 100 seeds of the susceptible plant to be grown (subterranean or white clover is a good indicator plant where it is not practical to use the susceptible plant) into the soil. Keep in a warm and well-lit location and ensure the soil does not dry out. After plant emergence, check the number of plants that have germinated and seedling vigour. Symptoms of Grazon Extra residues include non-germination or low plant emergence, leaf cupping, leaf whitening, stem elongation and twisting. If these symptoms occur do not grow the susceptible plant. Repeat the bioassay again after a further time interval.