

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**ACCENSI
IMIDACLOPRID
Ready to Use Insecticide**

ACTIVE CONSTITUENT: 0.125g/L IMIDACLOPRID

For the systemic control of various insect pests on flowers, shrubs, trees,
fruit trees and vegetables in the home garden.

Contents: 750mL, 1L

HOW TO PREPARE

The product is ready to use. Shake well before use.

CAUTION

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.
DO NOT spray if bees are feeding on flowering plants. Will kill bees.

STORAGE AND DISPOSAL

Keep out of the reach of children. Store in the closed, original container in a cool, dry place. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet which is available from the supplier upon request.

CONDITIONS OF SALE

The use of Accensi Imidacloprid RTU Insecticide being beyond the control of the manufacturer no warranty expressed or implied is given by ACCENSI PTY LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and ACCENSI PTY LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Batch No:

D.O.M.

Accensi Pty Ltd
ABN: 94 079 875 184
60-76 Potassium Street
NARANGBA QLD 4504
Ph: (07) 3897 2000

APVMA Approval no: 60898/ 59888

Bar Code

HOW TO USE

DO NOT spray if it is raining or about to rain.

DO NOT spray when soil is dry and plants are suffering from moisture stress.

Plant	Pest	How to apply
Roses	Aphids	Spray foliage thoroughly when insects are first seen
Shrubs, plants and ornamental trees, including non-bearing and ornamental citrus	Aphids, Azalea lace bug, Greenhouse thrips, Hibiscus flower beetle, Mealybugs, Psyllids, Sap-sucking bugs (including bronze orange bug), Soft scales	or if insects re-appear. Spray underneath the leaves to kill pests which may be hiding out of sight. For hibiscus flower beetle, spray buds and flowers as needed. For longtailed mealybug, apply 3 sprays 2 weeks apart. For psyllids, spray at first sign and then a week later. For soft scales, spray in late spring or when small scales are first seen. Apply 3 sprays, 2 weeks apart. For best results, spray in the morning when there is little or no wind and little direct heat from the sun.
Peach, plum, nectarine, apricot and other stone fruit, capsicum, zucchinis, melons and other cucurbits, egg plant, potatoes, tomatoes	Green peach aphid	Avoid spraying blooms or flowers in full sun as slight petal marking may occur.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.

DO NOT PICK STONE FRUIT FOR 21 DAYS AFTER SPRAYING.

DO NOT PICK CAPSICUM, POTATOES OR EGG PLANTS FOR 7 DAYS AFTER SPRAYING.

DO NOT PICK TOMATOES FOR 3 DAYS AFTER SPRAYING.

DO NOT PICK CUCURBITS FOR 1 DAY AFTER SPRAYING.