

Product Name: BOXER GOLD HERBICIDE  
APVMA Approval No: 61234/136807



|             |                      |
|-------------|----------------------|
| Label Name: | BOXER GOLD HERBICIDE |
|-------------|----------------------|

|                  |   |
|------------------|---|
| Signal Headings: | POISON<br>KEEP OUT OF REACH OF CHILDREN<br>READ SAFETY DIRECTIONS BEFORE OPENING OR USING |
|------------------|---|

|                         |   |
|-------------------------|---|
| Constituent Statements: | 800 g/L PROSULFOCARB<br>120 g/L S-METOLACHLOR |
|-------------------------|---|

|                 |                    |
|-----------------|--------------------|
| Mode of Action: | GROUP 15 HERBICIDE |
|-----------------|--------------------|

|                      |   |
|----------------------|---|
| Statement of Claims: | For control of Annual Ryegrass ( <i>Lolium rigidum</i> ) and other grass and broadleaf weeds in Barley, Chickpeas, Faba Beans, Field Peas, Lentils, Lupins, Oats, Potatoes, Triticale and Wheat |
|----------------------|---|

|               |          |
|---------------|----------|
| Net Contents: | 20-2500L |
|---------------|----------|

|             |                     |
|-------------|---------------------|
| Restraints: | Refer to attachment |
|-------------|---------------------|

|                     |                     |
|---------------------|---------------------|
| Directions for Use: | Refer to attachment |
|---------------------|---------------------|

|                    |  |
|--------------------|--|
| Other Limitations: |  |
|--------------------|--|

|                      |   |
|----------------------|---|
| Withholding Periods: | Barley, Chickpeas, Faba Beans, Field Peas, Lentils, Lupins, Oats, Triticale, Wheat<br>Harvest: NOT REQUIRED WHEN USED AS DIRECTED<br>Grazing: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER APPLICATION<br>Potatoes: NOT REQUIRED WHEN USED AS DIRECTED |
|----------------------|---|

|               |  |
|---------------|--|
| Trade Advice: |  |
|---------------|--|

|                       |  |
|-----------------------|--|
| General Instructions: | <p>BOXER GOLD Herbicide is a short residual, soil applied herbicide that is absorbed by the roots and shoots (coleoptile) of germinating seedlings with inhibition of growth in the meristematic region. Foliar uptake is possible but of lower effect. Upstream translocation in the plant occurs but movement in the phloem is very unlikely. Best activity can be expected from applications before or shortly after the germination of target weeds. For pre-emergent application, the product should be applied to a moist smooth seedbed free of soil clods and emerged weeds. Product efficacy may be reduced by improper incorporation, high levels of crop or trash residues, stones or foreign matter and in areas of extremely high weed seed density such as header rows. For optimal efficacy and crop safety in barley, oats, triticale and wheat, avoid shallow seeding and ensure accurate seed placement that avoids placement of seed in the herbicide band. Avoid water repellent soils, soils subject to water logging or where furrow walls are likely to collapse. Duration and effectiveness of weed control will depend on use rate, soil type and rainfall or irrigation after application. Adequate rainfall or irrigation should occur within 10 days of application, as this herbicide requires sufficient available soil moisture to ensure soil movement and uptake by emerging weed seeds.</p> <p>For early post-emergent application in wheat and barley to suppress annual ryegrass, the product should be applied to a moist soil. Annual Ryegrass should be free of stress from marginal moisture or from water-logging between crop planting and application. BOXER GOLD is less effective on larger emerged weeds and application to Ryegrass which has commenced tillering should be avoided. Follow-up rainfall within 14 days of application will be beneficial to herbicide activity as it will move the herbicide into the soil around the shoot base and roots where it is most effective. Mild, transient crop effects may occur with wheat and barley after post-emergent applications. On sandy soils, if substantial rain occurs within the week following application crop phytotoxicity may be more severe.</p> <p>A registered post-emergent selective graminicide may be required for the control of later germinating Annual Ryegrass cohorts, especially in pulse crops. BOXER GOLD should be used as part of an integrated weed management program that utilises all available weed control tactics. Consider using a registered selective spray-topping treatment such as GRAMOXONE 360 Pro Herbicide in pulse crops to reduce weed seed set.</p> |
|-----------------------|--|

### Crop Tolerance

The selectivity of BOXER GOLD is a combination of positional and physiological selectivity. Positional selectivity must be maintained by sowing at an adequate depth below the herbicide band. The physiological selectivity of BOXER GOLD is given by the ability of cereals to metabolise the herbicide to inactive compounds more rapidly than susceptible weed species. Environmental factors such as rainfall events following application and soil type will influence product movement into the seed zone. Crop injury may occur when used in sandy soils with high leaching potential or in all soil types when heavy rainfall is received between sowing and emergence. Crop injury may also result from tank mixes with other herbicides. Shallow seeding is not recommended due to the greater potential for movement of herbicide within close proximity of the emerging crop. Conditions resulting in poor root development or the occurrence of crop stresses including waterlogging, drought, frost, nutrient deficiency or disease can result in unacceptable crop damage and yield loss as a result of root uptake of herbicide. Although crop tolerance has been clearly demonstrated in a wide range of registered crop cultivars, differences in relation to the tolerance of recent and future release cultivars may exist. Risk of crop damage may be exacerbated where varieties with short coleoptile length are planted in conjunction with the use of some fungicide seed treatments. In general pulse crop tolerance is greatest in Chickpeas>Lentils>Lupins>Faba Beans>Field Peas (most susceptible). Consult your local agronomist, advisory service or National Variety Trial (NVT) herbicide tolerance screening results for the latest information of specific varietal tolerance.

### Tillage System

IBS with knife or blade points: Working speed should ensure adequate incorporation of product but avoid soil throw into the adjacent seeding row. Use of presswheels will minimise potential for herbicide to be dragged back into seeding rows. Weed control may be reduced in seeding rows as a result of concentration of herbicide in crop inter-rows. A knife or blade point is defined as being <12 mm in width, having no wings, inverted T or blade and is generally on a minimum 200 mm row spacing.

Full disturbance or conventional cultivation, including trailing harrows/prickle chain: Care should be taken to ensure seed placement below the herbicide band as crop damage may result where seed is sown too shallow. Variable efficacy may be experienced with seeding systems using anything other than knife point and press wheels. Consult trifluralin label for tank mix rate selection.

**PRODUCT MUST BE INCORPORATED INTO THE SOIL WITHIN 7 DAYS OF APPLICATION**

### Mixing

BOXER GOLD is an emulsifiable concentrate that mixes readily with water. Fill the spray tank to one quarter full. Add BOXER GOLD and continue adding water to make up to the final spray volume. Agitate while mixing and spraying.

When tank mixing, wettable powder or water dispersible granule formulations should be added to the tank first followed by suspension concentrates (flowables), water soluble salts then BOXER GOLD or other emulsifiable concentrate formulations. Maintain thorough agitation during mixing and application. Agitate tank mixes vigorously if allowed to stand.

Note: Tank mix spray solutions should NOT be left standing in the vat overnight.

### Sprayer Clean Up

After using BOXER GOLD, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

### Compatibility

Physical compatibility has been assessed for a range of products and, providing the correct mixing order is followed and accompanied with strong agitation of the spray solution, compatibility will be acceptable. Refer to your local Syngenta representative for the most up to date information relating to the compatibility and crop safety of herbicide tank mixtures or visit our website at [www.syngenta.com.au](http://www.syngenta.com.au).

As formulations of other manufacturers' products are beyond the control of Syngenta and water quality varies with location, all mixtures should be tested prior to mixing commercial quantities.

#### Application

Barley, Chickpeas, Faba Beans, Field Peas, Lentils, Lupins, Oats, Triticale, Wheat: Apply by ground rig only in a minimum of 50 L water/ha. Water rate selection should be based on soil type and stubble load. Stubble loads above 40 to 50% ground coverage can reduce weed control below acceptable levels. Water volumes greater than 70 L/ha are recommended in order to reduce the impact of stubble in direct drill or minimum tillage systems. Use a nozzle delivering spray quality in the coarse spray range.  
Potatoes: Apply by ground rig only in a minimum of 100 L water/ha. Use a nozzle delivering spray quality no smaller than a coarse droplet size.

#### Resistance Warning:

##### Resistant Weeds Warning - GROUP 15 HERBICIDE

BOXER GOLD Herbicide is a member of the thiocarbamates and  $\alpha$ -chloroacetamides groups of herbicides. This product has the inhibitors of very long chain fatty acids synthesis modes of action. For weed resistance management, this product is a Group 15 herbicide. Some naturally occurring weed biotypes resistant to this product and other Group 15 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 15 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

##### Resistance Management

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the surviving weeds. DO NOT make more than 1 application of a Group 15 herbicide with the inhibition of very long chain fatty acid synthesis mode of action to a crop in the same season, excluding use of a split application at planting as per Directions for Use.

If the user suspects that the target weed population is resistant to herbicides with this mode of action, BOXER GOLD or other Group 15 herbicides should not be used. Strategies to minimise the risk of herbicide resistance are available. The above recommendations should be incorporated into an Integrated Weed Management (IWM) Program that uses a diverse range of options to manage grass weeds and avoids over reliance on any 1 method of control. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Syngenta representative for details.

##### Tank Mixes

Application of BOXER GOLD at less than 2.5 L/ha when tank mixing for Annual Ryegrass control WILL NOT be effective in the control of Group 3 resistant populations nor to delay the onset of herbicide resistance development. Tank mixing of herbicides is only effective in managing resistance where a lethal dose rate of each herbicide is applied. Whilst a tank mixture of 2 herbicides of differing modes of action at sub-lethal dose rates may still provide a high level of weed control, the risk of developing resistance to multiple modes of action is increased. Tank mixtures of BOXER GOLD with trifluralin are only recommended where targeting additional weed species, on sandy soils where soil moisture is limiting or where product movement into the seeding zone poses a crop safety risk.

#### Precautions:

Re-entry Period: DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

#### Protections:

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS  
DO NOT sow susceptible crops within 6 months of herbicide application.

|  |  |
|--|--|
|  | <p>DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.<br/> DO NOT allow spray to drift onto adjacent fallow land.<br/> DO NOT apply on or near shrubs, trees, lawns or crops other than barley, chickpeas, faba beans, field peas, lentils, lupins, potatoes or wheat.<br/> DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend, or in situations where by movement of soil or by seepage absorption of the herbicide may occur.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT<br/> HIGHLY TOXIC TO AQUATIC ORGANISMS. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.</p> <p>.</p> |
|--|--|

|                       |  |
|-----------------------|--|
| Storage and Disposal: | <p>Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.</p> <p>Refillable containers<br/> Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.</p> <p>Non-refillable containers<br/> Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.</p> |
|-----------------------|--|

|                    |   |
|--------------------|---|
| Safety Directions: | <p>Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear:</p> <ul style="list-style-type: none"> <li>• cotton overalls, over normal clothing, buttoned to the neck and wrist</li> <li>• washable hat</li> <li>• elbow-length chemical resistant gloves</li> <li>• face shield or goggles.</li> </ul> <p>When using the prepared spray wear:</p> <ul style="list-style-type: none"> <li>• cotton overalls, over normal clothing, buttoned to the neck and wrist</li> <li>• washable hat</li> <li>• elbow-length chemical resistant gloves.</li> </ul> <p>If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After each day's use wash gloves, face shield or goggles and contaminated clothing. Wash hands after use.</p> |
|--------------------|---|

|                         |   |
|-------------------------|---|
| First Aid Instructions: | If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126. |
|-------------------------|---|

|                     |  |
|---------------------|--|
| First Aid Warnings: |  |
|---------------------|--|

## RESTRAINTS

DO NOT apply to vetch or other pulse crops not listed.  
DO NOT use in seeding/tillage systems that cannot ensure accurate seed placement and adequate spatial separation of seed and herbicide.  
DO NOT apply to soils prone to waterlogging, sodic soils or soils affected by physical compaction.  
DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 2 days of application.  
DO NOT irrigate treated fields to the point of run off within three days of application  
DO NOT apply to potato fields where slopes exceed 4%  
Potatoes: DO NOT apply more than once per crop  
Wheat, Barley: DO NOT apply more than 2.5 L/ha per single growing season when applied pre-emergent, IBS or as a split application.  
DO NOT apply more than 3.0 L/ha per single growing season when applied post-emergent  
Oats, Triticale: DO NOT apply more than 2.5 L/ha per single growing season when applied pre-emergent  
DO NOT apply by aircraft.

## SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

**DO NOT** allow bystanders to come into contact with the spray cloud.

**DO NOT** apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

**DO NOT** apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

**DO NOT** apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**DO NOT** apply by a boom sprayer unless the following requirements are met:

- Spray droplets not smaller than a **COARSE** spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

### Buffer zones for boom sprayers

| Application rate         | Mandatory downwind buffer zones |                  |
|--------------------------|---------------------------------|------------------|
|                          | Natural aquatic areas           | Vegetation areas |
| Up to maximum label rate | 20 metres                       | 10 metres        |

## DIRECTIONS FOR USE

| Crop  | Weeds   | Rate/ha   | Critical Comments   |
|---|---|---|---|
| <b>Barley,<br/>Wheat,<br/>Oats,<br/>Triticale</b> | Annual Ryegrass ( <i>Lolium rigidum</i> ) including control of Group 3 resistant populations, Silver Grass ( <i>Vulpia</i> spp.), Stone Crop ( <i>Crassula</i> spp.)<br><b>Suppression</b> of Barley Grass ( <i>Hordeum</i> spp.) | 2.5 L   | <p>Apply pre-emergent and incorporate mechanically by sowing operation (IBS). Application should be made to a moist seedbed up to 7 days prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse. Refer to crop tolerance, incorporation and tillage requirements under GENERAL INSTRUCTIONS.</p> <p><b>Barley Grass Management</b><br/>           Where a high weed density is expected, delay application and sowing until after the emergence of the first Barley Grass germination following planting rains. Control with cultivation or the application of a registered non-selective herbicide such as glyphosate. Where sowing following a pasture phase, it is recommended that spray topping with GRAMOXONE® 360 Pro Herbicide or glyphosate is carried out in the previous year to reduce Barley Grass seed set.</p> |
| <b>Barley,<br/>Wheat</b>                          |   | 1.75 L pre-plant followed by 750 mL post plant pre-emergent | <p>Apply pre-plant and incorporate mechanically by the sowing operation (IBS). Application should be made to a moist seedbed up to 7 days prior to planting. Follow with the post-plant application as soon after sowing as possible, but before the crop and weeds emerge.</p> <p>Sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after each application. At planting, avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse.</p> <p><b>Barley Grass Management:</b> Refer to IBS recommendation</p>  |

| Crop                                     | Weeds  | Rate/ha   | Critical Comments  |
|--|--|---|--|
| <b>Barley, Wheat</b><br><i>continued</i> | Suppression of Annual Ryegrass ( <i>Lolium rigidum</i> ) (1 to 3 leaf growth stage)  | 2.5 to 3.0 L  | <p>The use of BOXER GOLD® is intended as part of an integrated Annual Ryegrass management strategy that includes varied means to reduce ryegrass survival and seed-set, to sustain low seed bank numbers. Early post-emergent use of BOXER GOLD® should not be the primary means of managing ryegrass numbers. Surviving plants not controlled by BOXER GOLD® may be stunted and uncompetitive but may still set seed. Apply only to Annual Ryegrass growing where the soil moisture profile and growing conditions since planting have been good. A follow-up rainfall within 14 day of application is required to achieve a high level of weed suppression.</p> <p>Use the higher rate under moderate to high ryegrass densities. Use of the higher rate will not restore efficacy where ryegrass has been stressed or where soil moisture conditions are not conducive to good activity.</p> <p><b>DO NOT apply BOXER GOLD® after growth stage Z25 (GS25).</b></p> <p><b>DO NOT apply BOXER GOLD® post-emergent to crops that have been treated with a pre-emergent or split application of BOXER GOLD®.</b></p> <p><b>DO NOT apply to crops that are water-logged or may be subject to water-logging as unacceptable crop injury may result.</b></p> <p><b>DO NOT apply to Annual Ryegrass that has been stressed due to either a period of marginal moisture or water-logging as the level of weed control may be reduced.</b></p> <p><b>DO NOT apply to crops under stress, such as herbicide damage, due to increased risk of further crop injury.</b></p>                          |
|  | Annual Ryegrass ( <i>Lolium rigidum</i> ), Paradoxa Grass (Canary Grass), <i>Phalaris</i> spp., Red and White Fumitory, Sand Fescue, Silver Grass ( <i>Vulpia</i> spp.), Stone Crop ( <i>Crassula</i> spp.), soil surface Wild Oats, Wireweed (Hogweed)<br><br><b>Suppression of</b> Barley Grass ( <i>Hordeum</i> spp.), Brome Grass, Deadnettle, Rough Poppy, Yellow Burr Weed | 1.5 to 2.5 L plus 0.8 to 1.5 L of a 480 g/L trifluralin | <p>DO NOT use less than 2.5 L/ha BOXER GOLD® where Group 3 resistance is confirmed or suspected. Use of BOXER GOLD® below 2.5 L/ha alone or in a tank mix WILL NOT be effective in the control of Group 3 resistant Annual Ryegrass nor to delay the onset of herbicide resistance development. Refer to Tank Mixes under GENERAL INSTRUCTIONS.</p> <p>Apply pre-emergent and incorporate mechanically by sowing operation (IBS). Application should be made to a moist seedbed up to 24 hours prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse.</p> <p>Use 1.5 L/ha of a 480 g/L trifluralin in minimum tillage knife/blade point systems only as per trifluralin label. Attention to sowing speed and soil throw is important to ensure crop safety. This is especially critical at higher use rates. Risk of crop injury may increase where greater than 1 L/ha trifluralin 480 g/L is applied with 2.5 L/ha BOXER GOLD®. Always refer to use recommendations on trifluralin label regarding soil type restrictions and incorporation timing and rate selection under different tillage systems.</p> <p>This mixture is recommended for control or suppression of additional weeds, on sandy soils where product movement into the seeding zone poses a crop safety risk or where good soil moisture cannot be assured. Refer to crop tolerance, incorporation and tillage requirements under GENERAL INSTRUCTIONS.</p> |



| Crop                                     | Weeds  | Rate/ha                                | Critical Comments   |
|--|--|--|---|
| <b>Barley, Wheat</b><br><i>continued</i> | Toad Rush<br>( <i>Juncus bufonius</i> )  | 1.25 to 2.5 L                          | Apply pre-emergent and incorporate mechanically by the sowing operation (IBS). Application should be made to a moist seedbed up to 7 days prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse.<br>Use upper label rate where longer residual control is required or in situations where crop competition is minimal. Refer to crop tolerance, incorporation and tillage requirements under GENERAL INSTRUCTIONS.   |
| <b>Wheat</b>                             | Annual Ryegrass ( <i>Lolium rigidum</i> ),<br>Silver Grass ( <i>Vulpia</i> spp.),<br>Stone Crop ( <i>Crassula</i> spp.),<br>Wild Turnip<br><b>Suppression</b> of Barley Grass,<br>Brome Grass  | 2.5 L plus<br>25 g Monza*<br>Herbicide | Apply pre-emergent and incorporate mechanically by sowing operation (IBS). Application should be made to a moist seedbed prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse. Refer to crop tolerance, incorporation and tillage requirements under GENERAL INSTRUCTIONS. Observe crop rotation guidelines detailed on Monza* label.   |
| <b>Potatoes</b>                          | Annual Ryegrass including control of Group 3 resistant populations,<br>Barnyard Grass,<br>Blackberry Nightshade,<br>Fat Hen, Fumitory,<br>Glossy Nightshade,<br>Redroot Amaranth,<br>Summer Grass,<br>Toad Rush<br><b>Suppression</b> of Common Thornapple,<br>Fierce Thornapple | 4 to 5 L                               | Apply after planting, after the first cultivation but no later than 25% potato shoot emergence. Application should be made to moist soil. Good soil moisture in the top 3 to 5 cm of soil is required for weed uptake to occur. Drying of the top soil can result in less effective weed control. Cultivation after herbicide application may be detrimental to weed control and crop safety if it results in the herbicide being unevenly distributed in the soil. Crop yellowing may occur where BOXER GOLD® is used on soils that have a low organic matter content and which contain more than 60% of fines and silt (crops will recover fully). On these soil types, use rates of BOXER GOLD® towards the lower end of the rate range. DO NOT apply more than once per crop. |
|  | Capeweed   | 5 L                                    | DO NOT apply after 25% potato shoot emergence. May be tank mixed with registered knockdown herbicide (eg SPRAY.SEED® 250 Herbicide) at recommended label rates and timing for control of additional weeds.  |

| Crop  | Weeds  | Rate/ha | Critical Comments   |
|---|--|---------|---|
| <b>Chickpeas,<br/>Faba Beans,<br/>Field Peas,<br/>Lentils,<br/>Lupins</b> | Annual Ryegrass<br>( <i>Lolium rigidum</i> ),<br>Silver Grass<br>( <i>Vulpia</i> spp.),<br>Stone Crop<br>( <i>Crassula</i> spp.),<br>Toad Rush<br>( <i>Juncus bufonius</i> ) | 2.5 L   | <p>Apply as a pre-emergent application to the soil surface up to 7 days prior to sowing and incorporate mechanically by the sowing operation (IBS). Application should be made to a moist seedbed up to 7 days prior to sowing and sufficient rain to thoroughly wet the top 3 to 4 cm of soil should occur within 10 days after spraying.</p> <p><b>Accuracy of seed placement is critical in ensuring crop selectivity. Unacceptable crop injury, including a reduction in crop vigour and yield loss, may occur where adequate positional selectivity of the herbicide is not maintained or where heavy rainfall occurs during the early stages of crop establishment. Avoid soil throw into adjacent seeding rows or sites where furrow walls may collapse. Shallow sowing is not recommended due to the greater potential for movement of herbicide within close proximity of the emerging crop, especially in sandy soils.</b></p> <p>Application of BOXER GOLD® to crops sown in soils of high leaching potential and those low in clay or organic matter may result in crop damage, especially in the event of heavy rainfall during the early stages of establishment. Conditions resulting in poor root development and the occurrence of crop stresses including waterlogging, drought, nutrient deficiency or disease may result in unacceptable crop damage where adequate positional selectivity of the herbicide has not been maintained.</p> <p><b>Avoid double spraying (over-lapping) of the crop with herbicide, including on headlands.</b></p> <p>Field Peas and Faba Beans have been found to be more susceptible to herbicide injury. Refer to crop tolerance, incorporation and tillage requirements under GENERAL INSTRUCTIONS.</p> <p>Late germinating weeds may not be adequately controlled and a post-emergent graminicide or registered spray-topping herbicide treatment (eg GRAMOXONE® 360 Pro) may be required to control later germinating weed cohorts or to reduce weed seed set.</p> |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**