



90 mm

**MACSPRED**<sup>®</sup>

**Simazine 900 DF**

**Herbicide**

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

**GROUP C HERBICIDE**

**APPROVED LABEL**

**Infopest Verified**

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

**GROUP C HERBICIDE**

For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

**MACSPRED PTY LTD**  
ACN 011 029 495  
13 Kennedys Drive  
Delacombe Vic 3356  
Telephone: (03) 5335 8522

**Net Contents: 500g**

\*Macspred Registered Trademark

210 mm

**STORAGE AND DISPOSAL**

Keep out of reach of children. Store in the closed, original container in a dry, well-ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty container into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**NOTICE TO BUYER**

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

APVMA Approval No.: 61396/500g/0510

Date of Manufacture:

Batch No:

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

MAC simaz 500g 04/10 4m

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**MACSPRED®**

**Simazine  
900 DF**

**Herbicide**

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

**GROUP C HERBICIDE**

For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

®Macspred Registered Trademark

**MACSPRED PTY LTD**  
ACN 011 029 495  
13 Kennedys Drive  
Delacombe Vic 3356  
Telephone: (03) 5335 8522  
www.macspred.com.au

**Net Contents:**



**5kg**



**10kg**



**15kg**



**STORAGE AND DISPOSAL**

Keep out of reach of children. Store in the closed, original container in a dry, well-ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**NOTICE TO BUYER**

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

APVMA Approval No.: 61396/5-15kg/1009

Date of Manufacture:

Batch No:

MAC-simaz 10/09 2m

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Macspred

# Simazine 900 DF

## HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg SIMAZINE

**GROUP C HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

**CONTAINS: 126 x 5 kgs MEASURE PACKS  
WHICH IT IS ILLEGAL TO SELL SEPARATELY**

NET CONTENTS

# 630 kg

MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522



### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, well ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional information is listed in the Material Safety Data Sheet.

### NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

In a Transport Emergency  
Dial 000  
Police or Fire Brigade

MacspreD Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/630kg/0309

Date of Manufacture:

Batch No.:



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Macspred Simazine 900 DF HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg SIMAZINE

**GROUP C HERBICIDE**

For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

CONTAINS: 144 x 5 kg MEASURE PACKS  
WHICH IT IS ILLEGAL TO SELL SEPARATELY

NET CONTENTS

## 720 kg

MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522



#### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, well ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional information is listed in the Material Safety Data Sheet.

#### NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

In a Transport Emergency  
Dial 000  
Police or Fire Brigade

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/720kg/0209

Date of Manufacture:

Batch No.:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# Macspred Simazine 900 DF HERBICIDE

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

**GROUP C HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

**NET CONTENTS**

# 10kg



**MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522**

### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, well-ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

**Additional information is listed in the Material Safety Data Sheet.**

### NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

**In a Transport Emergency  
Dial 000  
Police or Fire Brigade**

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/10kg/0508

Date of Manufacture:

Batch No.:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Macspred Simazine 900 DF HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg SIMAZINE

**GROUP C HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

**NET CONTENTS**

# 15kg



MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522

## STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, well-ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional information is listed in the Material Safety Data Sheet.

## NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

In a Transport Emergency  
**Dial 000**  
Police or Fire Brigade

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/15kg/0508

Date of Manufacture:

Batch No.:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Macspred Simazine 900 DF HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg SIMAZINE

**GROUP G HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

**CONTAINS: 3 x 5 kgs MEASURE PACKS  
WHICH IT IS ILLEGAL TO SELL SEPARATELY**

NET CONTENTS

# 15 kg

MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522



## STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, well-ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional information is listed in the Material Safety Data Sheet.

## NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

In a Transport Emergency  
Dial 000  
Police or Fire Brigade

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/15/0508

Date of Manufacture:

Batch No.:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**Macspred**

# Simazine 900 DF

**HERBICIDE**

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

**GROUP C HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

**CONTAINS: 100 x 5 kgs MEASURE PACKS  
WHICH IT IS ILLEGAL TO SELL SEPARATELY**

**NET CONTENTS**

**500 kg**

**MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522**



## **STORAGE AND DISPOSAL**

Keep out of reach of children. Store in the closed, original container in a dry, well ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional information is listed in the Material Safety Data Sheet.

## **NOTICE TO BUYER**

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

In a Transport Emergency  
**Dial 000**  
Police or Fire Brigade

Macspre Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/500A/0508

Date of Manufacture:

Batch No.:





**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**MACSPRED**  
**SIMAZINE 900 DF**  
**HERBICIDE**

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

Not to be sold separately.  
Before use read all directions on the outer pack.

**5 kg NET**

APVMA Approval No. 61396/5M/0508



425x190mm 100% on A3

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Macspred

# Simazine 900 DF

## HERBICIDE

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

**GROUP C HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

**CONTAINS: 50 x 10 kgs MEASURE PACKS  
WHICH IT IS ILLEGAL TO SELL SEPARATELY**

**NET CONTENTS**

# 500 kg

MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522



### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated locked area, out of direct sunlight. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/500B/0508

Date of Manufacture:

Batch No.:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**MACSPRED**  
**SIMAZINE 900 DF**  
**HERBICIDE**

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

Not to be sold separately.  
Before use read all directions on the outer pack.

**10 kg NET**

APVMA Approval No. 61396/10M/0508



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Macspred

# Simazine 900 DF

HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg SIMAZINE

**GROUP C HERBICIDE**

**For the control of weeds in a range of horticultural and broadacre crops, forestry and in non-crop situations as per the Directions for Use table.**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

**CONTAINS: 74 x 6.7 kgs MEASURE PACKS WHICH IT IS ILLEGAL TO SELL SEPARATELY**

NET CONTENTS

# 495.8 kg

MACSPRED PTY LTD  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 5335 8522



#### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, well-ventilated locked area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilizers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional information is listed in the Material Safety Data Sheet.

#### NOTICE TO BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

In a Transport Emergency  
Dial 000  
Police or Fire Brigade

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522

APVMA Approval No. 61396/495.8/0508

Date of Manufacture:

Batch No.:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**MACSPRED**  
**SIMAZINE 900 DF**  
**HERBICIDE**

**ACTIVE CONSTITUENT: 900 g/kg SIMAZINE**

Not to be sold separately.  
Before use read all directions on the outer pack.

**6.7 kg NET**

APVMA Approval No. 61396/6.7M/0508



**70x100mm 100% on A4**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**MACSPRED<sup>®</sup>**

**SIMAZINE 900 DF HERBICIDE**

**ACTIVE CONSTITUENT : 900g/kg SIMAZINE**

**GROUP C HERBICIDE**




For the control of weeds in a range of horticultural and broad acre crops, forestry and in non-crop situations as per the directions for use table.

**READ THIS LEAFLET BEFORE USING THIS PRODUCT**

Macspred Pty Ltd  
ACN 011 029 495  
No. 13 KENNEDYS DRIVE  
DELACOMBE VIC 3356  
TEL: 03 53358522




IG



**DIRECTIONS FOR USE**  
**FRUIT AND VEGETABLES**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
Almonds	SA only	REFER TO WEED TABLE
Asparagus	All States	
Berry Fruits. Boysenberries, Currants, Loganberries, Raspberries established for 12 months		
Citrus established for 12 months		
Hops		
Pome Fruit, Apples, Pears	Qld only	
	NSW, ACT, Vic, SA, WA, Tas only	





RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
1 kg or 60g/100L	2 kg or 120g/100L	Use only if trees are at least 3 years old. Apply to bare, moist soil immediately after cultivation and before weeds emerge.
1.3 kg	2.5 kg	Apply to bare, moist soil prior to spear emergence.
		Do NOT apply to foliage or when fruit is present. Apply to bare, moist soil on established plants only.
		Apply to bare moist soil immediately after cultivation and before weeds emerge.
		Apply to bare, moist soil in late winter, early spring, before hop emergence. Hops should be covered by 50 mm of soil.
3 kg	4 kg	Use only if trees are at least 2 years old. Apply to bare soil.
1.8 kg/ha or 90g/100L	2.5 kg/ha or 120g/100L	



## FRUIT AND VEGETABLES cont...

CROP SITUATION	STATES	WEEDS
Strawberries	All States	REFER TO WEED TABLE
Vineyards	Qld only	
	NSW, ACT, Vic, SA, WA, Tas only	




RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
1.2 kg	1.2 kg	Apply between polythene covered beds. DO NOT use on white or grey sands in WA.
3.9 kg	3.9 kg	Use only if vines are at least 3 years old.
1.3 kg or 60g/100L	2.5kg/ha or 120g/100L	Use only if vines are at least 12 months old. In the first year of use, split applications are preferred. DO NOT use on excessively sandy soils, as crop damage may occur.



## FIELD CROPS AND PASTURES

CROP SITUATION	STATES	WEEDS
<p><b>Canola</b> (Triazine Tolerant varieties only) <b>Pre emergence or post sowing pre emergence only</b></p>	All States	<p>Capeweed, Charlock, Clover, Corn Cromwell, Doublegee, Fumitories, Geraniums, Ivy-leaf Speedwell, London Rocket, Mustards, , Turnips, Paterson's Curse, Shepherd's Purse, Silver Grass (Vulpia).</p> <p><b>Suppression</b> of Annual Ryegrass, Barley Grass, Brome Grass, Wild Oats and Wild Radish</p>
Chickpeas	NSW, Qld, ACT, SA, only	<p>Dead nettle, Indian Hedge Mustard, Lesser Swinecress, Milk Thistle, Prickly Lettuce, Purple Goosefoot, Shepherd's Purse, Turnip Weed, Wireweed and suppression of Black Bindweed and Paradoxa grass</p>



RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
1.1 - 2.2 kg		<p><b>This use is subject to an Integrated Weed Management Strategy for the use of triazine herbicides in triazine tolerant (TT) canola.</b></p> <p><b>See General Instructions: Integrated Weed Management Strategy for TT-Canola.</b></p> <p>Can be applied up to a week before sowing or post-sowing pre-emergence (ideally incorporated by harrows). For best results, apply to bare moist soil, either immediately before seeding or as a pre-emergence treatment at or within 7 days of planting. Sufficient rainfall (20-30 mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Application should not be made to ridged or excessively cloddy soil. When applied before seeding, incorporate to a depth of 5 cm.</p>
800g + 830g of 900g/kg prometryn		<p>Apply immediately post-planting. Application should not be made to ridged or excessively cloddy soil. For reliable results, significant rain (20-30mm) is necessary within 2-3 weeks of sowing.</p>




**FIELD CROPS AND PASTURES cont...**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
Chickpeas	Qld, NSW, ACT, Vic, SA only	Milk Thistle (Common Sowthistle), Indian Hedge Mustard, Turnip weed and suppression of Prickly Lettuce, Shepherd's Purse and Wireweed.
	Qld, NSW, ACT, Vic, SA, WA only	Annual ryegrass, Barley Grass, Capeweed, Charlock, Corn Gromwell, Fumitories, Geranium, Ivy-leaf Speedwell, Rough Poppy, Mustards, Turnips, Volunteer Canola (not triazine tolerant cultivars), Wireweed and suppression of Brome Grass and Wild Oats.



RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
800g – 1.1 kg		<p>For best results, apply to bare moist soil, immediately post planting. Sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application.</p> <p>Applications should not be made to ridged or excessively cloddy soil. Use the lower rate on light sandy soils.</p>
800g – 1.1 kg plus 800mL of 480 g/L trifluralin		<p>Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Use this mixture where annual Ryegrass and Wild Oats are the major problem.</p> <p>Sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application.</p> <p>Application should not be made to ridged or excessively cloddy soil. Use the lower rate on light sandy soils.</p>



**FIELD CROPS AND PASTURES cont...**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
Chickpeas	WA only	Capeweed, Clover, Dock, Doublegee Mustard, Radish, Self Sown Cereals, Silver Grass, Turnip Weed and suppression of Barley Grass, Ryegrass and Wild Oats.
Faba beans	NSW, ACT, Vic, SA, WA only	Annual Ryegrass, Barley Grass, Brome Grass, Capeweed, Corn Gromwell, (Sheepweed), Deadnettle, Fumitories, Geranium, Ivy-leaf Speedwell, Medics, Mustards, Paradoxa Grass, Saffron Thistle, Soursob, Volunteer Canola, Wireweed and suppression of Wild Oats






RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
550g to 1.1kg		<p>For best results, apply to bare moist soil, either immediately before seeding, or as a pre-emergent treatment at or within 7 days of planting. Sufficient rainfall to wet the soil through the weed root zone (20-30 mm) is necessary within 2-3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing, and Chickpeas are sown into a dry or low moisture seedbed.</p> <p>Application should be made to ridged or excessively cloddy soil. When applied before seeding, incorporation by the sowing operation should not be greater than 5 cm. Use 0.55 – 0.75 kg on lighter soils and in the Northern Agricultural areas, and up to 1.1 kg on heavier soil types. DO NOT use on whitish or grey sands.</p>
1.1 kg	1.4 kg	<p>Apply either pre-seeding or immediately post-sowing which is preferred on light soils. Sow the crop at least 5cm deep. Use the lowest rate on light soils. Application should not be made to ridged or excessively cloddy soil. For fully reliable results, significant rainfall (20-30 mm) is necessary within 2-3 weeks of sowing. Do not use rates higher than 1.1 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur.</p>



## FIELD CROPS AND PASTURES cont...

CROP SITUATION	STATES	WEEDS
Faba beans	NSW, ACT, Vic, SA, WA only	Annual Ryegrass, Barley Grass, Brome Grass, Capeweed, Corn Gromwell, (Sheepweed), Deadnettle, Fumitories, Geranium, Ivy-leaf Speedwell, Medics, Mustards, Paradoxa Grass, Saffron Thistle, Soursob, Volunteer Canola, Wireweed and suppression of Wild Oats
Lucerne (established)	Tas only	REFER TO WEED TABLE
Lupins	NSW, ACT, Vic, SA, Tas only	Annual Ryegrass, Barley Grass, Capeweed, Fumitories, Wireweed, Corn Gromwell (Sheep Weed), Ivy-Leaf Speedwell, Turnips, Mustards, Geraniums, suppression of Wild Oats, Brome Grass, and Soursob.




RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
550g plus 800mL of a 480g/L trifluralin	800g plus a 800mL of a 480g/L trifluralin	Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within hours of application. Use this mixture where Annual Ryegrass and Wild Oats are the major problem. Application should not be made to ridged or excessively cloddy soil. For fully reliable results. Significant rainfall (20-30mm) is necessary with 2-3 weeks of application.
1.1 kg	1.1kg	Use only if lucerne is more than 12 months old. Apply during the winter dormant period before weeds emerge.
800g – 1.1 kg	1.3 – 2.2 kg on loam soils	Apply 50 – 100 L spray mixture/ha. Apply to bare moist soil at or within 2 days of sowing. Application should not be made to ridged or excessively cloddy soil. Use the highest rate on heavy soil or where free organic matter such as trash is present. If dry conditions prevail after application, incorporation with light harrows up to one week after sowing may assist results.




## FIELD CROPS AND PASTURES cont...

CROP SITUATION	STATES	WEEDS
Lupins When no weeds are present at time of sowing	WA only	Capeweed, Clover, Dock, Doublegee, Mustard, Radish, Self Sown Cereals, Silver Grass, Turnip and suppression of Barley Grass, Brome Grass, Ryegrass and Wild Oats

LIGH  
SOIL550 g  
1.1 k550g  
1.1 kg  
1.25L  
480 g  
Liflure

RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
550 g - 1.1 kg	1.1 kg - 1.6kg on gravelly loam soils	Can be applied as a pre-emergent treatment before or within two days of planting. Application should not be made to ridged or excessively cloddy soil. For fully reliable results significant rainfall, e.g. 10-20 mm is necessary within 2-3 weeks of application <b>Note: - Do not use on white or grey sands.</b>
550g – 1.1 kg + 1.25L of 480 g/L trifluralin	550g – 1.1 kg + 1.25L of 480 g/L trifluralin	For best results apply the tank mix to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Application should not be made to ridged soil. Incorporation should be made within 4 hours of application. Use as a pre-emergence application only. Use this mixture where Annual Ryegrass and Wild Oats are the major problems. Sufficient rainfall to wet the soil through the weed root zone (20 to 30mm) is necessary within 2 – 3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and lupins are sown into dry or low moisture seedbed. Use 1 to 1.5 L/ha on yellow sands, 2L/ha on all other soil types. Where Brome Grass is a problem use 2 L/ha. DO NOT use on whitish or grey sands.




## FIELD CROPS AND PASTURES cont...

CROP SITUATION	STATES	WEEDS
Lupins Where weeds are present at time of sowing		
Lupins Post-emergence top up application following a pre-emergence application of Simazine		

LIGH  
SOIL550 l  
800g  
100 l  
160n  
of  
250g  
paraq400g  
1.1 k

RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
550 to 800g + 100 to 160mL of 250g/L paraquat	550 to 800g + 100 to 160mL of 250g/L paraquat	For best results, apply to bare moist soil 1 to 6 days prior to seeding in areas where the crop will be sown under conservation tillage system. Sufficient rainfall to wet the soil through the weed root zone (20 to 30 mm) is necessary within 2 to 3 weeks of application. Results can be variable if seasonal conditions are dry prior to sowing and Lupins are sown into a dry or low moisture seedbed. Use the lower rate when weeds have emerged for more than 2 weeks and the higher rate when application is made within 2 weeks of weed emergence. DO NOT use on whitish or grey sands.
400g – 1.1 kg	400g – 1.1 kg	To be used in addition to a pre-emergence application of Macspred Simazine 900 DF Herbicide, but not as an alternative to a pre-emergence application. Apply to moist soil. Use rates of 400g to 550g within 4 weeks after seeding. DO NOT apply to Lupins which are showing symptoms of simazine damage from pre-emergence application.



## FIELD CROPS AND PASTURES cont...

CROP SITUATION	STATES	WEEDS
Sub clover Established lucerne and perennial grass pastures	NSW, Vic, ACT, SA only	Vulpia (Silvergrass) Rat's tail fescue, Squirrel-tail Fescue, Sand Fescue
	NSW, ACT, Vic only	
	SA only	

LIGH  
SOIL

50

550  
100ml  
a 25040  
10  
of 250



RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
550 - 900 g		Apply 6-10 weeks after emergence of the pasture, but not before the 3rd trifoliolate leaf stage of Sub Clover. Best results are obtained from application to young, well grazed, and actively growing plants. Some damage to Sub Clover may occur especially at the higher rates, DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist soil conditions; rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and sub clover allowed to recover before the Simazine 900DF is applied. DO NOT use herbicides for broad-leaved weed control within 3 weeks of application.
550 to 800g plus 100mL to 160mLs of a 250 g/L paraquat.		Apply 6-10 weeks after emergence of the pasture, but not before the 3rd trifoliolate leaf stage of sub clover. Best results are obtained from application to young, well grazed, and actively growing plants. However, the addition of paraquat improves the control of well established plants. Add a non-ionic surfactant at 0.2% v/v (200 mL per 100L). Under conditions of good soil moisture, control of other grasses and some broadleaf weed seedlings may occur. DO NOT apply to sandy soils and use the lower rates on light textured soils. Apply under moist conditions: rain following application enhances results. DO NOT apply to stressed plants. Insect infestations should be controlled and Sub Clover allowed to recover before the Simazine 900DF plus paraquat is applied. DO NOT use herbicides for broadleaved weed control within 3 weeks of application.
400 to 550g + 100 to 160mL of 250 g/L paraquat.		



**FIELD CROPS AND PASTURES cont...**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
Sub clover Established lucerne and perennial pastures	WA only	Vulpia (Silvergrass) Rat's tail fescue

**ORNAMENTALS**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
Gladioli	All states	REFER TO WEED TABLE
Roses	NSW, ACT, Vic, SA, WA, Tas only	
Nurseries and propagating beds	All states	



RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
400 - 550g		<p>Apply within 8 weeks of emergence of Sub clover and grasses. Under conditions of good soil moisture control of other grasses and some broadleaf weeds may occur. Ensure that there is a good stand of subterranean clover present before spraying. Use the lower rate on light textured soils. DO NOT use on deep white sands. DO NOT use broadleaf weed herbicides or add crop oils or wetting agents.</p> <p>DO NOT use on medics, or red or white clover.</p> <p>DO NOT overlap when spraying, otherwise damage may be observed.</p>

RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
1.2 kg or 60g/100L	1.2 kg or 60g/100L	Apply to bare, moist soil after planting. May cause damage on sandy soils low in organic matter.
1.3 kg	2.5 kg	Plants should be at least 1 year old. Apply to bare, moist soil.
1.2 kg	1.2 kg	Use only on non-susceptible plants listed in the table.




**NON-CROP USES**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
Non-crop uses, Commercial, Industrial, Rights of Way, Public Utility areas, Road shoulders, Drains, Headlands, Driveways, Railway tracks, Aerodromes, Gutters, Footpaths	All states	REFER TO WEED TABLE
Dams, Tanks, Troughs	WA only	Filamentous blue green algae



RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
5kg OR 500g/100L	5kg OR 500g/100L	Provides seasonal control of annual weed germinations. Apply to bare moist ground.
10kg OR 1kg/100L	24kg OR 2.4kg/100L	Provides long residual control in Winter rainfall areas. Apply to bare moist ground.
24kg OR 2.4kg/100L	48kg OR 4.8kg/100L	Provides long residual control in Summer rainfall areas. Apply to bare moist ground. Read remarks under Crop Safety.
2g/1000 L water		Mix in a convenient amount of water and apply when algae development is first noticed.



**FORESTRY**

<b>CROP SITUATION</b>	<b>STATES</b>	<b>WEEDS</b>
<p><b>Forestry</b> Including farm tree plantations <i>Pinus, Araucaria, Corymbia</i> and <i>Eucalyptus</i> tree plantings pre- and post-planting [see General Instructions for further information on tree species]</p>	All states	<p>Pre-emergence control of many annual and some perennial grasses and broadleaved weeds.</p>



RATE		CRITICAL COMMENTS
LIGHT SOILS	HEAVY SOILS	
1.6 - 6.7 kg/ha		<p>Macspreed Simazine 900DF Herbicide may be applied prior to or after planting.</p> <p>For most situations use 5.0 to 5.5 kg/ha. On gravelly coarse textured soils use 2.8-4.0 kg/ha. In WA &amp; SA in medium rainfall zones [400—600 mm] use 3.0-4.0 kg/ha and for sandy low organic soils use 1.6-3.0 kg/ha. On sandy soils allow two weeks and at least 50 mm of rain between spraying and planting. Use a maximum rate of 1.6 kg/ha for overspraying eucalyptus plantations on sandy, low organic matter soils in WA &amp; SA. For short term weed control or when applied with other pre-emergent herbicides use half rates according to the situation.</p> <p>Apply to bare ground. Best results will be achieved if applied to moist, finely tilled soil. If weeds are present add a knockdown or other partner herbicide as a tank mix. If spraying after planting check tree tolerance to knockdown or other partner herbicide before using. Application in a band 1.5-2.0 metres wide, over the planting row is recommended although broadcast application can also be used. Rates above refer to sprayed hectares and not field hectares.</p> <p>Some tree damage may occur, especially with post-plant application. Recovery is normally rapid. Directed application is preferred for post-planting application especially when the higher rates are used.</p>



**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**HARVEST**

**ALL CROPS: NOT REQUIRED WHEN USED AS DIRECTED**

**GRAZING:**

**CHICKPEAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 9 WEEKS AFTER APPLICATION**

**FABA BEANS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.**

**CANOLA : DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 15 WEEKS AFTER APPLICATION.**

**SUB CLOVER: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION.**

**DO NOT CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.**

**WEED :**

Amarar  
Poa (W  
Barley  
Barnya  
Bindii  
\*Brome  
Capew  
Chickw  
Comm  
Corn gr  
Creepir  
Fat her  
Fumitor  
Gerani  
Ivy-leat  
Mustar  
Native  
Nettles  
Potato  
Powell  
Red roc  
Red sh  
Shephe  
Summe  
Thistle

Turnips  
Turnip  
Wild m  
\*Wild r  
\*Wild t  
\*Wimm  
Yellow  
\*See W





RY TO  
ION.**WEED TABLE Weeds controlled in all States.**

Amaranths	<i>Amaranthus spp.</i>
Poa (Winter grass)	<i>Poa annua</i>
Barley grass Annual	<i>Hordeum spp</i>
Barnyard grass	<i>Echinochloa spp</i>
Bindii	<i>Calotis hispidula</i>
*Brome grass	<i>Bromus spp</i>
Capeweed	<i>Arctotheca calendula</i>
Chickweed	<i>Stellaria media</i>
Common sowthistle	<i>Sonchus oleraceus</i>
Corn gromwell (Iron/Sheepweed)	<i>Buglossoides arvensis</i>
Creeping oxalis	<i>Oxalis corniculata</i>
Fat hen	<i>Chenopodium album</i>
Fumitories	<i>Fumaria spp.</i>
Geranium	<i>Geranium spp</i>
Ivy-leaf speedwell	<i>Veronica hederifolia</i>
Mustards	<i>Sisymbrium spp</i>
Native geranium	<i>Geranium solanderi</i>
Nettles	<i>Urtica spp</i>
Potato weed	<i>Galinsoga parviflora</i>
Powell's amaranth	<i>Amaranthus powellii</i>
Red root amaranth	<i>Amaranthus retroflexus</i>
Red shank	<i>Amaranthus cruentus</i>
Shepherds purse	<i>Capsella bursa-pastoris</i>
Summer grass	<i>Digitaria spp</i>
Thistles – annual	<i>Carduus spp</i>
	<i>Carthamus spp</i>
	<i>Centaurea spp</i>
	<i>Onopordum spp</i>
	<i>Picnoman acarna</i>
	<i>Silybum marianum</i>
	<i>Brassica spp.</i>
	<i>Rapistrum rugosum</i>
	<i>Sisymbrium spp</i>
	<i>Raphanus raphanistrum</i>
	<i>Brassica tournefortii</i>
	<i>Lolium rigidum</i>
	<i>Oxalis corniculata</i>
- soldier	
- variegated	
Turnips	
Turnip weed	
Wild mustard	
*Wild radish	
*Wild turnip	
*Wimmera (Annual) ryegrass	
Yellow wood sorrel	

\*See WA suppression only table on following pages



**Weeds suppressed at higher rates**

Soursob	<i>Oxalis pes-caprae</i>
Wild oats	<i>Avena spp</i>

**Weeds suppressed only in WA**

Annual ryegrass	<i>Lolium rigidum</i>
Brome grass	<i>Bromus spp</i>
Dock	<i>Rumex spp</i>
Double gee	<i>Emex australis</i>
Radish	<i>Raphanus raphanistrum</i>
Sorrel	<i>Rumex acetosella</i>
Soursob	<i>Oxalis pes-caprae</i>
Wild turnip	<i>Brassica tournefortii</i>
Wireweed	<i>Polygonum aviculare</i>

**GENERAL INSTRUCTIONS**

Simazine 900DF is a pre-emergent herbicide which selectively kills plants and seedlings by absorption through the root system. Very little, if any, foliage absorption takes place. It works best when applied to bare, moist soil. If vegetation is present at spraying then a knockdown herbicide should be added. Simazine 900DF is very insoluble in water and so usually remains in the top 5 cm of the soil. It is inactivated by absorption onto charcoal or other carbonaceous materials, muck solid or soils high in organic matter - higher rates are therefore required in these types of soil.

***Araucaria, Corymbia, Eucalyptus and Pinus plantations:*** Field trials and use experience has indicated that the following species are tolerant. Test tolerance before treating other species. ***Araucaria*** – all commercial species, ***Corymbia*** – *maculata*. ***Eucalyptus*** – *agglomerata, calophylla, camaldulensis, cladocalyx, cloeziana, dunnii, globulus, grandis, laevopinea, nitens, pilularis, pseudoglobulus, regnans, rudis, saligna, wandoo*. ***Pinus*** – *caribaea* and hybrids, *elliottii, pinaster, radiata*.

**INTEG**The u:  
Manag  
Strate

- 
- 

Compl  
outline  
practic  
weeds  
volunt

Copies

It is ad  
consul  
DF Her

To mir

- 

Broad  
900g/l  
(if ryeg  
Broad

### INTEGRATED WEED MANAGEMENT STRATEGY FOR TT CANOLA

The use of this product in TT canola is subject to an Integrated Weed Management Strategy for the use of triazine herbicides in TT canola' The Strategy encompasses:

- Integrated Weed Management;
- the specific management of crop plant volunteers and outcrossing to other plants; and
- triazine herbicide residue management (agronomic and environment).

Compliance with registered label directions and adoption of the principles outlined in the Strategy will assist with implementation of crop management practices that **minimise** the development of herbicide resistance in treated weeds; **reduce** the levels of triazine residues in the environment; and **manage** volunteer plants and outcrossing.

Copies of the Strategy are available from a Macspred Pty Ltd representative.

It is advised that consultation be undertaken with an appropriate agronomist, consultant or Departmental adviser prior to the use of Macspred Simezone 900 DF Herbicide on TT Canola.

#### To minimise herbicide resistance:

- Avoid dry sowing in heavily weed infested paddocks. Wait for a weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage.
- Adapt the weed control program to the anticipated weed spectrum and pressure:

**Broadleaf Weeds and Ryegrass:** Use (*simazine 900g/kg*) or (*atrazine 900g/kg*) plus trifluralin pre-emergence. A follow-up with a Group A herbicide (if ryegrass is susceptible) or *atrazine 900g/kg* may be necessary.

**Broadleaf Weeds only:** Use *atrazine 900g/kg* post-emergence.



- **DO NOT** use *atrazine 900g/L* or *simazine 900g/kg* if the area to be treated had a triazine herbicide applied to it last season.
- Watch for escapes, especially in paddocks with a long history of Group C herbicide use.
- **DO NOT** use Group C herbicides in consecutive years.

### Resistant Weeds Reporting

Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine and simazine may be resistant, get them tested and seek professional advice.

### To avoid Triazine carry-over:

**On acid soils (pH less than 6.5)** - The maximum rate of atrazine 900g/kg or simazine 900g/kg or a combination of the two products to be applied to the crop during the growing season is **2kg a.i./ha**.

**On alkaline soils (pH greater than 6.5)** - The maximum rate of atrazine 900g/kg or simazine 900g/kg or a combination of the two products to be applied to the crop during the growing season is **1kg a.i./ha**.

– **Post-emergence use** - It is recommended that *atrazine 900g/L* only be used, and at rates of **1kg a.i./ha** or less, on both acid or alkaline soils.

## GROUP C HERBICIDE

### Resistant weeds warning

Simazine 900DF Herbicide is a member of the triazines group of herbicides. Simazine 900DF has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management Simazine 900DF is a Group C herbicide.

Some naturally occurring weed biotypes resistant to Simazine 900DF and other inhibitors of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used



repeat  
or other  
Since  
Macsp  
failure

Mixing  
This p  
equip  
produc  
mixing

1. Pou  
byp  
into

2. Afte  
DIS  
agil  
ope

N/B. S

Applic  
Use a  
hectar  
per he  
after u

TT-Ca  
DO NO  
with a  
lakes  
peren  
where

repeatedly. These resistant weeds will not be controlled by Simazine 900DF or other inhibitors of photosynthesis at photosystem II herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Macspred Pty Ltd accepts no liability for any losses that may result from the failure of Simazine 900DF to control resistant weeds.

### Mixing

This product must be mixed with water and applied by suitable spray equipment. Fill spray tank 60-80% full with clean water BEFORE adding the product. Begin agitating vigorously and continue agitation during the entire mixing and spraying operation.

1. Pour required amount of product steadily into the spray tank. Allow vigorous bypass agitation to completely disperse the product. DO NOT dump product into spray tank all at once.
2. After adding required quantity of product and obtaining COMPLETE DISPERSION continue to fill tank to desired level for spraying. Thorough agitation of the spray liquid should continue during the entire spraying operation.

N.B. Spray solution should NOT be left standing in the tank overnight.

### Application

Use a boom sprayer or knapsack for small areas. 50-200L of mixture per hectare should be applied with the boom and a minimum equivalent to 500L per hectare through the knapsack sprayer. Clean spray equipment immediately after use with a tank and equipment cleaner.

### TT-Canola: Application

DO NOT apply to TT-canola by aircraft. Apply only with a low boom sprayer with a 60 m buffer zone downwind of treated fields to natural or impounded lakes or dams, and a 20 m buffer zone for any well, sink hole, intermittent or perennial stream. Apply only to areas where run-off is unlikely to occur or where run-off may be captured by farm earthworks.



**Soil Moisture**

If sufficient rain to thoroughly wet the weed root zone has not fallen within 14 days of application, water in with the equivalent of 13mm of rain. If applied under dry conditions and irrigation is not available, mechanical incorporation to a depth of not more than 4cm within 7 days of sowing may assist results.

**Compatibility**

Simazine 900DF is compatible with the following products at the recommended label rates and dilutions: Fusion, Fusilade, Crop Care Shirquat 250, Gramoxone 250, Reglone, Spray.Seed 250, Crop Care Gladiator, Striker, Revolver, trifluralin, Hammer.

Always add the Simazine 900DF to the tank first as per the mixing instructions and ensure that it has completely dispersed before adding the other product.

**PRECAUTION****RE-ENTRY PERIOD**

DO NOT enter treated area without protective clothing until spray has dried.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT spray foliage of desired plants, except lupins (WA only).

DO NOT make more than two applications during any one year in Asparagus, Berry fruit, Citrus, Almonds, Gladioli, Hops, Apples, Pears, Roses, Vineyards and Lupins.

DO NOT use under Peach, Apricot, Plum trees, ornamental Prunus or near newly planted shrubs.

DO NOT plant crops other than those recommended on this label for at least nine months following treatments at rates up to 2.5kg/ha. When rates exceed 2.5kg/ha, plantings may not be possible for very long periods afterwards.

DO NOT use on sandy or porous soils near desirable plants or trees, or where roots of such may extend, or are near the surface.

Crops such as red beet, lettuce, cabbage, tomato, cucumber and carrots are particularly sensitive to simazine residues.



DO NO  
DO NO  
DO NC  
extend

**DRIFT**  
DO NO  
cause  
pastur

**PROTE**  
DO NO  
contain

**STORA**  
Keep c  
cont  
from c  
into sp  
shred  
availa  
marke  
roots.

DO NOT exceed highest rate recommended for any particular crop.  
 DO NOT use on white or grey sands (WA only).  
 DO NOT use in channels or drains where the roots of desirable plants may extend.

#### **DRIFT WARNING**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways, with the chemical or used containers.

#### **STORAGE AND DISPOSAL**

**Keep out of reach of children.** Store in the closed, original container in a dry cool, well-ventilated area, as cool as possible out of direct sunlight and away from children, animals, food, feedstuffs, seed and fertilisers. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.



**SAFETY DIRECTIONS**

Avoid contact with eyes and skin. Do not inhale spray mist. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26)

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**NOTICE TO BUYER**

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Macspred or Seller is hereby excluded. Macspred hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.



Crop C  
trader  
mark C

Shirqu

Spray:  
marks

Macsp  
ACN 0

No. 13

DELAC

TEL: 0:

APVMA



s after

Crop Care Australia Pty. Ltd. is a licensed user of Gladiator which is a registered trademark of Nufarm Technologies USA Pty Ltd. Hammer is a registered trade mark of FMC (Chemicals) Pty Ltd.

Shirquat and Spark are registered Trade Marks of Crop Care Australia.

Spray.Seed, Fusion, Gramoxone, Reglone and Fusilade are registered trade marks of Syngenta Limited.

Macspred Pty Ltd

ACN 011 029 495

No. 13 KENNEDYS DRIVE

DELACOMBE VIC 3356

TEL: 03 53358522

APVMA Approval No. 61396/0508

tory or  
against  
buyer  
ect or  
advice  
t such  
s been  
ces be  
e paid

