

TEMPLATE FOR RELEVANT LABEL PARTICULARS (RLPs) (Veterinary Products)

Select appropriate:

- New Product (include all applicable RLPs) OR
 Variation (highlight instructions that are being varied). Approval no. of label being varied:
61663/0509

Signal heading:	Read Safety Directions Before Opening or Using For Animal Treatment Only
Product name:	WSD Cy-Guard Spray On Sheep Blowfly Protection
Active constituent/s:	60 g/L Cyromazine
Statement of claims:	When applied as directed will protect long wool sheep against fly strike for 11 weeks.
Net contents:	1L, 5L, 20L
Directions for Use Heading:	Directions for use
Restrains:	Do not use on sheep which are producing or may in the future produce milk or milk products for human consumption.
Contraindications:	This product should not be used on sheep with less than 6 weeks wool as a reduced period of protection may result. This product should not be used on open wounds e.g. those resulting from marking or mulesing operations. This product should not applied during gusty or windy weather as spray drift will reduce the period of protection.
Precautions:	--
Side effects:	-



Dosage & administration:	<table border="1"> <thead> <tr> <th>MONTHS SINCE SHEARING</th> <th>TOTAL VOLUME / SHEEP (ML)</th> <th>VOLUME PER TREATMENT BAND (ML)</th> <th>NO. OF BANDS</th> <th>SHEEP TREATED PER 1L</th> <th rowspan="15">CRITICAL COMMENTS 1. Use a WSD recommended Spray-On applicator gun. 2.Ensure the applicator gun is clean. 3.Calibrate applicator gun using an accurate volume measure. 4.Ensure all areas to be protected are treated. 5.When preventing body strike ensure bands overlap along the midline. 6.This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. 7.Heavy rain following the application of this product could diminish the period of protection. 8.WSD CY-GUARD SPRAY is colored for band identification. This scourable dye will disappear shortly after treatment. SEE DIAGRAM</th> </tr> </thead> <tbody> <tr> <td colspan="5">FOR THE PREVENTION OF BODY STRIKE</td> </tr> <tr> <td>Up to 6</td> <td>34</td> <td>17</td> <td>2</td> <td>29</td> </tr> <tr> <td>7</td> <td>38</td> <td>19</td> <td>2</td> <td>26</td> </tr> <tr> <td>8</td> <td>44</td> <td>22</td> <td>2</td> <td>22</td> </tr> <tr> <td>9</td> <td>50</td> <td>25</td> <td>2</td> <td>20</td> </tr> <tr> <td>10</td> <td>56</td> <td>28</td> <td>2</td> <td>17</td> </tr> <tr> <td>Lambs less than 15 kg</td> <td>17</td> <td>17</td> <td>1</td> <td>58</td> </tr> <tr> <td colspan="5">FOR THE PREVENTION OF BODY AND CRUTCH STRIKE</td> </tr> <tr> <td>Up to 6</td> <td>51</td> <td>17</td> <td>3</td> <td>19</td> </tr> <tr> <td>7</td> <td>57</td> <td>19</td> <td>3</td> <td>17</td> </tr> <tr> <td>8</td> <td>66</td> <td>22</td> <td>3</td> <td>15</td> </tr> <tr> <td>9</td> <td>75</td> <td>25</td> <td>3</td> <td>13</td> </tr> <tr> <td>10</td> <td>84</td> <td>28</td> <td>3</td> <td>11</td> </tr> <tr> <td>Lambs less than 15 kg</td> <td>34</td> <td>17</td> <td>2</td> <td>29</td> </tr> </tbody> </table>	MONTHS SINCE SHEARING	TOTAL VOLUME / SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS	SHEEP TREATED PER 1L	CRITICAL COMMENTS 1. Use a WSD recommended Spray-On applicator gun. 2.Ensure the applicator gun is clean. 3.Calibrate applicator gun using an accurate volume measure. 4.Ensure all areas to be protected are treated. 5.When preventing body strike ensure bands overlap along the midline. 6.This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. 7.Heavy rain following the application of this product could diminish the period of protection. 8.WSD CY-GUARD SPRAY is colored for band identification. This scourable dye will disappear shortly after treatment. SEE DIAGRAM	FOR THE PREVENTION OF BODY STRIKE					Up to 6	34	17	2	29	7	38	19	2	26	8	44	22	2	22	9	50	25	2	20	10	56	28	2	17	Lambs less than 15 kg	17	17	1	58	FOR THE PREVENTION OF BODY AND CRUTCH STRIKE					Up to 6	51	17	3	19	7	57	19	3	17	8	66	22	3	15	9	75	25	3	13	10	84	28	3	11	Lambs less than 15 kg	34	17	2	29
	MONTHS SINCE SHEARING	TOTAL VOLUME / SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS	SHEEP TREATED PER 1L	CRITICAL COMMENTS 1. Use a WSD recommended Spray-On applicator gun. 2.Ensure the applicator gun is clean. 3.Calibrate applicator gun using an accurate volume measure. 4.Ensure all areas to be protected are treated. 5.When preventing body strike ensure bands overlap along the midline. 6.This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. 7.Heavy rain following the application of this product could diminish the period of protection. 8.WSD CY-GUARD SPRAY is colored for band identification. This scourable dye will disappear shortly after treatment. SEE DIAGRAM																																																																							
	FOR THE PREVENTION OF BODY STRIKE																																																																												
	Up to 6	34	17	2	29																																																																								
	7	38	19	2	26																																																																								
	8	44	22	2	22																																																																								
	9	50	25	2	20																																																																								
	10	56	28	2	17																																																																								
	Lambs less than 15 kg	17	17	1	58																																																																								
	FOR THE PREVENTION OF BODY AND CRUTCH STRIKE																																																																												
	Up to 6	51	17	3	19																																																																								
	7	57	19	3	17																																																																								
	8	66	22	3	15																																																																								
	9	75	25	3	13																																																																								
	10	84	28	3	11																																																																								
Lambs less than 15 kg	34	17	2	29																																																																									
FOR THE PREVENTION OF POLL STRIKE Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of WSD CY-GUARD SPRAY to the areas requiring protection, around the poll and horns.																																																																													
Not to be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.																																																																													
General directions:	Do not dilute. Do not mix with any other product. Apply directly from the container using WSD recommended spray-on applicator gun. Except for small lambs under 15Kg, the dose volume is determined by the number of months shearing. Apply WSD Cy-Guard according to the diagrams and tables in the Directions for Use. WSD Cy-Guard Spray may be applied to all breeds of sheep, providing protection against flystrike. WSD Cy-Guard Spray is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25cm above the fleece, to achieve a 15cm wide band at each pass.																																																																												
Withholding Period/s:	Withholding periods: Meat: Do not use less than 7 days before slaughter for human consumption. Milk: DO not use on sheep which are producing or may in the future produce milk or milk products for human consumption. WOOL HARVEST INTERVAL: DO NOT use less than 2 months before shearing or fibre collection.																																																																												
Trade Advice:	Export Slaughter Interval (ESI): Do not use less than 28 days before slaughter for export. Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.																																																																												
Safety Directions:	Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.																																																																												
First Aid:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.																																																																												

Additional user safety:	Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling. Material Safety Data Sheet: If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone WSD (08) 9321 2888 or visit www.westernstock.com.au
Environmental statements:	Do not contaminate dams, rivers or streams with the product or used containers.
Disposal:	Triple or preferably pressure rinse containers and dispose of rinsate in a disposal pit as described below. Do not dispose of undiluted chemicals on site. Break crush or puncture and bury empty container in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and setup for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned.
Storage:	Store below 30C (room temperature). Store in original container tightly closed in a dry, cool place. Do not store in direct sunlight. Do not re-use the container. Keep out of reach of children.
Name & address:	Western Stock Distributors ABN 58 009 372 525, Koojan Avenue South Guildford WA 6055 Telephone (08) 9321 2888 Email sales@westernstock.com.au Website www.westernstock.com.au

The following is for APVMA use only:

APVMA approval no.	APVMA No: 61663/0614
--------------------	-----------------------------