

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# EMAZ

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present as the ammonium salt

**GROUP B HERBICIDE**

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.**

**eChem (Australia) Pty Ltd**

(ABN 23 089 133 095)  
13833 New England Highway,  
CAMBOOYA QLD 4358  
Ph: (07) 4696 1054 Fax: (07) 4696 1057



1  
Litre



## STORAGE AND DISPOSAL

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

## SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank-mixing with STOMP® or Paraquat, consult also the safety directions for those products.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.



## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier.

## CONDITIONS OF SALE:

eChem (Australia) Pty. Ltd. accepts responsibility for the consistent quality of the product however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

APVMA Approval Number: 61822/1L/1009

Batch Number:

Date of Manufacture:

Barcode

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# EMAZ

## HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present as the ammonium salt

**GROUP B HERBICIDE**

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.**

**eChem (Australia) Pty Ltd**

(ABN 23 089 133 095)  
13833 New England Highway,  
CAMBOOYA QLD 4358  
Ph: (07) 4696 1054 Fax: (07) 4696 1057



**5  
Litres**



Barcode

APVMA Approval Number: 61822/5L/1009

Batch Number:

Date of Manufacture:



### STORAGE AND DISPOSAL

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

### SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank-mixing with STOMP® or Paraquat, consult also the safety directions for those products.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier.

### CONDITIONS OF SALE:

eChem (Australia) Pty. Ltd. accepts responsibility for the consistent quality of the product however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# EMAZ

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present as the ammonium salt

**GROUP B HERBICIDE**

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.**

**eChem (Australia) Pty Ltd**  
(ABN 23 089 133 095)  
13833 New England Highway,  
CAMBOOYA QLD 4358  
Ph: (07) 4696 1054 Fax: (07) 4696 1057



**10  
Litres**

**STORAGE AND DISPOSAL**

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank-mixing with STOMP® or Paraquat, consult also the safety directions for those products.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier.

**CONDITIONS OF SALE:**

eChem (Australia) Pty. Ltd. accepts responsibility for the consistent quality of the product however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

APVMA Approval Number: 61 822/10L/1009

Batch Number:

Date of Manufacture:



**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# EMAZ

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present as the ammonium salt

**GROUP B HERBICIDE**

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.**

**eChem (Australia) Pty Ltd**  
(ABN 23 089 133 095)  
13833 New England Highway,  
CAMBOOYA QLD 4358  
Ph: (07) 4696 1054 Fax: (07) 4696 1057



**20  
Litres**

**STORAGE AND DISPOSAL**

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank-mixing with STOMP® or Paraquat, consult also the safety directions for those products.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier.

**CONDITIONS OF SALE:**

eChem (Australia) Pty. Ltd. accepts responsibility for the consistent quality of the product however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

APVMA Approval Number: 61 822/20L/1009

Batch Number:

Date of Manufacture:



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# EMAZ

HERBICIDE  
ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present  
as the ammonium salt

GROUP **B** HERBICIDE

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use.

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

APVMA Approval Number: 61822/1009

**eChem (Australia) Pty Ltd**

(ABN 23 089 133 095)  
13833 New England Highway,  
CAMBOOYA QLD 4358  
Ph: (07) 4696 1054 Fax: (07) 4696 1057

## DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT apply to soils saturated with water.  
DO NOT apply by aircraft.

| SITUATION  | WEEDS CONTROLLED   | RATE/ha  | CRITICAL COMMENTS   |
|--|--|--|---|
| Fallow prior to planting winter crops as directed in the follow crops table. (Qld and Nthn NSW only) | Barnyard grass ( <i>Echinochloa colona</i> ), Liverseed grass ( <i>Urochloa panicoides</i> ), Stink grass ( <i>Eragrostis cilianensis</i> ), Blowaway grass ( <i>Panicum decompositum</i> ), Button grass ( <i>Dactyloctenium radulans</i> ), Pigweed ( <i>Portulaca oleracea</i> ), Yellow vine ( <i>Tribulus terrestris</i> ), Mintweed ( <i>Salvia reflexa</i> ), Boggabri weed ( <i>Amaranthus mitchellii</i> ), Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ), Peachvine/cowvine ( <i>Ipomoea lonchophylla</i> )   | 150 to 200 mL  | Treatments may be applied from the end of the previous crop harvest but not less than 3 months prior to planting of winter crop (refer to follow crops table). Best results are obtained where treatments are applied to dry, weed-free soil prior to weed germination. Treatments can be applied to standing crop residues. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Emerged weeds must be controlled by prior cultivation, or by application of a knockdown herbicide. Where infestation of emerged weeds is light, tank mix treatment with glyphosate or paraquat. Cultivation following application of treatments may reduce pre-emergence weed control provided by Emaz Herbicide. Prolonged wet soil conditions following treatment will reduce the effective life of pre-emergence treatments. Weed escapes may require follow up application of knockdown herbicides. DO NOT apply more than 200 mL/ha per season on cropland. DO NOT apply in more than two successive seasons on cropland. DO NOT overspray headlands.   |
| Sugarcane (Qld, NSW and WA Only)   | Awnless barnyard grass ( <i>Echinochloa colona</i> ), Barnyard grass ( <i>E. crus-galli</i> ), Blackberry nightshade ( <i>Solanum nigrum</i> ), Blue billygoat weed ( <i>Ageratum houstonianum</i> ), Common sida ( <i>Sida rhombifolia</i> ), Giant/Black pigweed ( <i>Triathema portulacastrum</i> ), Glossy nightshade ( <i>Solanum americanum</i> ), Green amaranth ( <i>Amaranthus viridis</i> ), Green summer grass ( <i>Brachiaria subquadriflora</i> ), <i>Ipomoea</i> spp., Guinea grass ( <i>Panicum maximum</i> ), Liverseed grass ( <i>Urochloa panicoides</i> ), Milkweed (Mexican fire plant) ( <i>Euphorbia heterophylla</i> ), Pigweed ( <i>Portulaca oleracea</i> ), Summer grass ( <i>Digitaria ciliaris</i> ) | 300 to 400 mL  | DO NOT apply more than once a year to the same crop. Pre-emergence application:<br>PLANT CANE: Apply after planting and before emergence of the cane. On sandy soil, ensure thorough coverage of sets to a minimum depth of 5 cm.<br>RATOON: Apply after harvest and before new cane growth appears. Use the higher rates of Emaz Herbicide for increased residual control of weeds or anticipated high weed pressure. Heavy trash layers may prevent the product from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Emerged weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Weed control can be improved under dry conditions or where incorporating moisture has been insufficient, by cultivating the treated soil above sets. Care needs to be taken to minimise exposure of untreated soil or sets. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Heavy rain and/or irrigation within 2 days following application may concentrate Emaz Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Weed escapes may require follow up application of an appropriate herbicide. Contact eChem (Australia) Pty Ltd for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label). |
| Sugarcane (Qld, NSW and WA only)   | As above plus Crowsfoot grass ( <i>Eleusine indica</i> )   | 400 ML plus 3 L STOMP+ 330E or 2.2 L STOMP+ Xtra                                   | Incorporate into the soil within 7 days of application, either by overhead irrigation, rainfall or mechanical incorporation, to a minimum depth of 5 cm. Read all directions for use in sugarcane as well as the Restraints and Follow Crops restrictions on the STOMP 330E or STOMP Xtra label.  |
|  | As above plus control of those weeds expected with these rates of 250 g/L paraquat products alone.   | 300 to 400 mL plus 1.2 to 1.6 L of 250 g/L paraquat product                        | Post-emergence application (in emerged cane): ALWAYS tank mix Emaz Herbicide with registered and appropriate rates of paraquat for the emerged weeds. Use the higher rates of Emaz Herbicide for increased residual control of weeds or anticipated high weed pressure. Use the higher rates of paraquat for dense, more mature weeds. Apply as a carefully directed inter-row spray to weeds and at the base of the plant, minimising contact with sugarcane foliage. Contact with sugarcane foliage may cause temporary crop damage observed as inter veinal yellowing and reduction in crop biomass or crop height. Refer to the Paraquat label re spray boom types and precautions. Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis. DO NOT cultivate within 1 hour of application. Heavy rain and/or irrigation within 2 days following application may concentrate Emaz Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. DO NOT spray plants covered with heavy dew. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. <b>Contact eChem (Australia) Pty Ltd for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label).</b>  |
| Peanuts (Qld, NSW and NT Only)   | Awnless barnyard grass ( <i>Echinochloa colona</i> ), Barnyard grass ( <i>E. crus-galli</i> ), Blackberry nightshade ( <i>Solanum nigrum</i> ), Blue billygoat weed ( <i>Ageratum houstonianum</i> ), Common sida ( <i>Sida rhombifolia</i> ), Giant/Black pigweed ( <i>Triathema portulacastrum</i> ), Glossy nightshade ( <i>Solanum americanum</i> ), Green amaranth ( <i>Amaranthus viridis</i> ), Green summer grass ( <i>Brachiaria subquadriflora</i> ), <i>Ipomoea</i> spp., Guinea grass ( <i>Panicum maximum</i> ), Liverseed grass ( <i>Urochloa panicoides</i> ), Milkweed (Mexican fire plant) ( <i>Euphorbia heterophylla</i> ), Pigweed ( <i>Portulaca oleracea</i> ), Summer grass ( <i>Digitaria ciliaris</i> ) | 300 to 400 mL  | Pre-emergence application. Optimum weed control will occur when moisture is maintained in the top 5 cm of soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. DO NOT apply to any one crop more than once per season.  |
|  | As above plus Nutgrass* ( <i>Cyperus rotundus</i> )  | 400 mL plus Hasten+ or Kwicken+ at 1 L/100L (or a similar esterified oil adjuvant) | post-emergence application. Apply prior to crop flowering. DO NOT apply to any one crop more than once per season.<br>* Nutgrass: suppression only when applied to nutgrass at the 3 to 4 leaf stage.   |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS:

**HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.**

### GRAZING/STOCKFEED WHP:

**Wheat, Barley, Chickpea, Peanuts: DO NOT GRAZE TREATED VEGETATION, STUBBLE OR WEEDS, OR CUT FOR STOCKFOOD, FOR 4 WEEKS AFTER APPLICATION.**

**Sugarcane: DO NOT GRAZE CROP OR SPRAYED VEGETATION, OR CUT FOR STOCKFOOD, FOR 6 WEEKS AFTER APPLICATION.**

### GENERAL INSTRUCTIONS

#### All Applications

Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis.

#### Fallow application:

Emaz Herbicide can be applied in fallow prior to the planting of various crops as outlined in the Follow Crops table. Emaz Herbicide is a pre-emergence herbicide, which is recommended to provide short term (6 to 12 weeks) pre-emergence control of summer germinating grasses and broadleaved weeds in cultivated or uncultivated fallow. A green weed cover of greater than 50% will significantly reduce the length of pre-emergence weed control.

#### Sugarcane application

##### Pre-emergence

Where possible, apply to soil two days prior to incorporation. Incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm. Incorporation with flood irrigation may result in loss of weed control. Emaz Herbicide should be applied to plant or ratoon cane before the cane has emerged. When applying Emaz Herbicide in moist conditions, allow weeds to germinate prior to application and apply mixed with an appropriate knockdown herbicide. Whenever emerged weeds are present, Emaz Herbicide should be mixed with an appropriate knockdown herbicide.

##### Post-emergence

Emaz Herbicide MUST always be mixed with registered rates of paraquat and applied as a carefully directed inter-row spray to weeds and at the base of the sugarcane stalks, minimising spray contact with sugarcane foliage. DO NOT add crop oil to this mixture. Other surfactant or adjuvant is not required.

#### Peanut application

##### Pre-emergence:

Apply to soil and incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm.

##### Post-emergence:

Emaz Herbicide must always be mixed with an adjuvant as directed in the DIRECTIONS FOR USE section of the label.

##### Soil Type:

Pre-emergence weed control with Emaz Herbicide can be limited when applied onto soils with pH <5.0 and/or which contain high concentrations of iron (Fe) and/or aluminium (Al). These situations are common in soil types such as Krasnozems and red brown Ferrosols.

##### MIXING

Emaz Herbicide is an aqueous solution. Part fill the spray tank with water, then with agitator running, add the required amount of product, then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product. For post-emergence applications in peanuts, add adjuvant after the addition of Emaz Herbicide to the tank.

No adjuvants or surfactants are required when Emaz Herbicide is used alone for pre-emergence weed control or with paraquat. DO NOT add crop oil when applied post-emergence with paraquat.

##### APPLICATION

DO NOT apply by aircraft.

##### Fallow application:

Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

##### Sugarcane:

Ensure thorough coverage of soil and weeds with well-calibrated spray equipment. Apply in a minimum of 200 L water/ha using a nozzle configuration to generate a medium to coarse spray pattern for pre-emergence applications or medium spray pattern for post-emergence applications. Replace worn or old nozzles and calibrate equipment on a regular basis.

**Peanuts:**

Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands. For post-emergence applications, Emaz Herbicide should be applied a minimum of 2 hours before rainfall.

**EQUIPMENT CLEAN-UP**

Thoroughly flush all spray equipment with water following the use of Emaz Herbicide and before use with other products.

**COMPATIBILITY**

Emaz Herbicide is compatible with STOMP<sup>+</sup>, atrazine, glyphosate, paraquat, paraquat + diquat, Ally<sup>+</sup>, Garlon<sup>+</sup>, Touchdown<sup>+</sup> and 2,4-D.

It is NOT compatible with dicamba.

**RESISTANT WEEDS WARNING****GROUP B HERBICIDE**

Emaz Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group B herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population, if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, eChem (Australia) Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

**FOLLOW CROPS**

Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

**Following use in summer fallow:-**

| MONTHS AFTER APPLICATION   |  |   |                             |                 |
|--|--|---|-----------------------------|-----------------|
| 3  | 4  | 10  | 24                          | 36              |
| Faba beans, peanuts, mungbeans, maize varieties with CLEARFIELD Technology ONLY:<br>-Pacific Hycorn 621T<br>-Pacific Hycorn 531T<br>-Pioneer 33951R;<br>wheat varieties with CLEARFIELD Technology only. | Wheat <sup>①</sup> , (except for wheat varieties with CLEARFIELD Technology ONLY), barley <sup>①</sup> , lucerne <sup>①</sup> , chickpea | Sorghum <sup>②</sup> , maize <sup>②</sup> (except for maize varieties with CLEARFIELD Technology ONLY), dryland cotton <sup>③</sup> | Irrigated cotton, sunflower | Any other crops |

① The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), barley or lucerne during the next winter season.

DO NOT apply Emaz Herbicide in fallow later than the end of December.

DO NOT apply Emaz Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

② The following additional requirements apply if it is intended to plant sorghum or maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply Emaz Herbicide in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

③ The following additional requirements apply if it is intended to plant dryland cotton during the next summer season.

DO NOT apply Emaz Herbicide in areas where rainfall from spraying to sowing of dryland cotton is expected to be below 550 mm.

**Following use in sugarcane and peanuts:-**

| MONTHS AFTER APPLICATION   |  |                         |  |                      |                             |                |
|--|--|-------------------------|--|----------------------|-----------------------------|----------------|
| 0  | 4  | 5                       | 10   | 18                   | 24                          | 36             |
| Mung beans, Peanuts, soybeans, maize varieties with CLEARFIELD Technology ONLY:<br>-Pacific Hycorn 621T<br>-Pacific Hycorn 531T<br>-Pioneer 33951R | Chick peas, lucerne <sup>①</sup> , barley <sup>①</sup> , wheat <sup>①</sup> (except for wheat varieties with CLEARFIELD Technology ONLY) | Lupins, pasture legumes | Maize <sup>②</sup> , (except varieties with CLEARFIELD Technology ONLY), dryland cotton <sup>③</sup> | Sorghum <sup>②</sup> | Irrigated cotton, sunflower | Any other crop |

① The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), barley or lucerne during the next winter season.

DO NOT apply Emaz Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

② The following additional requirements apply if it is intended to plant sorghum or maize (except for varieties with CLEARFIELD Technology), during the next summer season.

DO NOT apply Emaz Herbicide in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

③ The following additional requirements apply if it is intended to plant dryland cotton during the next summer season.

DO NOT apply Emaz Herbicide in areas where rainfall from spraying to sowing of dryland cotton is expected to be below 550 mm.

**RE-ENTRY PERIOD FOR Emaz Herbicide PLUS STOMP Xtra**

When tank-mixing Emaz Herbicide with STOMP Xtra, DO NOT allow entry into treated areas for 12 hours. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50 m of wetlands or waterways.

**CROP SAFETY**

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth in sugarcane but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under conditions where soil moisture is high within 2 days following application and on light/sandy soil. Contact eChem (Australia) Pty Ltd for advice on variety susceptibility before using this product.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank-mixing with STOMP<sup>+</sup> or Paraquat, consult also the safety directions for those products.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**CONDITIONS OF SALE:**

eChem (Australia) Pty. Ltd. accepts responsibility for the consistent quality of the product however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

\* Other trademarks

