



**POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**CONQUEST**

# HATCHET

**HERBICIDE**

ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester.  
100 g/L PICLORAM present as hexyloxypropylamine salt.

**GROUP | HERBICIDE**

For the control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.  
SHAKE WELL BEFORE USE.**



Conquest Agrochemicals Pty Ltd.  
ABN 84 098 814 932  
Unit 1, 46 Hasler Road  
Osborne Park WA 6017  
Telephone (08) 9347 0500, Facsimile: (08) 9347 0551

**CONTENTS**

 **5L**  
 **20L**

**GENERAL INSTRUCTIONS**

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near feedstuffs, fertilizers or seed. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

**CONDITIONS OF SALE**

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL  
**1 800 033 111**  
24 hours Australia wide.

Batch No.

DOM

APVMA Approval No: 61863/5U/0507





RECEIVED  
By Leslie Roberts at 2:12 pm, Jan 11, 2008

APVMA Approval No: 61893/5L/0507  
Application Number: 41728  
Product Name: Conquest Hatched Herbicide  
Pack Size: 5L Label  
Dimensions: 145mm x 150mm  
Print Size: A4 - 100%

**POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**CONQUEST**

# HATCHET

**HERBICIDE**

ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester.  
100 g/L PICLORAM present as hexyloxypropylamine salt.

**GROUP | HERBICIDE**

For the control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.  
SHAKE WELL BEFORE USE.**



Conquest Agrochemicals Pty Ltd.  
ABN 84 098 814 932  
Unit 1, 46 Hasler Road  
Osborne Park WA 6017  
Telephone (08) 9347 0500, Facsimile: (08) 9347 0551

**CONTENTS**

■ 5L  
■ 20L

**GENERAL INSTRUCTIONS**

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near feedstuffs, fertilizers or seed. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

**CONDITIONS OF SALE**

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL  
**1 800 033 111**  
24 hours Australia wide.

Batch No.

DOM

APVMA Approval No: 61863/20/0507





APVMA Approval No: 61893/20L/0507  
Application Number: 41728  
Product Name: Conquest Hatched Herbicide  
Pack Size: 20L Label  
Dimensions: 145mm x 150mm  
Print Size: A4 - 100%

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**CONQUEST**  
**HATCHET**  
HERBICIDE


ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester.  
100 g/L PICLORAM present as hexyloxypropylamine salt.

**GROUP I HERBICIDE**

For the control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use Table.

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING**

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL  
**1800 033 111**  
ALL HOURS AUSTRALIA WIDE



**Conquest Agrochemicals Pty. Ltd**  
ABN 84 098 814 932  
Unit 1, 46 Hasler Road  
Osborne Park, WA, 6017

Telephone (08) 9347 0500 Facsimile (08) 9347 0551

APVMA Approval Number : 61863/0507



Conquest Agrochemicals Pty Ltd.  
ABN 84 098 814 932  
Unit 1, 46 Hasler Road  
Osborne Park, WA, 6017  
Telephone (08) 9347 0500, Facsimile: (08) 9347 0551

**DIRECTIONS FOR USE RESTRAINTS:**

- DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.
- DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.
- DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.
- DO NOT apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C.
- In areas prone to flooding, treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

**1. WOODY WEED SITUATIONS**

**Table A: High Volume Spraying**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEEDS GROWTH STAGE	STATE	RATE/100L WATER	CRITICAL COMMENTS
African boxthorn	Less than 2m tall	All States	500 mL	Apply when bushes have good leaf cover, growth and <b>no leaf fall</b> .
<i>Angophora</i> spp.	1 to 3m tall		350 mL	
Australian blackthorn	Less than 2m tall		500 mL	Apply from late spring to early autumn.
<i>Banksia</i> spp.	1 to 3m tall		350 mL	
Biddy bush (Chinese shrub), (Sifton bush)	Autumn when actively growing	ACT & NSW only	500 mL	Add Conquest Wetter 1000 at 125mL/100 L of water for best results.
Blackberry in association with: Docks, Ragwort, Smartweed Thistles	Late spring to autumn	All States except NT	350 mL or 500 mL	<b>Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry.</b> Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Conquest Hatchet herbicide.
Blue heliotrope	Flowering	NSW & Qld only	500 mL	Apply in a minimum spray volume of 1250L/ha
Brooms: Cape, English Flax leaf, Montpellier	Spring to mid- summer prior to pod formation	All States except NT	250 mL	Apply as a thorough foliage spray
	Autumn to winter		350 mL	
Camphor laurel	Less than 2m tall	All States		
	Above 2m tall		500 mL	
<i>Casuarina</i> spp.	1 to 3m tall		350 mL	
Chinese apple	Less than 2m tall	Qld & WA only	350 mL	Add Conquest Wetter 1000 at 100mL/100 L of water for best results.
Cockspur thorn	Spring to Autumn	ACT, NSW & Qld only		Apply as thorough foliage spray
Common sensitive plant	Any time when actively growing	NT, Qld & WA only	200 mL	To avoid leaves closing during application, spray plants while moving forward. Add Conquest Wetter 1000 at 100mL/100 L of water for best results.
Crofton weed	Spring to autumn	ACT, NSW & Qld only	350 mL	Apply as thorough foliage spray.
Eastern cotton bush ( <i>Maireana microphylla</i> )	Spring to autumn	NSW & Qld only	500 mL	Add Uptake* Spraying Oil at 500mL/100 L of water. Some bushes may require a follow up spray to control regrowth.
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3m tall	All States	350 mL or 500 mL	<b>Apply the high rate where difficult to control species of Eucalyptus regrowth is present.</b> Addition of an adjuvant may improve results- contact Conquest Agrochemicals Pty Ltd for details.
Galenia	Fresh growth in spring to summer	NSW only	500 mL	Use 200L of spray mixture/ha
Giant bramble	Spring to autumn	NT, Qld & WA only		Penetration of thick clumps may be difficult and respraying may be necessary. Add Conquest Wetter 1000 at 100mL/100 L of water for best results.
Gorse	1 to 1.5m tall	All States except NT	250 mL	Spring and Summer treatment only. Add Conquest Wetter 1000 at 100 mL/100 L of water for best results.
	Over 1.5m tall or autumn treatment		350 mL	Add Conquest Wetter 1000 at 100 mL/100 L of water for best results.
	Winter treatment		500 mL	Brownout may not be complete until summer. Add Conquest Wetter 1000 at 100mL/100 L of water for best results.
Groundsel bush ( <i>Baccharis halimifolia</i> )	1 - 1.5m tall in spring to summer	All States except NT	250 mL	Apply as a thorough foliar spray.
	Over 1.5 m tall or autumn treatment		350 mL	
Green cestrum	Late spring to early autumn	ACT, NSW & Qld only	500 mL	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
Hawthorn	Less than 2m tall	All States		Apply from late spring to early autumn.
Horehound	Pre- flowering		350 mL	Apply as a thorough foliar spray.
Japanese sunflower		NSW & Qld only		
Lantana ( <i>Lantana camara</i> ) ( <i>Lantana montevidensis</i> )	Up to 1m tall in summer to autumn 1 to 2m tall in summer to autumn	All States	500 or 750 mL	Add one of the following adjuvants, when using 350mL rate: Uptake* Spraying oil @ 0.5% v/v, Pulse*. Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
Lion Tail ( <i>Leonotis nepetifolia</i> )	Pre-flowering	Qld only	200 mL	Apply as a thorough foliar spray. Add Conquest Wetter 1000 at 100mL/100L of water for best results.
Limebush	Any time of the year with good leaf cover and soil moisture	NT & Qld only	350 mL	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results – contact Conquest Agrochemicals Pty Ltd for details.
Manuka	At flowering	Vic only	500 mL	For optimum results, add Pulse* Penetrant at 200mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite ( <i>Prosopis</i> spp.) ( <i>Prosopis velutina</i> )	Seedling, full leaf and flowering before podding	NSW, NT, Qld & WA only	350 mL	<b>DO NOT</b> spray plants bearing pods. Add Conquest Wetter 1000 at 100 mL/100L of water for best results.
		Qld only	670 mL	
Mistflower	Spring to autumn	ACT, NSW & Qld only		Apply as a thorough foliar spray.
Mother-of-millions	Flowering	NSW & Qld only	500 mL	Add Conquest Wetter 1000 at 100mL/100L of water for best results.
Paddy's lucerne	Active growth	NSW only		Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2m tall	NT, Qld & WA only	350 mL	Add Uptake* Oil at 500mL/100L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Prickly pear (common) Smooth tree pear	Active phyllode growth	All States	500 mL	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.
Rubber vine (Not infected with rust)	Up to 1.5m tall at flowering	NT, Qld & WA only	350 mL	Spray all leaves and stems just to the point of run off and thoroughly spray the base of the plant. With larger, more dense stands regrowth may appear
	Dense stands greater than 1.5m tall at flowering		500 mL	Subsequent control of any regrowth should be done by basal bark spraying.
Siam weed	Active growth	Qld & WA only	350 mL	Add Conquest Wetter 1000 at 100mL/100 L of water for best results.
Sicklepod	Up to flowering	NT, Qld & WA only	200 mL	<b>DO NOT</b> apply to podding plants. Add Conquest Wetter 1000 at 100 mL/100L of water for best results.
St. John's wort	From flowering to early seed set	All States	500 mL	Late spring to early summer.
Sweet briar	Up to 1.5m tall	All States except NT	350 mL	Add Conquest Metsulfuron 600 at 10g/100L water to obtain more reliable results with the lower rate of Conquest Hatchet.
			500 mL	Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Tobacco weed	Actively growing plants	NT, Qld & WA only	300 mL	Add Conquest Wetter 1000 at 100mL/100L of water for best results.
Wattle ( <i>Acacia</i> spp.) (except corkwood wattle)	1 to 3m tall	All States	350 mL	
Wild rosemary ( <i>Cassinia laevis</i> )	Active growth, 0.5 to 1.0m tall	Qld only	350 to 500 mL	Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.
Wild tobacco tree	Spring to autumn up to 2m tall	ACT, NSW & Qld only	350 mL	

**Table B: Aerial Application**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / ha	CRITICAL COMMENTS
Blackberry	Summer to autumn	NSW, Qld, SA, Vic & WA only	10L	Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Conquest Hatchet. <b>WARNING:</b> <i>Eucalyptus</i> species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 to 20 m tall may be partially defoliated but are likely to recover.
Gorse		Tas only		Helicopter application only
Cockspur thorn Crofton weed Lantana Mistflower	Late autumn	NSW, NT & Qld only (helicopter only)	1.5 L plus 6L Conquest Amine 625	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha. <b>Follow-up respraying will be required.</b>
Lantana			10L	Helicopter application only
Rubber vine (Not infected with rust)	When flowering	NT & Qld only (helicopter only)	3L to 5L	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands. However, complete coverage and penetration may be difficult. Follow-up respraying will be required. Any regrowth should be sprayed with a suitable basal bark herbicide.
St. John's wort	Flowering to early seed set (Nov-Jan)	NSW only	4L	Helicopter application only. Follow up spraying will be required in the following season.

AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / ha	CRITICAL COMMENTS
Parkinsonia	Seedlings 1 – 2m tall, or 12 - 24 months old	Qld and NT only (helicopter only)	3 L	Add Uptake* Oil at 1L/ha


**Table C: Controlled Droplet Application (CDA)**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / ha	CRITICAL COMMENTS
Blackberry in association with: Docks, Ragwort, St. John's wort, Thistles	Summer to autumn	All States except NT	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be re-sprayed after hardening off. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Conquest Hatchet.

**Table D: Low Volume High Concentrate Application Techniques (GAS POWERED GUN, SPRINKLER SPRAYER)**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / 10L WATER	CRITICAL COMMENTS
Blackberry	Late Spring to autumn	ACT, NSW, Qld, SA, TAS and WA only	335 mL	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Camphor laurel Cockspur thorn Crofton weed	Less than 1.5m high	ACT, NSW and Qld only	500 mL	
Eucalyptus species	Seedlings up to 2m tall	All States	335 mL	
Mistflower	Less than 1.5m high	ACT, NSW and Qld only	500 mL	
Sweet briar	1.5m tall, full leaf to ripe fruit	NSW only		<b>Gas powered Gun only:</b> Apply to actively growing bushes not more than 1.5m tall that have not more than 5 stems from the crown.
St. John's wort	During flowering to early seed set	NSW, Tas and Vic only		<b>Gas powered Gun only:</b> One application should provide control. Minor regrowth and seedlings may be re-treated the following summer.
Wild tobacco tree	Less than 1.5m high	ACT, NSW and Qld only		Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

**Table E: Boom Application**  
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.



AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Galenia	Fresh growth during spring to summer	NSW only	5L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	NT and Qld only	3L	DO NOT apply to podding plants. Add Conquest Wetter 1000 at 100 mL/100L of water.
St. John's wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4 L	Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

**2. FALLOW SITUATIONS**

**Table A: Boom Application**  
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blackberry nightshade - <b>Suppression only</b>	10 to 25 cm tall prior to flowering	NSW and Qld only	200 to 400 mL + 1.2L Conquest Glyphosate 450 + Conquest Wetter 1000	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing
Camel melon Prickly paddy melon Cucumber melon ( <i>Cucumis melo</i> )	From 2 leaf to 50 cm diameter			Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5cm diameter for <i>Polymeria pusilla</i> ).
Common sowthistle	From 8 leaf to flowering			Refer to Conquest Glyphosate 450 label for use of adjuvant
Cow vine	From 2 to 5 leaf up to 15 cm diameter, prior to flowering			<b>DO NOT</b> plant susceptible crops for up to nine months after application, as specified in General Instructions – Minimum Re-cropping Periods - Black Cracking Clay Soils, Northern NSW & Qld.
Lucerne (established)	Active growth, 15 to 25 cm high, during spring		300 to 500 mL + 1.2L Conquest Glyphosate 450 + Conquest Wetter 1000	Dry conditions after application will increase the re-cropping interval.
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20cm diameter, prior to flowering		200 to 400 mL + 1.2L Conquest Glyphosate 450 + Conquest Wetter 1000	

**Table B: Blanket Wiper Application**  
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Bitterbark ( <i>Alistonia constricta</i> )	From summer to end of autumn	Qld	1.4 (1 part Conquest Hatchet to 4 parts water)  2% solution for spot spray (eg. 100 mL Conquest Hatchet in 5L water)	For use with blanket wipers only. For best results apply in autumn to tall (>60 cm) plants using two opposite directional passes (up and back) Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks. Blanket wiper application can be made in summer when plants are smaller but follow up spot spraying may be necessary. Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months). <b>Spot spraying</b> "missed" plants: thoroughly wet all stems and leaves without producing any solution run off. Avoid any spray reaching the soil surface.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA FOR BLACKBERRY:**  
**DO NOT** treat bushes carrying mature or near mature fruit.

**FOR NATIVE VEGETATION:**  
Use of Conquest Hatchet on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

**WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.**

**GENERAL INSTRUCTIONS**

**RESISTANCE WEEDS WARNING**



Conquest Hatchet Herbicide is a member of the pyridines group of herbicides. Conquest Hatchet has the disruptors of plant cell growth mode of action. For weed resistance management, Conquest Hatchet is a Group I herbicide.  
Some naturally-occurring weed biotypes resistant to Conquest Hatchet and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Hatchet or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Agrochemicals Pty Ltd accepts no liability for any losses that may result from the failure of Conquest Hatchet to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Conquest Agrochemicals Pty Ltd representative.

**MINIMUM RECROPPING PERIODS- Black Cracking Clay Soils, Northern NSW & Qld.**

**Table A: Boom Application**

RATE mL/ha	Plant-back periods for crops following the application of Conquest Hatchet for rates up to 600 mL/ha		
	200	300	600
CROP	MONTHS		
Wheat	2	2	4
Barley	2	2	4
Canola	2	4	4
Faba Beans	4	4	6
Chickpeas	4	6	6
Lucerne	6	9	9

These plantback periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plantback period may be significantly longer.

**Table G: Blanket Wiper Application**

Plant-back periods for crops following blanket wiper application.		
CROP	MONTHS	
Broadleaf crop	18	
Lucerne	6	
	This will allow any potential soil residues to dissipate, if any, and allow effective control.	

**NOTE: Before using Conquest Hatchet in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product with the longest plant-back period, will determine the time between spraying and planting.**

**COMPATIBILITY**

Follow any regional restrictions, and all directions and restrictions on the label, for any chemical mixed with Conquest Hatchet (e.g 2,4-D amine).  
Conquest Hatchet is compatible with the following herbicides: Amicide\*625, Lantana\* DP600, Brushkiller\*/Associate\*, Nufarm Glyphosate CT\*, Roundup\*CT, Credit\* plus Bonus\* and Roundup PowerMAX\*.  
Conquest Hatchet is compatible with the following adjuvants, as per Directions for Use: Conquest Wetter 1000, Pulse\*Penetrant, LI 700\* and crop oils.

**MIXING**

Mix only with water.  
Half fill the spray unit with water, and add the required amount of Conquest Hatchet. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

**APPLICATION**

**1. WOODY WEED SITUATIONS**  
Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

**A. High Volume Spraying**

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

**Hand Gun**

Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa (400 to 500 kPa for St John's wort). A spray volume of 3000 to 4000 L per infested hectare of 1 to 2 metre high blackberry (30 to 40 L/100m2) should be used. Use 2000 L of spray mixture/ha of galenia infestation (i.e. 20L/100 m2 infested area).

**Knapsack**

Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun.  
A spray volume of 3 to 4 L/10m2 infested area should be used.  
A spray volume of 2L/10m2 should be used for an area infested with galenia.

**B. Aerial Application**

Apply in 200 L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter. The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets. **DO NOT** spray when wind exceeds 15 km/hr and/or air temperature reaches 35°C.

**C. Controlled Droplet Application (CDA)**

Results similar to high volume spraying can be obtained using Micron Herbi\* or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm2. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with CDA unit.

**D. Low Volume High Concentrate Application Techniques**

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

**Gas powered gun**

Apply 50mL shots to obtain uniform coverage of 4 to 5 m2 of surface area of bush. This relates to 20 droplets/cm2 of leaf surface.

**Sprinkler sprayer**

This technique involves using a micro sprinkler that is connected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

**E. Boom Application**

Application in a minimum spray volume of 200 L/ha for galenia and St. John's wort and 600 L of water/ha for sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200 - 300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

**2. FALLOW SITUATIONS**

**F. Boom Application**

Application of Conquest Hatchet in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200 - 300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

**Table G. Blanket Wiper Application**

Blanket needs to be made from durable and wettable material with a rigid backing.  
Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (re-wetting). Ideally, a scraper bar should be mounted at the front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10 - 15 kph are ideal for blanket wiping application.

**CLEANING SPRAY EQUIPMENT**

**Rinsing**

After using Conquest Hatchet empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsings procedure twice.

**Decontamination**

Before spraying cotton and other sensitive crops with equipment that has been used to apply Conquest Hatchet, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent such as Spraymate Tank and Equipment Cleaner at 500 mL/100 L of water and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Crops susceptible to Conquest Hatchet include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and Pinus spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines. Conquest Hatchet is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram, one of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

**DO NOT** allow physical spray drift onto waterways, native vegetation and susceptible crops.

**DO NOT** apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants. **DO NOT** apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. **DO NOT** move soil, which may have been treated to areas where desirable plants are to be grown.

**PROTECTION OF LIVESTOCK**

Poisonous plants may become palatable after spraying and stock should be kept away from these plants until they have died down. Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

**PROTECTION OF WILDFISH, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

**DO NOT** store near feedstuffs, fertilizers or seed.  
Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.  
If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

**SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS). Which can be obtained from the supplier.

**CONDITIONS OF SALE**

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.





APVMA Approval No: 61893/0507  
Application Number: 41728  
Product Name: Conquest Hatched Herbicide  
Pack Size: Leaflet  
Dimensions: 200mm x 700mm  
Print Size: A4 - 41%  
A3 - 58%