


MAIN PANEL

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



For non-residual control of certain broadleaf weeds in Liberty Link[®] cotton varieties as specified in the DIRECTIONS FOR USE table



RLP
Approved

*** L**

(label code)

* 20, 100, 110 litre

REAR PANEL - continued**Export of Treated Produce**

Maximum Residue Limits (MRLs) may be lower or not be established in export markets for cottonseed. Glufosinate-ammonium residues are not found in cottonseed oil when Liberty 200 Herbicide is used according to label directions. Check with Bayer CropScience Pty Ltd prior to using this product if the cottonseed is destined for export.

GENERAL INSTRUCTIONS

Liberty 200 Herbicide is a non-volatile herbicide with selective activity against many annual and perennial weeds in Liberty Link cotton varieties that are specially developed to be tolerant to Liberty 200.

Liberty 200 is absorbed by plant foliage and green stems. It is inactive in soil and does not provide residual weed control. Liberty 200 is not significantly translocated as an active herbicide throughout the plant, and therefore will only kill that part of a green plant that is contacted by spray. Best results are achieved when applications are made under good growing conditions to young, actively growing weeds. Liberty 200 will have an effect on weeds that are larger than the recommended leaf stage, but speed of activity and control may be reduced. Application to weeds under stress should be avoided. Visible symptoms of control appear in 3 to 7 days, but complete desiccation may take 20 to 30 days under cool conditions.

Liberty Link Cotton and Liberty 200 Herbicide Crop Management Plan

A detailed management plan has been developed that outlines sound agronomic practices and integrated weed management principles and requires growers to implement a preventative weed resistance management strategy. The requirements of the plan are designed to optimise the performance of Liberty Link cotton and minimise the potential for the development of herbicide resistance in weed populations.

The *Liberty Link Cotton and Liberty 200 Herbicide Crop Management Plan* should be consulted prior to the use of this product on Liberty Link cotton. Implementation of the management plan is an essential part of herbicide resistance management. This product is supplied by Bayer CropScience on condition that it is used in conjunction with the *Liberty Link Cotton and Liberty 200 Herbicide Crop Management Plan* (available from www.bayercropscience.com.au).

Pre-plant Paddock Preparation

Control all existing weeds by cultivation or by using a sufficient rate of knockdown herbicide. This may be a glyphosate or paraquat plus diquat based product.

Climatic Conditions

Best results are achieved when Liberty 200 Herbicide is applied under warm humid conditions, (e.g. temperatures below 33° C with a relative humidity above 50 %). Under any other conditions efficacy and speed of action of Liberty 200 may be reduced.

Use of Surfactant/Wetting agent

The addition of a surfactant is not required with Liberty 200 Herbicide.

Mixing Instructions

Liberty 200 Herbicide mixes easily with water. Clean water should always be used for mixing with Liberty 200. Ensure that the spray tank is free of any residues of previous spray materials. Two-thirds fill the spray tank with clean water, and with agitator operating add the required amount of Liberty 200. Add other relevant compatible products. Top the tank up to the required volume with clean water with agitator running.

REAR PANEL - continued**PRECAUTIONS**

Minimum recropping intervals apply for crops following Liberty 200 application.

For advice on crops not listed below, contact the manufacturer, Bayer CropScience Pty Ltd.

Crop	Minimum recropping interval
Cotton	24 hours

Re-entry period

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overall buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT apply on desirable foliage (other than Liberty Link® cotton varieties) or allow drift onto desirable plants, as damage may occur.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with this product or the used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

20 litre pack

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. Do not re-use empty container for any other purpose.

100 and 110 litre pack

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Bayer CropScience Pty Ltd.