



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FAUDEL FUNGICIDE

ACTIVE CONSTITUENT: 720 g/L CHLOROTHALONIL

GROUP	Y	FUNGICIDE
--------------	----------	------------------

**For control of fungus diseases on Almonds, Bananas, Cucurbits, Grapes,
Ornamentals, Peanuts, Stone Fruits, Tobacco
and Vegetables as specified in the Directions for Use table.**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

Contents: 1L [5L, 20L]

AAKO AUSTRALIA PTY LTD
ACN 122 279 109
Suite 1, Level 4, Building B, 207 Pacific Highway
St Leonards NSW 2065 Australia
Tel: (02) 9431 7800 Fax: (02) 9431 7700



STORAGE AND DISPOSAL (1 L)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.



STORAGE AND DISPOSAL (5 L, 20 L)

Store in the closed, original container in a well-ventilated area as cool as possible, out of direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Add rinsings to spray tank. Dispose of the rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. **For refillable containers**, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Attacks eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When preparing spray and using the prepared spray wear cotton overalls, buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves and goggles and disposable mist mask. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FAUDEL is available from Aako Australia Pty Ltd on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FAUDEL FUNGICIDE being beyond the control of the manufacturer, no warranty expressed or implied is given by Aako Australia Pty Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Aako Australia Pty Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

**IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY
PHONE 1800 024 973 TOLL FREE – ALL HOURS – AUSTRALIA WIDE**

Batch No: Date of Manufacture:

APVMA Approval No: 62030/1L/0507
 62030/5L/0507
 62030/20L/0507



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FAUDEL FUNGICIDE

ACTIVE CONSTITUENT: 720 g/L CHLOROTHALONIL

GROUP	Y	FUNGICIDE
--------------	----------	------------------

**For control of fungus diseases on Almonds, Bananas, Cucurbits, Grapes, Ornamentals,
Peanuts, Stone Fruits, Tobacco
and Vegetables as specified in the Directions for Use table.**

**IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER.
READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.**

AAKO AUSTRALIA PTY LTD
ACN 122 279 109
Suite 1, Level 4, Building B, 207 Pacific Highway
St Leonards NSW 2065 Australia
Tel: (02) 9431 7800 Fax: (02) 9431 7700



DIRECTIONS FOR USE

RESTRAINTS: DO NOT mix spraying oils or wetting agents with FAUDEL.

For all stone fruit except Cherries, following additional restraints apply:

DO NOT use oils after bud-swell and during the season if FAUDEL is to be used after shuck fall.

DO NOT tank mix FAUDEL with EC formulations when spraying after shuck fall.

DO NOT exceed 3.2 L of FAUDEL per hectare of crop.

1. TREE AND VINE CROPS

RATE This table shows rates for dilute spraying. For concentrate spraying refer to the Mixing/Application Section.				CRITICAL COMMENTS For uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use at rates greater than 5 times the dilute spraying rate.	
SITUATION & CROP	DISEASE	RATE	WHP (DAYS)		
Almonds (ACT, NSW, Vic, SA, WA, Tas only)	Shot-hole (<i>Stigmia carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)	160 mL/100L	-	Apply at bud-swell, bud-burst, pink-bud, shuck fall, cap fall. Apply every 10-14 days. Apply 1 week pre-harvest.	
Apricots (ACT, NSW, Vic, SA, WA, Tas only)	Brown Rot - Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)		7	Apply at bud-swell, bud-burst, pink bud, full bloom.	Spray applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.
	Stone Fruit Rust (<i>Tranzschelia discolor</i>), Shot-hole (<i>Stigmia carpophila</i>) Freckle (<i>Venturia carpophila</i>)			Apply at bud-swell, bud-burst, pink bud, shuck fall, cap fall, then apply every 10-14 days	
Cherries	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)	160 mL/100 L	7	Apply at bud-swell, bud-burst, pink-bud, early blossom, full bloom. Apply 3 weeks pre-harvest and 1 week pre-harvest.	
	Shot-hole (<i>Stigmia carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)			Apply at budswell, bud-burst, pink bud, shuck fall, cap fall, then every 10-14 days. Apply 1 week pre-harvest.	
	Transit Rot (<i>Rhizopus stolonifer</i>)			Apply 1 week pre-harvest.	
Grapes Note: Russetting of some table grape varieties may occur	Downy Mildew (<i>Plasmopara viticola</i>), Bunch Rot (<i>Botrytis cinerea</i>)	160-210 mL/100 L	Dessert 7 Wine 14	Make the first application as soon as conditions favour the development of he disease. Apply at 7-14 day intervals particularly during wet weather and periods of rapid growth. For Bunch Rot, applications at early full bloom, two weeks later, at veraisson and two weeks prior to harvest. Table Grapes – Russetting of some varieties may occur. Use only up to cap fall in these varieties.	
	Black Spot (Anthracnose) (<i>Elsinoe ampelina</i>)	210 mL/100 L	Dessert 7 Wine 14	If Black Spot has occurred in the vineyard in any of the last three years, apply at budburst and 14 days later when shoots are 10-15 cm long. If wet/humid conditions persist spray again when shoots are 20-30 cm long to protect new growth. Some initial minimal leaf damage on the new shoots of sensitive varieties can occur if spray treatments are applied at budburst and in the first 14 days after budburst. In vineyards with no history of Black Spot, make the first application as soon as conditions favour the development of the disease on the foliage and then at 7-14 day intervals if wet/humid conditions persist. Applications up to cap fall will not cause russetting in table grape and dried fruit varieties.	
Nectarines (NSW, ACT, Vic, Tas, SA, WA, NT only)	Shot-hole (<i>Stigmia carpophila</i>)	160 mL/100 L	7	Apply at bud-swell, pink bud, shuck fall and then repeat each 14 days if weather conditions favour disease.	Spray applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.
	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)			Apply at bud-swell, bud-burst, pink bud, early blossom and full bloom.	
Peaches	Brown Rot – Fruit (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)	160 mL/ 100 L	7	Apply at bud-swell, pink bud, early blossom and full bloom.	Spray applications later than 35 days prior to harvest may result in unacceptable fruit skin damage. See also Additional Restraints for stone fruits.
	Shot-hole (<i>Stigmia carpophila</i>), Stone Fruit Rust (<i>Tranzschelia discolor</i>)			Apply at bud-swell and continue at 7-14 day intervals.	



	Leaf Curl (<i>Taphrina deformans</i>)			Apply at bud-swell and continue at 7-14 day intervals. In Qld spray only at budswell.	
Plums	Brown Rot – Fruit (<i>Monilinia fructicola</i>) Blossom Blight (<i>Monilinia laxa</i>)	160 mL/100 L	1	Apply at bud-swell, bud-burst, early blossom, pink bud and full bloom.	
	Stone Fruit Rust (<i>Tranzschelia discolor</i>), Shot-hole (<i>Stigmina carpophila</i>)			Apply at bud-swell, bud-burst, pink bud, shuck fall and cap fall.	

2. LEGUMES

SITUATION & CROP	DISEASE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Beans – Broad (Faba)	Chocolate Spot (<i>Botrytis fabae</i>), Rust (<i>Uromyces viciae-fabae</i>)	1.4-2.3 L/ha	7	Make the first application as soon as conditions favour the development of the disease. Repeat at 7-14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.
Peanuts (Qld, WA, NSW, NT only)	Early Leaf Spot (<i>Cercospora arachidicola</i>), Late Leaf Spot (<i>Cercosporidium personatum</i>), Peppery Leaf Spot (<i>Leptosphaerulina trifolii</i>), Net Blotch (<i>Didymosphaeria arachidicola</i>), Peanut Rust (<i>Puccinia arachidis</i>)	1.1-1.8 L/ha	–	Use higher rate for Peanut Rust. South Burnett, NSW and WA: commence application as soon as conditions favour the development of the disease and repeat as necessary to maintain control. Atherton Tableland: commence application six weeks after planting – repeat every two weeks.
Peas (Tas, Qld, WA, NT only)	Downy Mildew (<i>Peronospora viciae</i>)	1.1-1.8 L/ha	7	Make the first application as soon as conditions favour the development of the disease. Repeat at 7-14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.

3. VEGETABLES & FRUIT

SITUATION & CROP	DISEASE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Artichokes – Globe	Grey Mould (<i>Botrytis cinerea</i>)	1.8-2.3 L/ha	1	Apply at 7-10 day intervals as required. Use the higher rate if disease pressure is high.
Bananas (Qld, NSW, NT, WA only)	Leaf Spot (<i>Mycosphaerella musicola</i>)	1.8 L/ha	1	DO NOT apply to bananas within 14 days of applying an oil based spray or crop damage will occur. For North Qld – Spray every 14 days late January to end of March. Extend to 21 days after March and then 28-35 days in the drier months of the year. For South Qld – Spray every 21 days from start of December to end of April. For NSW – Spray every 21 days from end of November to end of May. For WA – Spray every 10-14 days from November to end of May and extend to 14-21 days from May to November.
Bananas (WA only)		1.5 L/ha		
Bananas (Qld, NSW, WA, NT only)	Leaf Speckle (<i>Mycosphaerella musae</i>)	3L/ha		
Broccoli, Brussels Sprouts, Cabbage, Cauliflowers	Ring Spot (<i>Mycosphaerella brassicicola</i>)	1.8-3.5L/ha	7	Make the first application as soon as conditions favour the development of the disease. Repeat at 14 day intervals. Use higher rate if disease pressure is high
Capsicums, Peppers	Grey Mould (<i>Botrytis cinerea</i>), Anthracnose (<i>Colletotrichum acutatum</i>)	1.8-2.3 L/ha	1	Apply at 7-10 day intervals as required. Use the higher rate if disease pressure is high.
Carrots	Leaf Blight (<i>Alternaria dauci</i>)	1.8 L/ha	7	Commence application at first sign of disease and then spray every 7-14 days.
Celery	Septoria Leaf Spot (<i>Septoria apiicola</i>)		1	Apply every 7-14 days. Apply also to seed beds.
Celery (NSW, WA, ACT only)	Cercospora Early Blight (<i>Cercospora apii</i>)	1.2 L/ha		Apply every 10-14 days.



Cucurbits including Cantaloupe, Cucumbers, Honeydew, Muskmelon, Pumpkin, Squash, Rockmelon, Watermelon, Zucchini	Downy Mildew (<i>Peronospora cubensis</i>), Gummy Stem Blight (<i>Didymella bryoniae</i>), Anthracnose (<i>Collectotrichum orbiculare</i>), Leaf Blight (<i>Alternaria cucumerina</i>), Target Leaf Spot (<i>Cercospora citrulina</i>)	1.6-2.5 L/ha		Apply on a 7-14 day schedule. Apply rates towards the higher end of the recommended range and at 7 day intervals when disease pressure is high. Commence spray program early, at first true leaf, and before vining or flowering. When used at these rates, FAUDEL FUNGICIDE will suppress Powdery Mildew
	Belly Rot (<i>Rhizoctonia solani</i>)			If applying Faudel Fungicide for the above foliar disease useful suppression of Belly Rot will be obtained where fruit form on soil covered by earlier recent Faudel Fungicide spraying.
Endive	Grey Mould (<i>Botrytis cinerea</i>)	1.8-2.3 L/ha		Apply at 7-10 day intervals as required. Use the higher rate if disease pressure is high.
Leeks, Shallots	Purple Blotch (<i>Alternaria porri</i>)			Make the first application as soon as conditions favour the development of the disease. Repeat at 7-10 day intervals, particularly during wet weather.
Okra (Qld, WA only)	Leaf Disease/Spot			Make the first application as soon as conditions favour the development of the disease. Repeat at 7-10 day intervals.
Onions (excluding spring onions)	Downy Mildew (<i>Peronospora destructor</i>)	2.3 L/ha	14	Make the first application as soon as conditions favour the development of the disease. Repeat at 14 day intervals.
Potato	Early Blight or Target Spot (<i>Alternaria solani</i>), Late Blight or Irish Blight (<i>Phytophthora infestans</i>)	1.1-1.8 L/ha	–	Commence applications at flowering time or earlier if weather conditions are favourable for disease infection. Spray every 7-14 days until within 14 days of desiccation or harvest.
Rhubarb	Ascochyta Leaf Spot (<i>Ascochyta rhei</i>)	140 mL/ 100 L Apply 2000 L/ha	7	Apply every 10-14 days.
Radish	Grey Mould (<i>Botrytis cinerea</i>)	1.8-2.3 L/ha	1	Apply at 7-10 day intervals as required. Use the higher rate if disease pressure is high.
Sweet corn	Turcicum Leaf Blight (<i>Exserohilum turcicum</i>)			Make the first application as soon as conditions favour the development of the disease. Repeat at 7-10 day intervals.
Tomato	Early Blight or Target Spot (<i>Alternaria solani</i>), Leaf Mould (<i>Fulvia fulva</i>) Grey Leaf Spot (<i>Stemphylium solani</i>) Botrytis Rot or Grey Mould (<i>Botrytis cinerea</i>)			Make the first application as soon as conditions favour the development of the disease. Repeat at 7-10 day intervals. Continue into the picking season to obtain best results against Leaf Mould and Botrytis Rot.
Water Cress (Qld, WA only)	Cercospora Leaf Spot			Apply at 7-10 day intervals as required. Use the higher rate if disease pressure is high

4. MISCELLANEOUS

SITUATION & CROP	DISEASE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Ornamentals, Chrysanthemums, Geraniums, Iris and other ornamentals	Botrytis, Grey Mould, Rust diseases, Septoria and Alternaria Leaf Spot diseases	100 mL/100L	-	Apply at 7-10 day intervals. Use 7 day under severe conditions. Avoid application to blooms in Azalea, Hydrangea and Petunia..
Tobacco (Qld, WA only)	Tobacco Leaf Spot (<i>Ascochyta arida</i>)	220 mL/100L	–	Apply every 7 days in seed bed application only.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS (WHP):

Artichoke, Bananas, Capsicum, Celery, Cucurbits, Endive, Leek, Okra, Peppers, Plums, Radish, Shallot, Sweet Corn, Tomatoes, Watercress:

DO NOT HARVEST FOR 1 DAY AFTER APPLICATION



Apricots, Beans, Broccoli, Brussels
Sprouts, Cabbages, Carrots,
Cauliflowers, Cherries, Nectarines,
Peaches, Peas, Rhubarb, Table
Grapes:
Onions, Wine Grapes:
Beans, Peanuts, Peas:
Potatoes:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.
DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.
DO NOT GRAZE LIVESTOCK ON TREATED CROPS.
**WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS
DIRECTED**

GENERAL INSTRUCTIONS

Note: FAUDEL FUNGICIDE is a protectant fungicide only. Therefore to achieve best results, FAUDEL FUNGICIDE must be applied before diseases occur/become established. Use disease warning services or past history to determine timing of first spray. If diseases are already present then an eradicant product should be used first and mixed with FAUDEL FUNGICIDE.

MIXING / APPLICATION

Slowly invert container several times to ensure uniform mixture. Mix the required quantity of product with water in the spray tank with agitator operating. Apply in high volume, low volume or through mister or aircraft spray equipment. Thorough uniform coverage is essential for effective disease control and protection.

Dilute Spraying:

Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions For Use table for each 100 L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying:

Use a sprayer designed and set up for the concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.

Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above:
For example 1500 L/ha.
2. Your chosen concentrate spray volume:
For example 500 L/ha.
3. The concentration factor in this example is:
 $3 \times$ (ie $1500 \text{ L} \div 500 \text{ L} = 3$).
4. If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3×10 , that is 30 mL/100 L of concentrate spraying.
The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

SUGGESTED SPRAY VOLUMES PER HECTARE

High Volume: 500 litres or more.
Mister: 75 to 255 litres.
Low Volume: 200 litres or more.
Aerial Application: 20 litres or more.

COMPATIBILITY

This product is compatible with wettable powder formulations of the most commonly used fungicides, insecticides



and miticides. DO NOT combine with oil-based emulsifiable or flowable pesticides, unless prior experience has shown the combination to be physically compatible and non-injurious to your crop.

This product should not be mixed with spraying oils or sprayed onto crops that have been sprayed with oil for at least 10 days after the oil spray. Oils should not be sprayed on crops treated with this product for at least 10 days after the last spray. Wetting agents have not improved performance. Under some conditions, certain surfactants may cause plant injury.

FUNGICIDE RESISTANCE WARNING

GROUP	Y	FUNGICIDE
-------	---	-----------

For fungicide resistance FAUDEL Fungicide is a Group Y fungicide. Some naturally occurring individual fungi resistant to FAUDEL FUNGICIDE and other Group Y fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by FAUDEL FUNGICIDE and other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of FAUDEL FUNGICIDE to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT contaminate feed or foodstuffs with this product or empty containers.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL (1 L)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE AND DISPOSAL (5 L, 20 L)

Store in the closed, original container in a well-ventilated area as cool as possible, out of direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Add rinsings to spray tank. Dispose of the rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. **For refillable containers**, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Attacks eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When preparing spray and using the prepared spray wear cotton overalls, buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves and goggles and disposable mist mask. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water.

**MSDS**

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FAUDEL FUNGICIDE is available from Aako Australia Pty Ltd on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE:

The use of FAUDEL FUNGICIDE Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Aako Australia Pty Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Aako Australia Pty Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval No: 62030/0507