



Product Name: GENFARM TOLEX LIQUID FUNGICIDE
APVMA Approval No: 62158/122794

Label Name:	GENFARM TOLEX LIQUID FUNGICIDE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	500 g/L TOLCLOFOS-METHYL
Mode of Action:	GROUP 14 FUNGICIDE
Statement of Claims:	For the control of Rhizoctonia solani in Cotton and seed borne Rhizoctonia solani In Potatoes and Rhizoctonia solani in lettuce.
Net Contents:	1L 5L 20L
Restrains:	
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	COTTON, POTATOES: NOT REQUIRED WHEN USED AS DIRECTED. LETTUCE: HARVEST: NOT REQUIRED WHEN USED AS DIRECTED. GRAZING: DO NOT FEED TREATED PRODUCE TO LIVESTOCK.

Trade Advice:	
---------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS</p> <p>COTTON: TOLEX Liquid does not control Pythium spp. and should only be applied to seed previously treated with a fungicide such as Apron* for control of Pythium spp. TOLEX Liquid may be applied to seed previously treated with quintozone, thiodicarb or acephate.</p> <p>Mixing Add the required amount of TOLEX Liquid to the partly filled tank with the agitator running and complete filling the tank with water. Do not mix with alkaline water. Continue thorough agitation during spraying and after stoppage. Do not let prepared spray solution sit in spray tank overnight.</p> <p>Lettuce: Add the required amount of TOLEX Liquid to the partly filled tank with the agitator running and complete filling the tank with water. Do not mix with alkaline water. Continue thorough agitation during spraying and after stoppage. Do not let prepared spray solution sit in spray tank overnight.</p> <p>Application TOLEX Liquid can be applied as an in furrow spray or by water injection using 20 to 350 L/ha of water irrespective of spray band width.</p> <p>POTATOES: TOLEX Liquid ensures protection of potatoes from seed borne Rhizoctonia solani. Best results will be obtained where other crop management practices (including crop rotation) are employed to reduce Rhizoctonia solani infection in the soil.</p> <p>Application Mix the required quantity of TOLEX Liquid with 5 L of water per tonne of seed. Ensure constant agitation of the spray tank during application. Apply only when seed pieces are being rotated using spraying methods designed to ensure complete coverage. Only treat enough seed to meet planting needs. Ensure the seed is dry prior to planting. DO NOT plant wet seed. Ensure that treated seed remains separated from untreated seed at all times.</p>
-----------------------	--

Resistance Warning:	<p>FUNGICIDE RESISTANCE WARNINGS</p> <p>GROUP 14 FUNGICIDE For fungicide resistance management TOLEX Liquid is a Group 14 Fungicide. Some naturally occurring individual fungi resistant to TOLEX Liquid and other Group 14 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by TOLEX Liquid or any other Group 14 fungicides, thus resulting in a reduction of efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Landmark Operations Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.</p>
---------------------	--

Precautions:	<p>PRECAUTIONS</p> <p>DO NOT use treated seed for animal or human consumption. DO NOT allow treated seed to contaminate grain or other seed intended for animal or human consumption. DO NOT feed treated seed or otherwise expose to wild life or domestic birds. When treated seed potatoes are stored they should be kept apart from other potatoes and the bags or containers should be clearly marked to indicate the contents have been treated</p>
--------------	---

	<p>with this product. Bags and containers which have held treated seed potatoes are not to be used for any other purpose.</p> <p>RE-ENTRY TO TREATED AREA The following conditions MUST be observed when re-entry is required before the spray has dried: Wear impervious boots. Do not re-use footwear until thoroughly aired.</p> <p>RE-HANDLING TREATED SEED The following conditions MUST be observed when handling treated seed prior to planting. Wear cotton overalls buttoned to the neck and wrists (or equivalent clothing) and elbow-length PVC gloves. After each day's use, wash gloves and clothing.</p>
--	--

Protections:	<p>PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT feed treated seed to animals including poultry. Dangerous to fish. DO NOT contaminate streams, rivers or watercourses with the product or used containers.</p>
--------------	--

Storage and Disposal:	<p>Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.</p> <p>This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple rinse containers before disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap.</p> <p>Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection point or similar container management program site. The cap should not be replaced but may be taken separately.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.</p> <p>For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.</p>
-----------------------	---

Safety Directions:	<p>May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use. When mixing and loading wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. When using in enclosed areas wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. Ensure adequate ventilation during use. After each day's use, wash gloves and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

DC12-39329154_Genfarm Tolex Liquid Fungicide_DFU_20191125

DIRECTIONS FOR USE

CROP	DISEASE	STATES	RATE	CRITICAL COMMENTS
Cotton	<i>Rhizoctonia solani</i>	QLD and NSW only	120 mL/ha or 120 mL/10,000 m row	Apply as an in-furrow spray or by water injection at time of planting.
Lettuce (Field grown)	Bottom rot (<i>Rhizoctonia solani</i>)	All States	120 mL/ha OR 120 mL/10,000 m row	Apply a maximum of one application per crop by in furrow spray or equivalent with spray volume of 20-350 L/ha. Apply as an in-furrow spray or plug hole drench at transplanting.
Potatoes	Seed borne <i>Rhizoctonia solani</i>	All States	0.4 L/tonne of seed	Do NOT apply to freshly cut seed potatoes as emergence may be affected. Allow sufficient time for cut wounds to dry and suberise before applying TOLEX Liquid or use "round" seed. DO NOT plant wet seed. Apply immediately prior to planting (within 1 day) by spray methods and allow seed to dry before planting. Seed pieces should be rotated on a roller system (e.g. grading table) to ensure complete and thorough coverage. For further information refer to the General Instructions. If conditions (e.g. high soil temperature and moisture) or seed quality favour seed piece breakdown, the addition of any moisture to the seed may enhance the problem. Under these circumstances the use of Tizolex 100D is recommended.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.