

100% on A3

**YOUR PROOF**

job#: 155339  
name: 02 QPE 100 5L Front  
client: Genfarm  
date: 04/3/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.  
PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC**® Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

Dieline

PMS 3995

PMS 348

PMS 654

Black

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Genfarm**  
**QPE 100**  
**HERBICIDE**

ACTIVE CONSTITUENT:  
99.5g/L QUIZALOFOP-P-ETHYL  
SOLVENT: 801 g/L HYDROCARBON LIQUID

**GROUP A HERBICIDE**

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE



**CONTENTS: 5 LITRES**



140mm deep



140mm wide



100% on A3



# Genfarm QPE 100 HERBICIDE

#### STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

#### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

APVMA Approval No: 62508/5L/0608


Genfarm Crop Protection Pty Ltd  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

#### CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

ENVIRONMENTALLY HAZARDOUS SUBSTANCES,  
LIQUID, N.O.S.  
(CONTAINS Quizalofop-p-ethyl)

UN NO. 3082  
Packaging Group III  
Hazchem 2X



FOR SPECIALIST ADVICE  
IN AN EMERGENCY DIAL  
**1800 033 111**  
24 HOURS AUSTRALIA WIDE

IN A TRANSPORT EMERGENCY  
**DIAL 000**  
POLICE OR FIRE BRIGADE



PHILSTIC MULTIFOLD

137mm deep

125mm wide

### YOUR PROOF

job#: 155339  
name: 03 QPE 100 5L Base  
client: Genfarm  
date: 11/06/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.

PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC** Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

Dieline

Book area (138mm x 110mm)

Black



**Genfarm**

**QPE 100**  
**HERBICIDE**

PEEL  
HERE

140mm deep

**STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

APVMA Approval No: 62508/5L/0608

Genfarm Crop Protection Pty Ltd  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92



**CONDITIONS OF SALE**

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

<p><b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS Quizalofop-p-ethyl)</b></p>	
<p><b>UN NO. 3082</b> Packaging Group III Hazchem 2X</p>	<p><b>FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111</b> 24 HOURS AUSTRALIA WIDE</p>
<p><b>IN A TRANSPORT EMERGENCY DIAL 000</b> POLICE OR FIRE BRIGADE</p>	

Batch No:

Date of Manufacture:



140mm wide

100% on A3

**YOUR PROOF**

job#: 155339  
name: 03 QPE 100 5L Top  
client: Genfarm  
date: 11/06/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC** Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

Dieline

PMS 3995

PMS 348

PMS 654

Black

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Genfarm**  
**QPE 100**  
**HERBICIDE**

ACTIVE CONSTITUENT:  
99.5 g/L QUIZALOFOP-P-ETHYL  
SOLVENT: 801 g/L HYDROCARBON LIQUID

**GROUP A HERBICIDE**

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE



**CONTENTS: 10 LITRES**



170mm deep

170mm wide

100% on A3

**YOUR PROOF**

job#: 1555339  
name: 03 QPE 100 10L Front  
client: Genfarm  
date: 6/06/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.  
PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC**® **Philstic Labels Ltd**  
Phone +64 9 630 5822  
Fax +64 9 630 0957

 Dieline

 PMS 3995

 PMS 348

 PMS 654

 Black



**APVMA**  
**APPROVED LABEL**

# Genfarm

# QPE 100

## HERBICIDE

### STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

APVMA Approval No: 62508/10L/0608

**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

### CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**ENVIRONMENTALLY HAZARDOUS SUBSTANCES,  
LIQUID, N.O.S.  
(CONTAINS Quizalofop-p-ethyl)**

**UN NO. 3082**  
**Packaging Group III**  
**Hazchem 2X**



**FOR SPECIALIST ADVICE  
IN AN EMERGENCY DIAL  
1800 033 111  
24 HOURS AUSTRALIA WIDE**

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**



PHILSTIC MULTIFOLD

167mm deep

155mm wide

100% on A3

### YOUR PROOF

job#: 155339  
name: 03 QPE 100 10L Base  
client: Genfarm  
date: 11/06/2008

**This proof is NOT a true representation of the final colours that will be printed**

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED**

**PHILSTIC** Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

Dieline

Book area (168mm x 140mm)

Black



**APVMA  
APPROVED  
LABEL**

**Genfarm**

# QPE 100

## HERBICIDE

**STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

**CONDITIONS OF SALE**

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**ENVIRONMENTALLY  
HAZARDOUS SUBSTANCES,  
LIQUID, N.O.S.  
(CONTAINS Quizalofop-p-ethyl)**

**UN NO. 3082  
Packaging Group III  
Hazchem 2X**

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**



**FOR SPECIALIST ADVICE  
IN AN EMERGENCY DIAL  
1800 033 111  
24 HOURS AUSTRALIA WIDE**

APVMA Approval No: 62508/10L/0608  
**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92



Batch No:

Date of Manufacture:



PEEL  
HERE

170mm deep

170mm wide

100% on A3

**YOUR PROOF**

job#: 155339  
name: 03 QPE 100 10L Top  
client: Genfarm  
date: 11/06/2008






**This proof is NOT a true representation of the final colours that will be printed**

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO responsibility will be taken for errors not noted on this signed proof.**

**PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED**

**PHILSTIC® Philstic Labels Ltd**  
Phone +64 9 630 5822  
Fax +64 9 630 0957

-  Dieline
-  PMS 3995
-  PMS 348
-  PMS 654
-  Black

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Genfarm**  
**QPE 100**  
**HERBICIDE**

ACTIVE CONSTITUENT:  
99.5g/L QUIZALOFOP-P-ETHYL  
SOLVENT: 801 g/L HYDROCARBON LIQUID

**GROUP A HERBICIDE**

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE



**CONTENTS: 20 LITRES**



170mm deep

170mm wide

100% on A3

**YOUR PROOF**

job#: 155339  
name: 02 QPE 100 20L Front  
client: Genfarm  
date: 04/03/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.  
PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC**® **Philstic Labels Ltd**  
Phone +64 9 630 5822  
Fax +64 9 630 0957


Dieline

PMS 3995

PMS 348

PMS 654

Black



**APVMA  
APPROVED  
LABEL**

# Genfarm QPE 100 HERBICIDE

**STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

APVMA Approval No: 62508/20/0608

**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

**CONDITIONS OF SALE**

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**ENVIRONMENTALLY HAZARDOUS SUBSTANCES,  
LIQUID, N.O.S.  
(CONTAINS Quizalofop-p-ethyl)**

**UN NO. 3082  
Packaging Group III  
Hazchem 2X**



**FOR SPECIALIST ADVICE  
IN AN EMERGENCY DIAL  
1800 033 111  
24 HOURS AUSTRALIA WIDE**

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**



PHILSTIC MULTIFOLD

167mm deep

155mm wide

100% on A3

**YOUR PROOF**

job#: 155339  
name: 04 QPE 100 20L Base  
client: Genfarm  
date: 11/06/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.

**PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED**

**PHILSTIC** Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

Dieline

Book area (168mm x 140mm)

Black





# Genfarm

# QPE 100

## HERBICIDE

PEEL HERE

**STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

**CONDITIONS OF SALE**

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

<p><b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS Quizalofop-p-ethyl)</b></p>	
<p><b>UN NO. 3082</b> Packaging Group III Hazchem 2X</p>	
<p><b>IN A TRANSPORT EMERGENCY DIAL 000</b> POLICE OR FIRE BRIGADE</p>	<p><b>FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111</b> 24 HOURS AUSTRALIA WIDE</p>

APVMA Approval No: 62508/20/0608

**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92



Batch No:

Date of Manufacture:



170mm deep

170mm wide

100% on A3

**YOUR PROOF**

job#: 155339  
 name: 04 QPE 100 20L Top  
 client: Genfarm  
 date: 11/06/2008

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required

sign: \_\_\_\_\_  
 date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC** Philstic Labels Ltd  
 Phone +64 9 630 5822  
 Fax +64 9 630 0957

- Dieline
- PMS 3995
- PMS 348
- PMS 654
- Black



# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Genfarm

# QPE 100

## HERBICIDE

ACTIVE CONSTITUENT:  
99.5 g/L QUIZALOFOP-P-ETHYL  
SOLVENT: 801 g/L HYDROCARBON LIQUID

**GROUP A HERBICIDE**

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE**

Batch No:  
Date of Manufacture:



**CONTENTS: 110 LITRES**



170mm deep

170mm wide

100% on A3

### YOUR PROOF

job#: 169854  
name: 01 QPE 100 110L Front  
client: Genfarm  
date: 28/10/2009

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_  
date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.  
**PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED**

**PHILSTIC** Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

-  Dieline
-  PMS 3995
-  PMS 348
-  PMS 654
-  Black

100% on A3

# Genfarm

# QPE 100

## HERBICIDE

**STORAGE AND DISPOSAL**  
Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage

**SAFETY DIRECTIONS**  
Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**  
If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.


**MATERIAL SAFETY DATA SHEET**  
Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.


APVMA Approval No: 62508/110/0809

**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

**CONDITIONS OF SALE**  
The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**ENVIRONMENTALLY HAZARDOUS SUBSTANCES,  
LIQUID, N.O.S.  
(CONTAINS Quizalofop-p-ethyl)**

<p><b>UN NO. 3082</b> Packaging Group III Hazchem 2X</p>	
<p><b>FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 24 HOURS AUSTRALIA WIDE</b></p>	
<p><b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b></p>	



PHILSTIC MULTIFOLD

167mm deep

**169mm wide**

**YOUR PROOF**

job#: 169854

name: 01 QPE 100 110L Base

client: Genfarm

date: 28/10/2009

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed

Further proof required

sign: \_\_\_\_\_

date: \_\_\_\_\_

**NO** responsibility will be taken for errors not noted on this signed proof.  
PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

**PHILSTIC**® *Philstic Labels Ltd*  
Phone +64 9 630 5822  
Fax +64 9 630 0957

- Dieline
- Book area (168mm x 140mm)
- Silicone
- Black

100% on A3

# Genfarm QPE 100 HERBICIDE



#### STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. **For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage

#### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

#### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

APVMA Approval No: 62508/110/0809

**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park NSW 2113  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

#### CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS Quizalofop-p-ethyl)</b>	
<b>UN NO. 3082 Packaging Group III Hazchem 2X</b>	
<b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b>	<b>FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 24 HOURS AUSTRALIA WIDE</b>








170mm deep

170mm wide

YOUR PROOF	
job#:	169854
name:	01 QPE 100 110L Top
client:	Genfarm
date:	28/10/2009
<b>This proof is NOT a true representation of the final colours that will be printed</b>	
<input type="checkbox"/>	OK to proceed
<input type="checkbox"/>	Further proof required
sign:	_____
date:	_____
<b>NO responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED</b>	

**PHIL STIC** Philstic Labels Ltd  
Phone +64 9 630 5822  
Fax +64 9 630 0957

-  Dieline
-  PMS 3995
-  PMS 348
-  PMS 654
-  Black



# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# GENFARM QPE 100 HERBICIDE

ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL

SOLVENT: 801 g/L HYDROCARBON LIQUID

GROUP **A** HERBICIDE

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

IMPORTANT: READ THIS BOOKLET BEFORE USE

APVMA APPROVAL NO: 62508/A/0809

Genfarm Crop Protection Pty Ltd  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park,  
NSW, 2113  
Tel: (02) 9889 5400

## DIRECTIONS FOR USE

### Restrains

**DO NOT** apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).

**DO NOT** apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.

**DO NOT** apply under conditions of high temperature and low humidity.

**DO NOT** apply if rainfall is expected within 3 hours.

**DO NOT** apply less than 50L prepared spray/ha

**BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES.**

Crop	Weeds Controlled	Weeds Stage	Rate/ha	Withholding Period	
				Harvest	Grazing
Beetroot	Annual Ryegrass ( <i>Lolium rigidum</i> ) Brome Grasses ( <i>Bromus spp.</i> )	3 leaf to early tillering	300 mL or 375 mL	14 days	-
Cabbage				9 weeks	-
Canola				11 weeks	4 weeks
Carrots				10 weeks	-
Cauliflower	Barley grass ( <i>Hordeum spp.</i> ) Volunteer Barley Volunteer Wheat	3 leaf to early tillering	250 mL	14 days	-
Chickpeas				12 weeks	4 weeks
Clover & Sub-clover pastures				-	14 days
Cucumbers•				14 days	-
Faba Beans	Wild Oats ( <i>Avena spp.</i> )	3 leaf to mid tillering	125 mL or 250 mL	12 weeks	4 weeks
Field Peas				9 weeks	4 weeks
Honey Dew Melons•				9 weeks	-
Lentils				12 weeks	4 weeks
Lucerne				-	4 weeks
Lupins				6 weeks	4 weeks
Medic pasture				-	14 days
Mung beans				12 weeks	4 weeks
Navy Beans				12 weeks	4 weeks
Onions				18 weeks	-
Potatoes				10 weeks	-
Pumpkins•				9 weeks	-
Radish	21 days	-			
Tomatoes	4 weeks	-			
Vetch	12 weeks	4 weeks			

\* Always add a surfactant/wetting agent at the recommended Rate see section on “use of surfactant/wetting agent” for details.

• Important: DO NOT apply Genfarm QPE 100 Herbicide before the five true leaf stage of the crop.

CRITICAL COMMENTS
Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
Apply when weeds are actively growing*
Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control*

**BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES**

Crop	Weeds Controlled	Weeds Stage	Rate/ha	Withholding Period				
				Harvest	Grazing			
Beetroot	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ) Barnyard Grass ( <i>Echinochloa crusgalli</i> )	3 leaf to early tillering	500 mL or 750 mL	14 days	-			
Cabbage				9 weeks	-			
Cauliflower				14 weeks	-			
Carrots				10 weeks	-			
Cucumbers•				14 days	-			
Green Beans				5 weeks	4 weeks			
Honey Dew				9 weeks	-			
Melons•				Couch grass ( <i>Cynodon dactylon</i> )	2 – 4 leaf seedling	500 mL	-	4 weeks
Lucerne							12 weeks	4 weeks
Mung beans				Dinebra ( <i>Dinebra retroflexa</i> )	3 leaf to early tillering	250 mL or 375 mL	12 weeks	4 weeks
Navy Beans	18 weeks	-						
Onions	11 weeks	11 weeks						
Peanuts	Crowfoot grass ( <i>Eleusine indica</i> )	2 leaf to early tillering	500 mL	10 weeks	-			
Potatoes				7 days	-			
Pineapples				9 weeks	-			
Pumpkins•	Foxtail Millet ( <i>Setaria italica</i> )	2 leaf to early tillering	250 mL or 500 mL	21 days	-			
Radish				12 weeks	4 weeks			
Soybeans				9 weeks	11 weeks			
Sunflower				4 weeks	-			
Tomatoes	Columbus grass ( <i>Sorghum x almum</i> ) Johnson grass ( <i>Sorghum halepense</i> ) Liverseed grass ( <i>Orochloa panicoides</i> )	2 leaf to 3 tiller	500 mL or 750 mL	4 weeks	-			
						Rhodes grass ( <i>Chloris gayana</i> ) Stink grass ( <i>Eragrostis cilianensis</i> )	3 leaf to early tillering	250 mL or 500 mL

**CRITICAL COMMENTS**

Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering\*

Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed\*.

Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering\*

Apply when weeds are actively growing\*.

Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering\*



**BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES (Cont)**

Crop	Weeds Controlled	Weeds Stage	Rate/ha	Withholding Period	
				Harvest	Grazing
Beetroot	Queensland Blue grass ( <i>Dichanthium sericeum</i> )	3 leaf to early tillering	500 mL	14 days	-
Cabbage				9 weeks	-
Cauliflower				14 weeks	-
Carrots				10 weeks	-
Cucumbers•	Summer grass ( <i>Digitaria ciliaris</i> )	3 leaf to early tillering	500 mL or 1000 mL	14 days	-
Green Beans				5 weeks	4 weeks
Honey Dew				9 weeks	-
Melons•				-	4 weeks
Lucerne				12 weeks	4 weeks
Mung beans				12 weeks	4 weeks
Navy Beans				18 weeks	-
Onions				11 weeks	11 weeks
Peanuts				10 weeks	-
Potatoes				7 days	-
Pineapples				9 weeks	-
Pumpkins•				21 days	-
Radish				12 weeks	4 weeks
Soybeans				9 weeks	11 weeks
Sunflower	4 weeks	-			
Tomatoes					

\* Always add a surfactant/wetting agent at the recommended Rate see section on “use of surfactant/wetting agent” for details.

• Important: DO NOT apply Genfarm QPE 100 Herbicide before the five true leaf stage of the crop.

**CRITICAL COMMENTS**

Apply when weeds are actively growing\*.

Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering\*

**HAND GUN SPRAYING: ALL STATES**

Crop	Weeds Controlled	Weeds Stage	Rate/ 100L	Withholding Period				
				Harvest	Grazing			
Beetroot	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ) Canola Crowsfoot grass ( <i>Eleusine indica</i> )	3 leaf to early tillering	125 mL	14 days	-			
Cabbage				9 weeks	-			
Cauliflower				14 days	-			
Canola				11 weeks	4 weeks			
Carrots				10 weeks	-			
Chickpeas				12 weeks	4 weeks			
Clover & Sub- clover pastures				Paspalum ( <i>Paspalum dilatatum</i> )	Seedling to Established Clumps	-	14 days	
Cucumbers•						14 days	-	
Faba Beans				Johnson grass ( <i>Sorghum halepense</i> )	Seedling 2 leaf to early tillering	125 mL	12 weeks	4 weeks
Field Peas							9 weeks	4 weeks
Grapes	-	-						
Green beans	5 weeks	4 weeks						
Honey Dew	9 weeks	-						
Melons•	Rhizome 2 leaf to early tillering	250 mL	12 weeks				4 weeks	
Lentils			-				4 weeks	
Lucerne	Kikuyu ( <i>Pennisetum clandestinum</i> )	Seedling to Established plants	250 mL				6 weeks	4 weeks
Lupins							-	14 days
Medic pasture							12 weeks	4 weeks
Mung beans				12 weeks	4 weeks			
Navy Beans				-	-			
Non crop				18 weeks	-			
Onions				11 weeks	11 weeks			
Peanuts				7 days	-			
Pineapples				10 weeks	-			
Potatoes				9 weeks	-			
Pumpkins•	21 days	-						
Radish	12 weeks	4 weeks						
Soy beans	9 weeks	11 weeks						
Sunflowers	4 weeks	-						
Tomatoes	12 weeks	4 weeks						
Vetch								

**CRITICAL COMMENTS**

Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha. Thoroughly wet target weeds (especially well established clumps) without causing runoff. Repeat spraying may be necessary for perennial grass species.

Always add a surfactant/wetting agent at the recommended rate – see section on “Use of Surfactant/Wetting agent” for details.

- \* Always add a surfactant/wetting agent at the recommended Rate see section on “use of surfactant/wetting agent” for details.
- Important: DO NOT apply Genfarm QPE 100 Herbicide before the five true leaf stage of the crop.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIOD

##### HARVEST

**BEETROOT, CAULIFLOWER, CUCUMBERS: DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.**

**CABBAGE, FIELD PEAS, HONEY DEW MELON, PUMPKINS, SUNFLOWERS: DO NOT APPLY LATER THAN 9 WEEKS BEFORE HARVEST.**

**CARROTS, POTATOES: DO NOT APPLY LATER THAN 10 WEEKS BEFORE HARVEST.**

**CHICKPEAS, FABA BEANS, LENTILS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT APPLY LATER THAN 12 WEEKS BEFORE HARVEST.**

**CANOLA, PEANUTS: DO NOT APPLY LATER THAN 11 WEEKS BEFORE HARVEST.**

**GREEN BEANS: DO NOT APPLY LATER THAN 5 WEEKS BEFORE HARVEST.**

**LUPINS: DO NOT APPLY LATER THAN 6 WEEKS BEFORE HARVEST.**

**ONIONS: DO NOT APPLY LATER THAN 18 WEEKS BEFORE HARVEST.**

**PINEAPPLES: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.**

**RADISH: DO NOT APPLY LATER THAN 21 DAYS BEFORE HARVEST.**

**TOMATOES: DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST.**

##### GRAZING

**CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, GREEN BEANS, LENTILS, LUCERNE, LUPINS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.**

**CLOVER, MEDIC PASTURES (EXCEPT LUCERNE): DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.**

**PEANUTS, SUNFLOWERS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11 WEEKS AFTER APPLICATION.**

#### GENERAL INSTRUCTIONS

Genfarm QPE 100 Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops.

Genfarm QPE 100 Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates from the treated foliage to the root system and growing points of the plant.

Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

#### Resistant Weeds Warning

**GROUP A HERBICIDE**

Genfarm QPE 100 Herbicide is a member of the “Fop” (aryloxyphenoxy propionate) group of herbicides. Genfarm QPE 100 Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management Genfarm QPE 100 Herbicide is a Group A herbicide.

Some naturally occurring weed biotypes resistant to Genfarm QPE 100 Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Genfarm QPE 100 Herbicide or other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Genfarm Crop Protection accepts no liability for any losses that may result from failure of Genfarm QPE 100 Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

#### Time of Application

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to curcubit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and/or grazing or cutting for stockfeed.

**IMPORTANT:** DO NOT apply Genfarm QPE 100 Herbicide to curcubit crops before the five true leaf stage.

## Spray Preparation

If spray equipment has been previously used for the application of sulfonyleurea herbicides eg metsulfuron methyl, Harmony M, chlorsulfuron, triasulfuron, ensure that the recommended sprayer cleanup procedure for the product previously sprayed has been followed, before using the equipment for the application of Genfarm QPE 100 Herbicide.

Genfarm QPE 100 Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the amount of Genfarm QPE 100 directly to the spray tank and mix well. If tank mixing with carbendazim or mancozeb fungicides, add these to the spray tank and mix thoroughly BEFORE adding Genfarm QPE 100 Herbicide.

## Use of a Surfactant/Wetting Agent

Genfarm QPE 100 Herbicide may be used with either of the following adjuvant system:

- 1) A non-ionic surfactant (1000 gai/L strength) at 100 mL/100L (i.e. 0.1% v/v) and a mineral spray oil at 1 L/100 L of final spray volume (i.e. 1% v/v).
- 2) A non-ionic surfactant (1000 gai/L strength) at 200 mL/100L of final spray volume (i.e. 0.2% v/v).

DO NOT use the non-ionic surfactant + mineral oil adjuvant system on vegetable crops, use a non-ionic surfactant only.

DO NOT use a spray oil alone, a surfactant must always be added.

DO NOT use non-methylated spray oils of crop origin as spray adjuvants.

## Compatibilities

Genfarm QPE 100 Herbicide is compatible with the insecticides deltamethrin, omethoate, dimethoate, methomyl, endosulfan, the fungicides carbendazim and mancozeb and the herbicides diflufenican and clopyralid **when using the surfactant only adjuvant system.**

Genfarm QPE 100 Herbicide is compatible with the herbicide sethoxydim and clethodim **when using the surfactant plus spray oil adjuvant system.**

When tank mixing Genfarm QPE 100 Herbicide and broadleaf herbicides, DO NOT use a spray oil, use a surfactant only.

Application of Genfarm QPE 100 Herbicide /diflufenican tank mix may result in crop effects. The effect is temporary and usually appears as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected.

DO NOT add crop oil to tank mixtures of Genfarm QPE 100 Herbicide /diflufenican.

To minimise crop effects apply Genfarm QPE 100 Herbicide a minimum of 5 days prior to diflufenican application or a minimum 10 days after diflufenican application. (14-21 days if weeds have been stressed).

## Ground Spraying

Use sprayers with flat fan or hollow cone nozzles at 210-350 kPa to give an even distribution of droplets with no less than 50% of the spray volume in the 150-300 micron range. 110<sup>9</sup>

flat fan nozzles are preferred. Use between 50 and 100 litres of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150 litres prepared spray/ha. For proper spray coverage adjust the boom and nozzle height according to the specification listed by the nozzle manufacturers. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained. Avoid overlapping boomspray widths and shut off spray booms while starting, slowing or stopping.

## Sprayer cleanup

Thoroughly clean all traces of Genfarm QPE 100 Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops. If spray equipment has been previously used for the application of sulfonyleurea herbicides eg metsulfuron methyl, thifensulfuron methyl, chlorsulfuron, triasulfuron or terbutryn, ensure that the recommended sprayer cleanup procedure of the product previously sprayed has been followed before using the equipment for the application of Genfarm QPE 100 Herbicide.

## Precaution: Re-entry

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of Genfarm QPE 100 Herbicide.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.


### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

### CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**Genfarm Crop Protection Pty Ltd**  
**Suite 2, Level 3, 64 Talavera Road,**  
**Macquarie Park,**  
**NSW, 2113**  
**Tel: (02) 9889 5400**

<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S.</b> (CONTAINS Quizalofop-p-ethyl)	
<b>UN NO. 3082</b> Packaging Group III Hazchem 2X	
<b>IN A TRANSPORT EMERGENCY DIAL 000</b> POLICE OR FIRE BRIGADE	<b>FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111</b> 24 HOURS AUSTRALIA WIDE

**THIS PAGE DOES NOT PRINT**

**Pages @ 100% on A3**



# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# GENFARM QPE 100 HERBICIDE

ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL

SOLVENT: 801 g/L HYDROCARBON LIQUID

**GROUP A HERBICIDE**

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

**IMPORTANT: READ THIS BOOKLET BEFORE USE**

APVMA APPROVAL NO: 62508/B/0809

**Genfarm Crop Protection Pty Ltd  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park,  
NSW, 2113  
Tel: (02) 9889 5400**

## DIRECTIONS FOR USE

### Restrains

**DO NOT** apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).

**DO NOT** apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.

**DO NOT** apply under conditions of high temperature and low humidity.

**DO NOT** apply if rainfall is expected within 3 hours.

**DO NOT** apply less than 50L prepared spray/ha

**BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES.**

Crop	Weeds Controlled	Weeds Stage	Rate/ha	Withholding Period	
				Harvest	Grazing
Beetroot	Annual Ryegrass ( <i>Lolium rigidum</i> ) Brome Grasses ( <i>Bromus spp.</i> )	3 leaf to early tillering	300 mL or 375 mL	14 days	-
Cabbage				9 weeks	-
Canola				11 weeks	4 weeks
Carrots				10 weeks	-
Cauliflower	Barley grass ( <i>Hordeum spp</i> ) Volunteer Barley Volunteer Wheat	3 leaf to early tillering	250 mL	14 days	-
Chickpeas				12 weeks	4 weeks
Clover & Sub-clover pastures				-	14 days
Cucumbers*				14 days	-
Faba Beans	Wild Oats ( <i>Avena spp</i> )	3 leaf to mid tillering	125 mL or 250 mL	12 weeks	4 weeks
Field Peas				9 weeks	4 weeks
Honey Dew Melons*				9 weeks	-
Lentils				12 weeks	4 weeks
Lucerne				-	4 weeks
Lupins				6 weeks	4 weeks
Medic pasture				-	14 days
Mung beans				12 weeks	4 weeks
Navy Beans				12 weeks	4 weeks
Onions				18 weeks	-
Potatoes				10 weeks	-
Pumpkins*				9 weeks	-
Radish				21 days	-
Tomatoes				4 weeks	-
Vetch				12 weeks	4 weeks

CRITICAL COMMENTS
Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
Apply when weeds are actively growing*
Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control*

\* Always add a surfactant/wetting agent at the recommended Rate see section on “use of surfactant/wetting agent” for details.

• Important: DO NOT apply Genfarm QPE 100 Herbicide before the five true leaf stage of the crop.



**BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES**

Crop	Weeds Controlled	Weeds Stage	Rate/ha	Withholding Period	
				Harvest	Grazing
Beetroot	Awnless Barnyard Grass ( <i>Echinochloa colona</i> )	3 leaf to early tillering	500 mL or 750 mL	14 days	-
Cabbage				9 weeks	-
Cauliflower	Barnyard Grass ( <i>Echinochloa crusgalli</i> )			14 weeks	-
Carrots				10 weeks	-
Cucumbers*	Couch grass ( <i>Cynodon dactylon</i> )	2 – 4 leaf seedling	500 mL	14 days	-
Green Beans				5 weeks	4 weeks
Honey Dew	Dinebra ( <i>Dinebra retroflexa</i> )	3 leaf to early tillering	250 mL or 375 mL	9 weeks	-
Melons*				-	4 weeks
Lucerne	Crowfoot grass ( <i>Eleusine indica</i> )	2 leaf to early tillering	500 mL	12 weeks	4 weeks
Mung beans				12 weeks	4 weeks
Navy Beans	Foxtail Millet ( <i>Setaria italica</i> )	2 leaf to early tillering	250 mL or 500 mL	18 weeks	-
Onions				11 weeks	11 weeks
Peanuts	Columbus grass ( <i>Sorghum x alnum</i> )	2 leaf to 3 tiller	500 mL or 750 mL	10 weeks	-
Potatoes				7 days	-
Pineapples	Johnson grass ( <i>Sorghum halepense</i> )			9 weeks	-
Pumpkins*				21 days	-
Radish	Liverseed grass ( <i>Oryzopsis spiculata</i> )			12 weeks	4 weeks
Soybeans				9 weeks	11 weeks
Sunflower	Rhodes grass ( <i>Chloris gayana</i> )	3 leaf to early tillering	250 mL or 500 mL	4 weeks	-
Tomatoes				11 weeks	-
	Stink grass ( <i>Eragrostis ciliaris</i> )				
	Queensland Blue grass ( <i>Dichanthium sericeum</i> )	3 leaf to early tillering	500 mL		
	Summer grass ( <i>Digitaria ciliaris</i> )	3 leaf to early tillering	500 mL or 1000 mL		

CRITICAL COMMENTS
Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed*.
Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
Apply when weeds are actively growing*.
Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
Apply when weeds are actively growing*.
Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*

\* Always add a surfactant/wetting agent at the recommended Rate see section on “use of surfactant/wetting agent” for details.

- Important: DO NOT apply Genfarm QPE 100 Herbicide before the five true leaf stage of the crop.

**HAND GUN SPRAYING: ALL STATES**

Crop	Weeds Controlled	Weeds Stage	Rate/100L	Withholding Period				
				Harvest	Grazing			
Beetroot	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ) Crowsfoot grass ( <i>Eleusine indica</i> )	3 leaf to early tillering	125 mL	14 days	-			
Cabbage				9 weeks	-			
Cauliflower				14 days	-			
Canola				11 weeks	4 weeks			
Carrots				10 weeks	-			
Chickpeas				Paspalum ( <i>Paspalum dilatatum</i> )	Seedling to Established Clumps	12 weeks	4 weeks	
Clover & Sub-clover pastures						-	14 days	
Cucumbers•				Johnson grass ( <i>Sorghum halepense</i> )	Seedling 2 leaf to early tillering	250 mL	14 days	-
Faba Beans							12 weeks	4 weeks
Field Peas							9 weeks	4 weeks
Grapes	-	-						
Green beans	Rhizome 2 leaf to early tillering	5 weeks	4 weeks					
Honey Dew		9 weeks	-					
Melons•	Kikuyu ( <i>Pennisetum clandestinum</i> )	Seedling to Established plants	250 mL				12 weeks	4 weeks
Lentils							-	4 weeks
Lucerne							6 weeks	4 weeks
Lupins							-	14 days
Medic pasture				12 weeks	4 weeks			
Mung beans				12 weeks	4 weeks			
Navy Beans				-	-			
Non crop				18 weeks	-			
Onions				11 weeks	11 weeks			
Peanuts				7 days	-			
Pineapples	10 weeks	-						
Potatoes	9 weeks	-						
Pumpkins•	21 days	-						
Radish	12 weeks	4 weeks						
Soy beans	9 weeks	11 weeks						
Sunflowers	4 weeks	-						
Tomatoes	12 weeks	4 weeks						
Vetch								

CRITICAL COMMENTS
Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha. Thoroughly wet target weeds (especially well established clumps) without causing runoff. Repeat spraying may be necessary for perennial grass species.
Always add a surfactant/wetting agent at the recommended rate – see section on “Use of Surfactant/Wetting agent” for details.

\* Always add a surfactant/wetting agent at the recommended Rate see section on “use of surfactant/wetting agent” for details.

• Important: DO NOT apply Genfarm QPE 100 Herbicide I before the five true leaf stage of the crop.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIOD

##### HARVEST

**BETROOT, CAULIFLOWER, CUCUMBERS: DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.**

**CABBAGE, FIELD PEAS, HONEY DEW MELON, PUMPKINS, SUNFLOWERS: DO NOT APPLY LATER THAN 9 WEEKS BEFORE HARVEST.**

**CARROTS, POTATOES: DO NOT APPLY LATER THAN 10 WEEKS BEFORE HARVEST.**

**CHICKPEAS, FABA BEANS, LENTILS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT APPLY LATER THAN 12 WEEKS BEFORE HARVEST.**

**CANOLA, PEANUTS: DO NOT APPLY LATER THAN 11 WEEKS BEFORE HARVEST.**

**GREEN BEANS: DO NOT APPLY LATER THAN 5 WEEKS BEFORE HARVEST.**

**LUPINS: DO NOT APPLY LATER THAN 6 WEEKS BEFORE HARVEST.**

**ONIONS: DO NOT APPLY LATER THAN 18 WEEKS BEFORE HARVEST.**

**PINEAPPLES: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.**

**RADISH: DO NOT APPLY LATER THAN 21 DAYS BEFORE HARVEST.**

**TOMATOES: DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST.**

##### GRAZING

**CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, GREEN BEANS, LENTILS, LUCERNE, LUPINS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.**

**CLOVER, MEDIC PASTURES (EXCEPT LUCERNE): DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.**

**PEANUTS, SUNFLOWERS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11 WEEKS AFTER APPLICATION.**

##### GENERAL INSTRUCTIONS

Genfarm QPE 100 Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops.

Genfarm QPE 100 Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates from the treated foliage to the root system and growing points of the plant.

Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

#### Resistant Weeds Warning

**GROUP A HERBICIDE**

Genfarm QPE 100 Herbicide is a member of the "Fop" (aryloxyphenoxy propionate) group of herbicides. Genfarm QPE 100 Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management Genfarm QPE 100 Herbicide is a Group A herbicide.

Some naturally occurring weed biotypes resistant to Genfarm QPE 100 Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Genfarm QPE 100 Herbicide or other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Genfarm Crop Protection accepts no liability for any losses that may result from failure of Genfarm QPE 100 Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

##### Time of Application

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to curcubit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and/or grazing or cutting for stockfeed.

**IMPORTANT:** DO NOT apply Genfarm QPE 100 Herbicide to curcubit crops before the five true leaf stage.

##### Spray Preparation

If spray equipment has been previously used for the application of sulfonylurea herbicides eg metsulfuron methyl, Harmony M, chlorsulfuron, triasulfuron, ensure that the recommended sprayer cleanup procedure for the product previously sprayed has been followed, before using the equipment for the application of Genfarm QPE 100 Herbicide. Genfarm QPE 100 Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the amount of Genfarm QPE 100 directly to the spray tank and mix well. If tank mixing with carbendazim or mancozeb fungicides, add these to the spray tank and mix thoroughly BEFORE adding Genfarm QPE 100 Herbicide.

##### Use of a Surfactant/Wetting Agent

Genfarm QPE 100 Herbicide may be used with either of the following adjuvant system:

- 1) A non-ionic surfactant (1000 gai/L strength) at 100 mL/100L (i.e. 0.1% v/v) and a mineral spray oil at 1 L/100 L of final spray volume (i.e. 1% v/v).
  - 2) A non-ionic surfactant (1000 gai/L strength) at 200 mL/100L of final spray volume (i.e. 0.2% v/v).
- DO NOT use the non-ionic surfactant + mineral oil adjuvant system on vegetable crops, use a non-ionic surfactant only. DO NOT use a spray oil alone, a surfactant must always be added. DO NOT use non-methylated spray oils of crop origin as spray adjuvants.

### Compatibilities

Genfarm QPE 100 Herbicide is compatible with the insecticides deltamethrin, omethoate, dimethoate, methomyl, endosulfan, the fungicides carbendazim and mancozeb and the herbicides diflufenican and clopyralid **when using the surfactant only adjuvant system.**

Genfarm QPE 100 Herbicide is compatible with the herbicide sethoxydim and clethodim **when using the surfactant plus spray oil adjuvant system.**

When tank mixing Genfarm QPE 100 Herbicide and broadleaf herbicides, DO NOT use a spray oil, use a surfactant only. Application of Genfarm QPE 100 Herbicide /diflufenican tank mix may result in crop effects. The effect is temporary and usually appears as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected.

DO NOT add crop oil to tank mixtures of Genfarm QPE 100 Herbicide /diflufenican.

To minimise crop effects apply Genfarm QPE 100 Herbicide a minimum of 5 days prior to diflufenican application or a minimum 10 days after diflufenican application. (14-21 days if weeds have been stressed).

### Ground Spraying

Use sprayers with flat fan or hollow cone nozzles at 210-350 kPa to give an even distribution of droplets with no less than 50% of the spray volume in the 150-300 micron range. 110° flat fan nozzles are preferred.

Use between 50 and 100 litres of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150 litres prepared spray/ha.

For proper spray coverage adjust the boom and nozzle height according to the specification listed by the nozzle manufacturers. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained.

Avoid overlapping boomspray widths and shut off spray booms while starting, slowing or stopping.

### Sprayer cleanup

Thoroughly clean all traces of Genfarm QPE 100 Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops. If spray equipment has been previously used for the application of sulfonyleurea herbicides eg metsulfuron methyl, thifensulfuron methyl, chlorsulfuron, triasulfuron or terbutryn, ensure that the recommended sprayer cleanup procedure of the product previously sprayed has been followed before using the equipment for the application of Genfarm QPE 100 Herbicide.

### Precaution: Re-entry

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of Genfarm QPE 100 Herbicide. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store tank mixes of Genfarm QPE 100 Herbicide with other materials. When storing keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, relace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shields or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

### CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

**Genfarm Crop Protection Pty Ltd**  
**Suite 2, Level 3, 64 Talavera Road,**  
**Macquarie Park, NSW, 2113**  
**Tel: (02) 9889 5400**

<p><b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS Quizalofop-p-ethyl)</b></p>	
<p><b>UN NO. 3082 Packaging Group III Hazchem 2X</b></p>	
<p><b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b></p>	<p><b>FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 24 HOURS AUSTRALIA WIDE</b></p>

**THIS PAGE DOES NOT PRINT**

**Pages @ 100% on A3**