



**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **Sylon<sup>®</sup> 750 SG Herbicide**

**ACTIVE CONSTITUENT: 750g/kg CLOPYRALID**  
**present as the potassium salt**

**GROUP I HERBICIDE**

**For the control of a wide range of broadleaf weeds in wheat, barley, oats, triticale, canola, pastures and fallow land as specified in the Directions for Use table.**

**Important: Read the attached leaflet before using this product**

**Net Contents: 2kg / 3.5kg**

**Sipcam Pacific Australia Pty Ltd**  
**Level 1, 191 Malop Street**  
**GEELONG VIC 3220**

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

## SYLON 750 SG HERBICIDE

### STORAGE AND DISPOSAL

#### Storage for all containers

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store in area sheltered from rainfall. **DO NOT** store near food, feedstuffs, fertilisers or seed. This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SMALL SPILL MANAGEMENT

Sweep up material and contain in a refuse vessel for disposal in the same manner as for containers (see STORAGE AND DISPOSAL section).

### SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When mixing and loading wear cotton overalls, over normal clothing, buttoned to the neck and wrist and chemical resistant gloves. If applying by hand wear cotton overalls, or equivalent clothing, buttoned to the neck and wrist and chemical resistant gloves. Wash hands after use. After each days use, wash gloves and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone: Australia 13 11 26).

### MSDS

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier or from our web site, [www.sipcam.com.au](http://www.sipcam.com.au).

### NOTICE TO BUYERS

Sipcam Pacific Australia Pty Ltd (Sipcam) shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Sipcam's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk.

Batch No:

Date of Manufacture:



This product is NOT classified as a Dangerous Good under the Australian code for the transport of dangerous goods by road and rail.

For specialist advice in an emergency only, call 1800 033 111, all hours Australia wide

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **Sylon<sup>®</sup> 750 SG Herbicide**

**ACTIVE CONSTITUENT: 750g/kg CLOPYRALID**  
**present as the potassium salt**

<b>GROUP</b>	<b>I</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**For the control of a wide range of broadleaf weeds in wheat, barley, oats, triticale, canola, pastures and fallow land as specified in the Directions for Use table.**

**APVMA Approval Number: 62636/54041**

**Important: Read the attached leaflet before using this product.**

**Net Contents: 2kg / 3.5kg**

**Sipcam Pacific Australia Pty Ltd**  
**Level 1, 191 Malop Street**  
**GEELONG VIC 3220**

© Registered Trademark of Sipcam Pacific Australia Pty Ltd

## DIRECTIONS FOR USE

IT IS ESSENTIAL to select a rate appropriate for the weed size. Best results will be obtained when weeds are actively growing at treatment.

### Restraints:

**DO NOT** apply to weeds which may be stressed (inactive growth) due to prolonged periods of extreme heat or cold, moisture stress (water logging or drought) or previous herbicide treatment as reduced levels of control may result.

**DO NOT** apply immediately before sowing susceptible crops or sow susceptible crops into paddocks treated the previous year with Sylon 750 SG until after the required plantback period has elapsed - see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

**DO NOT** apply this product by air or mister within a Chemical Control Area in Victoria without a valid permit.

**DO NOT** spray if rain is likely within 3 hours.

**DO NOT** apply later than the 8 leaf stage of canola or the 1<sup>st</sup> node stage of winter cereals.

**DO NOT** compost material from treated plants or crops before reading the PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

Table 1. Winter Cereals and Canola: Pre-sowing Knockdown

WEED	WEED STAGE	RATE g/ha	CRITICAL COMMENTS
Capeweed, Chickpea (volunteer), Faba bean (volunteer), Vetch and Sub-clover	Up to 8 leaf and maximum 10 cm diameter	60 plus a knockdown herbicide	<b>Pre-sowing:</b> This rate should only be used in tank mixture with formulations of paraquat/diquat or glyphosate.

Table 2. Winter Cereals and Canola: Post-sowing Pre-emergence to 3 leaf crop stage

WEED	WEED STAGE	RATE g/ha	CRITICAL COMMENTS
Capeweed (In cereals only, WA only)	Pre-emergence to 8 leaf and maximum 10 cm diameter	60 plus diuron at 300 mL/ha	<b>Post sowing pre-emergent to 3 leaf:</b> This rate should only be used in tank mixture with diuron for control of transplants.  Rates of 120 – 200 g/ha give good suppression (reduced seed set and up to 80% weed control). 240 g/ha is required for good control of capeweed and sub-clover. Apply to moist soil and time treatment for major germination of weeds. Good soil moisture and application close to time of weed germination is essential for best control.
Capeweed, Faba bean (volunteer) and Sub-clover	Pre-emergence	120 – 240	

**Table 3. Winter Cereals: Early Post-emergence 2 leaf to 1<sup>st</sup> node crop stage**

WEED	WEED STAGE	RATE g/ha	CRITICAL COMMENTS
Capeweed (WA only)	Cotyledons to 6 leaf and maximum 5 cm diameter	60	<b>Early post-emergent:</b> Weeds should be young, actively growing and not larger than listed size. Weeds will become stunted and non- competitive soon after application, although final results may not show for some weeks.
Capeweed Soldier thistle St Barnaby's thistle	Up to 10 cm diameter (4 to 8 leaf)	120	
Chickpea Lentils and Safflower (volunteers)	Up to 6 leaf	100	
Faba bean and Lupins (volunteers)	Up to 4 leaf	100	Faba beans and lupins will usually survive, but will be stunted, uncompetitive and generally not set viable seed.
Field pea (volunteers)	Maximum 10 cm high or 6 nodes	60	
Medic and Lucerne seedling (volunteers)	Up to 8 leaf	60-80	For best control of hairy leaved medics such as snail medic, add 500 mL Uptake* Spraying Oil /100 L of water.
Sub-clover (volunteers)	Up to 6 leaf		
Vetch (volunteers)	Runners up to 10 cm and maximum 16 leaf	40	

**Table 4. Winter Cereals: Post-emergence tank mixtures WA, SA, VIC, TAS, NSW only (unless specified)**

Weeds should be young and actively growing. Weeds will become stunted and non-competitive soon after application although final results may not show for some weeks. Where a rate range is listed use low rate mixtures for small weeds to 5cm across and higher rate mixtures for weeds up to 10cm across. Use a surfactant such as BS-1000 for granular herbicides or the recommended adjuvant on the partner herbicide label.

WEED	WEED STAGE	RATE g/ ha	CRITICAL COMMENTS
Capeweed	Up to 4 leaf, 10cm diameter	80-120 plus 20 g/ha Chlorsulfuron	Chlorsulfuron mixes – 2 leaf to 1 <sup>st</sup> node crop stage.
		40 plus 5-7 g/ha Eclipse* + 0.35-0.5 L/ha MCPA LVE	Eclipse/MCPA LVE mixes – 3 leaf to 1 <sup>st</sup> node. Where 0.5 L/ha MCPA LVE added apply from 4-5 leaf to 1 <sup>st</sup> node crop stage.
		40 plus 5 g/ha Metsulfuron methyl + 0.5 L/ha MCPA LVE	Metsulfuron methyl/MCPA LVE mixes – 4 to 5 leaf to 1 <sup>st</sup> node crop stage.
		40 plus 0.75 L/ha Tigrex	Tigrex mixes – 3 leaf to 1 <sup>st</sup> node crop stage, but not on barley or kulin wheat in WA.
Field peas (volunteer)	Up to 6 node, 10cm diameter	40 plus 5-7 g/ha Eclipse + 0.5-0.7 L/ha bromoxynil / MCPA	bromoxynil/MCPA mixes – 3 leaf to 1 <sup>st</sup> node crop stage.
Vetch (volunteer)	Up to 4 branch, 10cm diameter	40 plus 5-7 g/ha Eclipse + 0.35-0.5 L/ha MCPA LVE	Eclipse/MCPA LVE mixes – 3 leaf to 1 <sup>st</sup> node. Where 0.5 L/ha MCPA LVE added apply from 4-5 leaf to 1 <sup>st</sup> node crop stage.
		40 plus 5g/ha Metsulfuron-methyl + 0.35L/ha MCPA LVE or 30 plus 0.7 L/ha MCPA LVE	Use 30 g/ha only in combination with MCPA LVE. Sylon + MCPA LVE mixes – 4 to 5 leaf to 1 <sup>st</sup> node crop stage.
Chickpea (volunteer)	Up to 4 branch, 10cm diameter	40 plus 5-7g/ha Eclipse + 0.5-0.7 L/ha bromoxynil / MCPA	bromoxynil/MCPA mixes – 3 leaf to 1 <sup>st</sup> node crop stage.
Faba bean (volunteer)	Up to 4 node, 10cm tall		
Lupin (volunteer)	Up to 6 leaf, 10cm tall	40 plus 5-7 g/ha Eclipse + 0.35-0.5 L/ha MCPA LVE	Eclipse/MCPA LVE mixes – 3 leaf to 1 <sup>st</sup> node. Where 0.5 L/ha MCPA LVE added apply from 4-5 leaf to 1 <sup>st</sup> node crop stage.
Sub-clover (volunteer)	Up to 5 trifoliolate, 5cm diameter	40 plus 5 g/ha Metsulfuron methyl + 0.35-0.7 L/ha MCPA LVE	Ally/MCPA LVE mixes – 4 to 5 leaf to 1 <sup>st</sup> node crop stage.
Prickly lettuce	Up to 6 leaf, max. 10cm diameter		
Medic (volunteer)	Up to 6 leaf, max. 5cm diameter		
Prickly lettuce	Up to 6 leaf, max. 10 cm diameter	60 plus 700 mL/ha MCPA LVE	Sylon 750 + MCPA LVE mixes – 4 to 5 leaf to 1 <sup>st</sup> node crop stage.
Thistles including: Nodding, Saffron Scotch, Slender Spear, Stemless, Variegated	Rosettes up to 10 cm max. diameter	20 plus 1 L/ha MCPA amine (500 g/L) or 20 + 700 mL/ha MCPA LVE	For thistle control, Sylon 750 SG rate will depend on density, growth stage, climatic conditions and time of application. Use higher rates for best control where high density and/or large weeds occur. MCPA or 2,4-D mixes apply from 4-5 leaf to 1 <sup>st</sup> node crop stage.
St Barnaby's Thistle	4 to 8 leaf, 5 to 10 cm across	20 - 40 + 2,4-D amine 0.5 – 1 L/ha or MCPA amine 1 – 1.5 L/ha	
Sowthistle (common) (WA, SA, Vic, Tas, NSW and QLD)	Young rosettes up to 8 true leaves	40 + 0.8 L/ha Torrent 242 or 5 g/ha metsulfuron (750g/kg) + 0.7 L/ha MCPA LVE	Apply to actively growing young rosettes. Use Uptake* Spraying Oil at 500 mL/100 L of water for improved control with Torrent 242 tank-mixes or BS-1000 with Metsulfuron /MCPA LVE tank-mixes. Apply tank-mixes from 4 - 5 leaf to 1 <sup>st</sup> node crop stage.
Skeleton weed (NSW, Vic and SA, WA only)	5 to 15 cm rosettes	200 plus 1 L/ha MCPA amine (500 g/L)	Weeds should be a minimum 5 cm in diameter, and growing actively. This rate will give control until harvest and substantially reduce weed numbers the following season. Apply from 4-5 leaf to 1 <sup>st</sup> node crop stage.

**Table 5. Canola Post-emergence 2 to 8 leaf crop stage.**

WEED	WEED STAGE	RATE g/ha	CRITICAL COMMENTS
Capeweed, Cotula, Saffron thistle, Skeleton weed, Soldier thistle	Up to 10 cm diam (4 to 8 leaf)	120	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks. Skeleton weed will only be controlled until harvest.
Chickpea, Lentils and Safflower (volunteer)	Up to 6 leaf	100	<b>For the control of annual grasses:</b> Sylon 750 SG is compatible with Inquest® Herbicide. Uptake Spraying Oil should be added to this tank-mix for best grass control. Sylon 750 SG + Inquest + Uptake Spraying Oil is compatible and selective to canola.
Faba beans and Lupins (volunteer)	Up to 4 leaf		
Field peas (volunteer)	Maximum 10 cm high or 6 nodes	60	Faba beans and lupins will usually survive, but will be stunted, uncompetitive and generally not set viable seed.
Medics and Lucerne seedlings (volunteer)	Up to 8 leaf		
Sub-clover (volunteer)	Up to 6 leaf	40	For best control of hairy leaved medics such as snail medic, add 500 mL Uptake Spraying Oil / 100 L water.  Will not control woolly pod vetch.
Vetch (volunteer)	Runners to 10 cm max. 16 leaf		
St Barnaby's thistle	4 to 8 leaf, 5 to 10 cm diameter	60-120	Sylon 750 SG rate will depend on weed density, growth stage, climatic conditions and time of application. Use higher rates for best control where high density and/or large weeds occur.

**Table 6. Herbicide Tolerant Canola: Post-emergence 2 to 8 leaf crop stage.**

WEED	WEED STAGE	RATE g/ha	CRITICAL COMMENTS
<b>Clearfield Canola</b>			
Cotula (common), Capeweed	Up to 6 leaf	60 + 40g OnDuty	Where capeweed is a significant component of the weed spectrum, a tank-mix with Sylon 750 SG Herbicide may be needed post-emergence. DO NOT exceed this rate as reduced control of grass weeds may occur.
<b>Triazine tolerant Canola</b>			
Capeweed, Lupins (volunteer), Saffron thistle, Skeleton weed, Soldier thistle and weeds from conventional canola	Up to 6 leaf	120	Sylon 750 SG is compatible with atrazine and simazine for use in triazine tolerant canola. Uptake Spraying Oil at 500 mL/100 L of water should be added to this mix for best grass and broadleaf weed control. For the control of annual grass weeds Sylon 750 SG + atrazine + Inquest + Uptake Spraying Oil are compatible and selective to triazine tolerant canola.

**Table 7. Pastures and Fallow Land – Post-emergence (Established perennial grass and sub-clover based pastures) (Boom spray application if not specified).**

WEED	WEED STAGE	RATE	STATE	CRITICAL COMMENTS
Hardhead thistle (creeping knapweed, Russian knapweed)	Actively growing plants	Hand gun: 200 g/100 L of water. Boom spray: 800 or 1600 g/ha	Vic & Qld only	See <b>CRITICAL COMMENTS</b> below for spraying thistles in pastures and fallow land.  Only use the 1600 g/ha rate in Qld by boom spray.
St Barnaby's thistle	5 to 8 leaf and 5 to 10 cm diameter	20 or 40 plus 0.5-1 L/ha 2,4- D amine or 1.5-2.5 L/ha 2,4- DB or 1 L/ha Inferno® or 1- 1.5 L/ha simazine + 1 L/ha 2,4-DB	NSW, Vic, Tas, SA and Qld only	
Thistles including: Nodding, Scotch, Spear, Slender Saffron, St Barnaby's, Variegated,	Rosette stage prior to stem elongation	20 or 28 g/ha plus 1 - 1.5 L/ha MCPA amine (500 g/L)/ ha Drench gun: 20 g/1 L of water Hand gun: 100 g/100 L of water	WA, NSW, Vic, Tas, SA and Qld only	
Nodding thistle	Rosettes up to 20 cm diameter	40	NSW only	Apply the spray from September to October. Apply by boom spray only. <b>DO NOT</b> apply to thistles over 20 cm in diameter. When thistles are over 20 cm in diameter use Sylon 750 SG plus MCPA (referred to above). <b>Clover Damage:</b> Damage to white clover will be no greater than damage with MCPA alone and less than damage from Sylon 750 SG plus MCPA mixtures. Damage to sub-clover may be greater than with MCPA or 2,4-D alone. <b>DO</b> <b>NOT</b> use for spot treatment.
Californian thistle	From early buds to flowering (December to February)	Hand gun: 100 g/100 L of water Boom spray: 800 g/ha	Vic and Tas only	Addition of a wetting agent at label rates is recommended. Re- treatment of regrowth in the year following treatment will usually be necessary to achieve a high level of control. <b>NOTE:</b> Clovers and medics will be eliminated for at least 1 year.
Lucerne	30 to 40cm high preflowering	120 plus 1.5 – 2 L/ha Glyphosate 450 + either 2 L/ha MCPA amine or 2 L/ha 2,4-D amine or 2 L/ha 2,4-D ester	Qld, NSW, Vic, SA, WA	Treat healthy, actively growing lucerne in early spring prior to flowering. After grazing or cutting, allow lucerne to regrow for approx. 4 weeks before treatment. For best control, do not re-graze for > 2 weeks after application. For complete control of lucerne in pasture, cultivate approx. 1 month after herbicide treatment.

**CRITICAL COMMENTS – Thistle control in pasture.**

- 1. Hardhead thistles - DO NOT USE HANDGUN APPLICATION ON LUCERNE, CLOVERS AND MEDICS AS THEY WILL BE ELIMINATED FOR AT LEAST ONE YEAR. Victoria only:** Use the lower rate only on light soils (sand and sandy loam) where a slightly lower degree of control is acceptable. Use the higher rate on all soil types where complete control is required. Addition of a wetting agent at label rates is recommended for treatment of hardhead thistle. Spray between September and April on actively growing plants for effective control. Thorough coverage is essential. Apply in 200 to 250 L of water/ha.



2. **BOOM SPRAYING:** Use the higher rates of Sylon 750 SG plus MCPA on multi-crowned plants or rosettes larger than 30 cm in diameter. Spraying may be done at any time during active growth, usually in early winter or spring. Avoid spraying during the dormant winter period or at any time when thistles are not actively growing. Do not spray flowering thistles.
3. **PRE-SPRAY MANAGEMENT:** The pasture should be slightly grazed prior to spraying to reduce clover and grass cover and expose the smaller thistles to the spray. The grazed pasture should be left seven days to allow thistles to freshen prior to treatment.
4. **POST-TREATMENT MANAGEMENT:** Response of thistles to treatment with the Sylon 750 SG plus MCPA mixture will be slow compared to the standard treatments with 2,4-D or MCPA. If possible delay grazing of sprayed thistles for 14 days after treatment.
5. **CLOVER DAMAGE:** Sylon 750 SG plus MCPA or 2,4-D mixtures can be damaging to clover. The low rate is no more damaging than label rates of 2,4-D or MCPA. Use 20g/ha mixes when clover is at the 6 trifoliate leaf stage to just prior to flowering. The 28 g/ha mix will reduce the clover component of the pasture for about two months. Use the 28 g/ha mix from 6 trifoliate leaf stage to flowering to minimise clover injury, and when clover has reached the 6 to 8 trifoliate leaf stage and where thistles are large due to early germination. Clover recovery will be quicker during periods of active growth. If clover damage is the major consideration, use the lower Sylon 750 SG rate to minimise damage.
6. **Gramoxone mixes are for lucerne pasture use only.** Simazine mixes are for silver grass control and for lucerne based pastures only.
7. **HANDGUN (Spot spray):** Treat from rosette stage to early flowering. Thorough spraying is necessary.
8. **DRENCHGUN:** Apply 10 mL to rosette crown. To multicrown plants, apply 10 mL to each crown.

**Table 8: Agricultural Non-crop Areas, commercial and Industrial Areas, Forests, Pastures and Rights-of-Way – Stem Injection Application on Acacia Species**

Mix 200g Sylon 750 SG with 2.5 litre of water and apply the diluted mix as directed below:

WEED GROWTH STAGE	APPLICATION RATE	CRITICAL COMMENTS
Single stems less than 25 cm diameter at base	1 mL of the diluted mix per cut @ 10 to 13 cm centres	Apply to waist high cuts. See GENERAL INSTRUCTIONS Application section for application method details. DO NOT exceed the recommended spacings from the centre of one cut to the centre of the next cut. Inject each stem of a multi stem tree where possible.
Multiple stems or more than 25 cm diameter at base	2 mL of the diluted mix per cut @ 10 to 13 cm centres	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

Situation	Withholding Period
PASTURES and FALLOW LAND	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION
CEREALS and CANOLA	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION
FORESTS	NOT REQUIRED when used as directed

#### GENERAL INSTRUCTIONS

##### MIXING

Measure the required quantity of granules by weighing on scales. Sylon 750 SG granules are highly soluble in water and will dissolve rapidly once added to fast moving water. Maintain agitation at all times, including during mixing as well as spraying.

##### Spray rigs with premix hoppers

For spray rigs that have a drop down chemical induction hopper, three-quarter fill this hopper with water and have the rinsing sprinkler operating. Add the Sylon 750 SG and when dissolved, transfer this batch into the quarter filled main tank. Continue to rinse the hopper until the entire product has washed through.

### **Spray rigs with limited bypass agitation**

For spray rigs that have limited bypass agitation, then as for most granulated formulations, pre-dissolve the Sylon 750 SG in a bucket before adding them to the main tank. Add Sylon 750 SG while stirring until the granules have dissolved.

### **Tank-mixes: The following order should be followed:**

1. **Quarter** fill the spray tank maintaining agitation
2. Add Sylon 750 SG granules, using the mixing procedure above.
3. Add Inquest **if it is to be used in the tank-mix.**
4. Add water to **Half** fill the spray tank.
5. Add wettable powders, water dispersible granules or suspension concentrates
6. Add other emulsifiable concentrates including other selective grass herbicides.
7. If Uptake\* Spraying Oil is to be used add this when spray tank is **Half** full.
8. If other adjuvants or a wetting agent is to be used then add these according to their label.
9. Add water to bring to the **final spray volume.**

Only mix sufficient spray solution for immediate use and avoid storing.

### **COMPATIBILITY**

**Conventional Canola:** Sylon 750 SG + Inquest + Uptake\* Spraying Oil are compatible and selective.

**Triazine Tolerant Canola:** Atrazine + Sylon 750 SG + Inquest + Uptake\* Spraying Oil are compatible and selective.

**Clearfield Canola:** OnDuty\* + Sylon 750 SG are compatible and selective.

Sylon 750 SG is compatible with the following:

**BROADLEAF HERBICIDES:** Acclaim®, Metsulfuron methyl, bromoxynil, bromoxynil/MCPA LVE, chlorsulfuron, diuron, glyphosate, MCPA amine, MCPA LVE, paraquat, Alarm®, terbutryn, 2,4-D amine, Broadstrike®, Eclipse®, Eclipse/MCPA LVE, Metsulfuron/MCPA LVE, Redeem®, atrazine, simazine, Torrent® 242, Tigrex\*

**GRASS HERBICIDES ON BROADLEAF CROPS:** Inquest Herbicide, Grasidim, OnDuty, atrazine, simazine

**GRASS HERBICIDES IN CEREAL CROPS:** Diclofop methyl, Density®, Wildcat®, Trend®, Tristar®.

Compatibilities for each herbicide and key grass weeds can be obtained from your Sipcam Representative.

**ADJUVANTS:** Uptake\* Spraying Oil, BS-1000

### **APPLICATION**

#### **BOOM SPRAYING CROP and PASTURES:**

Apply Sylon 750 SG in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivering 200 to 300 micron droplets and not less than 50 L/ha water volume for boom sprayers or not less than 20 L/ha for aerial applications.

Hardhead thistle - Use a spray volume of 200 to 250 L/ha of water.

#### **HIGH VOLUME HAND GUN:**

Apply the recommended mix to give full coverage of leaves and stems through a No. 6-8 tip at 700 to 1500 kPa. Spray volume for effective coverage of dense pasture weeds should be 10 to 15 litres of spray per 100 m<sup>2</sup> (10 m x 10 m) of infestation. For larger areas an equivalent would be 1000 to 1500 litres per infested hectare.

#### **STEM INJECTION**

To make a stem injection pocket at waist height, use a  $\frac{3}{4}$  length axe with a blade width of 5 to 7 cm. The axe cut must be through the bark and deep enough to place all the chemical in contact with the sap wood. The chemical must be applied immediately after the injection pocket is made. Apply chemical with a Phillips 5mL vaccinator fitted with a tree injector kit which can be accurately calibrated. Set vaccinator to deliver 1 mL of the diluted mix. When treating regrowth less than the width of the axe, ensure chemical does not run out the sides of the cut, as reduced control will result. This can be overcome by using the corner of the axe to make the pocket in the stem.

### CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

### PARTIAL CLEANING (before spraying crops that are selective to Sylon 750 SG):

After using Sylon 750 SG, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

### COMPLETE CLEANING (before spraying crops that are susceptible to Sylon 750 SG residues):

After using Sylon 750 SG, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate as above, then drain. Quarter fill the tank again and add an alkali detergent (e.g. Surf, Omo, Drive) at 500 mL/100 L water or 500g/100L water and circulate throughout the system for at least fifteen minutes. If using a concentrated laundry detergent use 250 g (or mL)/100 L water. Do not use chlorine based cleaners. Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using pressure hose and flush system with clean water. Chlorine based cleaners are NOT recommended.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and watercourses.

### RESISTANT WEEDS WARNING

#### GROUP I HERBICIDE

Sylon 750 SG Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide. Some naturally occurring weed biotypes resistant to the product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Sipcam representative.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do not apply under weather conditions, or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures.

**Composts and mulches** - Do not apply Sylon 750 SG Herbicide to crops or pastures that will be used for the production of compost or mulches or mushroom substrate to. Such compost or mulch made from plant material treated with Sylon 750 SG may cause damage to susceptible crops and plants.

**Susceptible crops** and plants include, but are not limited to chickpeas, clovers, cotton, faba beans, field peas, fruit trees, lentils, lupins, lucerne, medics, ornamentals, potatoes, safflower, tomatoes, vegetables, grape and kiwifruit vines, vetches, and wattles. Field peas, faba beans, lentils and vetches are particularly susceptible and should not be sown the season following an application of Sylon 750 SG at 200 g/ha.

*Where Sylon 750 SG Herbicide residue carryover from use rates of less than 200 g/ha is suspected and susceptible crops are to be planted, test the treated area as follows:*

*Field bioassay – where rain allows, plant a small area of the susceptible crop four to six weeks before desired planting date and take note of any symptoms of injury. If any herbicide symptoms are observed, only plant either canola or a cereal (see recommendation for northern and southern Australia below).*

*Pot bioassay – where not practical to do field bioassay, plant a small number of seeds of the susceptible crop into pots containing soil from the treated field. Do this four to six weeks before desired planting date. If any herbicide symptoms are observed, only plant either canola or a cereal (see recommendation for northern and southern Australia below).*

**Stubble from treated crops** - ensure that harvesters effectively spread crop straw and do not leave a heavy 'header trail' after harvest. Burn (if legal in the area), bale and remove, slash or incorporate stubble as soon as practical after harvest and as long as possible before planting next year to allow microbial breakdown of any residues in straw. Heavy stubble loads may carry more residues into the following season. Where heavy stubble burdens and/or non-wetting soils exist and less than

recommended rain amount have occurred from application to planting the susceptible crop (see below), only plant a winter and summer cereal or canola.

**Planting crops following use of Sylon 750 SG Herbicide in previous crop** - planting crops 'dry' without significant rain (see below) in the 'autumn break' of increases the risk of injury to susceptible crops. This practice should be avoided, or only plant a winter or irrigated summer cereal, or canola. In severely dry conditions, where less than 30% of average annual rainfall and/or less than the minimum rain has fallen between application and planting the next year, only plant a winter or summer cereal or canola.

#### **PLANTBACK PERIODS FOR SOUTHERN AUSTRALIAN WINTER DOMINANT RAINFALL AREAS (Sth NSW, VIC, SA, WA):**

**Required rainfall** - A minimum 25mm rain event in the post harvest summer to autumn period, with a subsequent extended period of at least 1 week where the top 10cm of the soil stays moist is required to enable breakdown of soil residues. Fastest residue breakdown will occur under good soil moisture and warm conditions, which promote microbial activity where significant rain (> 25mm) has fallen in summer to autumn, with soil wetting for at least one week, the following plantback periods apply:

Following Crops	Rate (g/ha) used previously	Plantback Interval
Clover, chickpea, faba bean, field pea, lentils, lupins, medics and vetch	Up to 120	9 months
	> 120 – 200	12 months
	> 200	24 months
Barley, canola, wheat, oats	All label rates	1 week

#### **PLANTBACK PERIODS FOR NORTHERN AUSTRALIA SUMMER DOMINANT RAINFALL AREAS (Nth NSW, QLD):**

Required rainfall before plantback:

**If planting susceptible summer crops** - at least 100 mm rain.

**If planting susceptible winter crops** - at least 150 mm rain.

This rain or irrigation should wet the soil for extended periods (at least one week).

This is essential for breakdown of soil residues prior to planting susceptible crops.

**If planting a cereal or canola crop** - at least 50 mm of rain or irrigation is required to enable soil wetting for at least one week.

Where these requirements have been met the following plantback periods apply:

Following Crops	Rate (g/ha) and Plantback Interval	
	Up to 30g/ha	>30 – 120g/ha
Chickpea, Cotton Soybean, Sunflower	3 months	6 months
Lucerne	9 months	9 months
Maize, sorghum	1 week	2 weeks
Wheat, barley, oats, canola	1 week	1 week

**Note:** Susceptible crops should not be sown for at least 2 years where Sylon 750 SG Herbicide at more than 120 g/ha has been used in northern Australia.

#### **PROTECTION OF LIVESTOCK**

**DO NOT** graze or cut treated crops for stock food except as specified under WITHHOLDING PERIODS.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Sylon 750 SG has low toxicity to fish, birds, honey bees, livestock, earthworms and aquatic organisms.

**DO NOT** contaminate streams, rivers or waterways with chemical or used containers.

## **STORAGE AND DISPOSAL**

### **Storage for all containers**

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store in area sheltered from rainfall. **DO NOT** store near food, feedstuffs, fertilisers or seed. This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **SMALL SPILL MANAGEMENT**

Sweep up material and contain in a refuse vessel for disposal in the same manner as for containers (see STORAGE AND DISPOSAL section).

### **SAFETY DIRECTIONS**

Will irritate the eyes. Avoid contact with eyes. When mixing and loading wear cotton overalls, over normal clothing, buttoned to the neck and wrist and chemical resistant gloves. If applying by hand wear cotton overalls, or equivalent clothing, buttoned to the neck and wrist and chemical resistant gloves. Wash hands after use. After each days use, wash gloves and contaminated clothing.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone: Australia 13 11 26).

### **MSDS**

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier or from our web site, [www.sipcam.com.au](http://www.sipcam.com.au).

### **NOTICE TO BUYERS**

Sipcam Pacific Australia Pty Limited (Sipcam) shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Sipcam's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk.

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

\* Non-Sipcam Trademark