

STORAGE AND DISPOSAL

Store in the closed, original container in a locked, well-ventilated area as cool as possible and away from children, animals, food, feedstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale vapour. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC, nitrile or neoprene gloves, goggles and disposable fume mask. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC, nitrile or neoprene gloves and goggles. If product is in eyes, wash it out immediately with water. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Ph Australia 13 11 26). If swallowed **DO NOT** induce vomiting. Give a glass of water.

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



IMTRADE CRACKER 250 EC FUNGICIDE

**ACTIVE CONSTITUENT: 250 g/L PROPICONAZOLE
SOLVENT: 640 g/L LIQUID HYDROCARBON**

GROUP 3 FUNGICIDE

For control of certain fungal diseases of Bananas, Oats, Peanuts, Perennial Ryegrass, Pineapples, Stone Fruit, Sugar Cane, Wheat and other crops in certain states as specified in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED
LEAFLET BEFORE OPENING OR
USING THIS PRODUCT.**

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799
Fax: 1800 171 788
Web: www.imtrade.com.au



MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet which is available from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITION OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgement in purchasing or using the same and every person dealing with this product does so at their own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

For Specialist Advice In An Emergency Dial
1800 033 111
24 Hours Australia Wide
IN A TRANSPORT EMERGENCY DIAL
000
POLICE OR FIRE BRIGADE

- CONTENTS 5L**
APVMA APPROVAL NO: 63820/5L/0909
- CONTENTS 10L**
APVMA APPROVAL NO: 63820/10L/0909



Batch No:

Date of Manufacture:



STORAGE AND DISPOSAL

Store in the closed, original container in a locked, well-ventilated area as cool as possible and away from children, animals, food, feedstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale vapour. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC, nitrile or neoprene gloves, goggles and disposable fume mask. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC, nitrile or neoprene gloves and goggles. If product is in eyes, wash it out immediately with water. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Ph Australia 13 11 26). If swallowed **DO NOT** induce vomiting. Give a glass of water.

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



IMTRADE CRACKER 250 EC FUNGICIDE

**ACTIVE CONSTITUENT: 250 g/L PROPICONAZOLE
SOLVENT: 640 g/L LIQUID HYDROCARBON**



GROUP 3 FUNGICIDE

For control of certain fungal diseases of **Bananas, Oats, Peanuts, Perennial Ryegrass, Pineapples, Stone Fruit, Sugar Cane, Wheat and other crops in certain states as specified in the Directions for Use table.**

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT.**

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au



MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet which is available from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITION OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgement in purchasing or using the same and every person dealing with this product does so at their own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

For Specialist Advice In An Emergency Dial

1800 033 111

24 Hours Australia Wide

IN A TRANSPORT EMERGENCY DIAL

000

POLICE OR FIRE BRIGADE

Batch No:

Date of Manufacture:

CONTENTS 20L
APVMA APPROVAL NO: 63820/20L/0909

CONTENTS 110L
APVMA APPROVAL NO: 63820/110L/0909

CONTENTS 200L
APVMA APPROVAL NO: 63820/200L/0909

CONTENTS 1000L
APVMA APPROVAL NO: 63820/1000L/0909



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE CRACKER 250 EC FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L PROPICONAZOLE
SOLVENT: 640 g/L LIQUID HYDROCARBON

GROUP 3 FUNGICIDE

For control of certain fungal diseases of Bananas, Oats, Peanuts, Perennial Ryegrass, Pineapples, Stone Fruit, Sugar Cane, Wheat and other crops in certain states as specified in the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET BEFORE
OPENING OR USING THIS PRODUCT**

APVMA Approval No: 63820/0909

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE:

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High volume		
Apricots	Prune Rust (<i>Tranzschelia discolor</i>)	SA only	-	<p>Dilute Spraying: 32 mL/100L</p> <p>Concentrate Spraying: Refer to the Mixing/ Application Section</p>	1 day	<p>Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.</p> <p>Curative Control: Apply when the disease first occurs. Further applications should be made when the disease occurs on new growth. DO NOT make more than 5 applications to any individual tree during the season.</p> <p>Protective Treatment: Spray mancozeb or zineb mixed with Imtrade Cracker 250 EC Fungicide at the full – recommended rates of application. This use is subject to a DMI anti-resistance strategy.</p>

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Bananas (including bananas inter-planted with avocados)	Leaf Spot (<i>Mycosphaerella musicola</i>), Leaf Speckle (<i>Mycosphaerella musae</i>), Cordana Leaf Spot (<i>Cordana johnstonii</i>)	Sth Qld, WA, NSW only	Ground Application: 200 to 400 mL + 3 to 5L of water miscible oil, in a convenient volume of water.	-	1 day	<p>This use is subject to a DMI anti-resistance strategy.</p> <p>Ground Application: Apply by misting machine or airblast sprayer. Use rates towards the higher end of the range where weather conditions favour diseases or where equipment or terrain does not permit thorough spray coverage of all foliage.</p> <p>This use is subject to a DMI anti-resistance strategy.</p> <p>NSW, Sth Qld: Ground and aerial application: Commence spraying at the start of the summer rainy season. Apply a maximum of 5 sprays per season at 21 to 28 day intervals. For effective control Imtrade Cracker 250 EC Fungicide must be applied at least 2 consecutive sprays at 21 to 28 day intervals before further treatments of an alternative recommended protectant fungicide are applied.</p>
	Leaf Spot (<i>Mycosphaerella musicola</i>), Leaf Speckle (<i>Mycosphaerella musae</i>), Cordana Leaf Spot (<i>Cordana musae</i>)	Nth Qld, WA, NT only	Aerial Application: 400 mL + 3 to 5L of a water miscible oil, in a minimum of 30L of water			
	Black Sigatoka (<i>Mycosphaerella fijiensis var difformis</i>)	Qld, WA, NT only	Aerial Application without water: 400 mL + 8 to 10L of spraying oil. (This use does not require further dilution with water).			

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Bananas (including bananas inter-planted with avocados)	<i>continued from previous page</i>					<i>continued from previous page</i> Nth Qld, WA, NT: Ground and aerial applications: Commence spraying at the start of the wet season and apply a maximum of 6 sprays per season at 14 to 21 day intervals. For effective control Imtrade Cracker 250 EC Fungicide must be applied for at least 2 consecutive sprays at 14 to 21 day intervals before further treatments of an alternative recommended fungicide are applied. Continue with treatments of an alternative recommended protective fungicide for the remainder of the season. Use the lower rate of oil in Nth Qld. DO NOT apply during July, August, September and October.
Barley	Powdery Mildew (<i>Blumeria graminis</i>)	All states	150 to 500 mL	-	Harvest 4 weeks Grazing 7 days	Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Powdery Mildew: Higher rates provide longer protection.
	Barley Scald (<i>Rhynchosporium secalis</i>)		500 mL			

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Barley (cont'd)	Spot Form Net Blotch (<i>Pyrenophora teres f. maculate</i>)	All states	500 mL	-	Harvest 4 weeks Grazing 7 days	Apply after flag leaf is around 70% emerged and before infection averages 10% on the flag-2 leaf. Ensure thorough coverage of stems and leaves.
Boronia	Rust (<i>Puccinia boroniae</i>)	Tas, WA only	500 mL to 1L	-	-	Apply 2 to 5 applications at 10 to 14 days intervals during the main disease period. Use the lower rate when application is made protectively before disease occurs. Use the higher rate when the disease is first observed and when the minimum number of applications are applied.
Oats	Stem Rust (<i>Puccinia graminis f. sp avenae</i>)	All states	500 mL	-	Harvest 4 weeks Grazing 7 days	Apply at the first sign of disease and before there is an average of over 2 pustules per tiller. Ensure thorough coverage of stems and leaves.

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Oats (cont'd)	Crown Rust (<i>Puccinia coronata f. sp avenae</i>)	All states	250 to 500 mL	-	Harvest 4 weeks Grazing 7 days	Apply after flag blade leaf is fully emerged or Z39 and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5-10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Suppression of Septoria Leaf Blotch (<i>Leptosphaeria avenaria</i>)				Apply after flag blade is fully emerged or Z39. If infection averages 10% on the flag-2 leaf. The high rate of application gives longer period of protection than the lower rates. Use higher rates on high potential crops when conditions favour severe disease development during flowering. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.	

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Peanuts	Early Leaf Spot (<i>Cercospora arachidicola</i>), Late Leaf Spot (<i>Cercosporidium personatum</i>)	Sth Qld, NSW, WA only	400 to 600 mL	-	14 days	This use is subject to a DMI anti-resistance strategy. Spray when disease symptoms are first observed. Apply at 14 days intervals while weather conditions favour disease. Use rates towards the higher end of the range when wet conditions prevail. Use a fungicide from a different activity group (non-DMI) after 3 consecutive sprays using Imtrade Cracker 250 EC Fungicide alone. Apply a maximum of 5 sprays per season. The leaves of peanuts sprayed may become darker green in colour and modified in shape. These effects will not adversely affect yield at recommended rates.
	Rust (<i>Puccinia arachidis</i>)	Sth QLD, WA only	600 mL			

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Peppermint, Spearmint grown for oil production only	Mint Rust (<i>Puccinia menthae</i>)	NSW, Vic, Tas only	500 mL	-	5 weeks	Apply 2 to 5 applications at 10 to 14 days intervals during the main disease period. DO NOT use on mint grown for the fresh market.
Perennial Ryegrass	Stem Rust (<i>Puccinia graminis</i>), Blind Seed Disease (<i>Gleotinia granigena</i>)	Vic only			4 weeks	Apply at ear emergence and again at anthesis.
Plums for Prune production	Prune Rust (<i>Tranzschelia discolor</i>)	NSW, Vic, SA, WA only	-	Dilute Spraying: 32 mL/100L Concentrate Spraying: Refer to the Mixing/ Application section	1 day	Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying method. Curative Control: Apply when the disease first occurs. Further applications should be made when the disease occurs on new growth. DO NOT make more than 5 applications to any individual tree during the season. Protective Treatment: Spray mancozeb or zineb mixed with Imtrade Cracker 250 EC Fungicide at the full recommended rates of application. This use is subject to a DMI anti-resistance strategy.

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Pineapples	Base Rot (<i>Thielaviopsis paradoxa</i>)	Qld, NT, WA only	-	10 to 20 mL/100L	-	Preplant dip: Ensure thorough coverage by totally immersing the planting material in the dip solution. Allow 50 mL of the dip solution per plant. Apply the higher rate under conditions of high disease pressure.
Poppies <i>Papaver somniferum</i>	Leaf Smut (<i>Entyloma fuscum</i>)	TAS only	500 mL	-	4 weeks	Usage recommended by poppy contract companies. Apply as mid season application in the full flower/petal drop period when disease is present.
Stone Fruit	Brown Rot (Blossom Blight) (Blossom phase) (<i>Monilinia laxa</i>)	Vic, Tas, WA only	-	Dilute spraying: 25 mL/100L Concentrate Spraying: Refer to the Mixing /Application section	1 day	Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. This use is subject to a DMI anti-resistance strategy. Apply at early (1 to 10%) blossom and again at full bloom. A further application is made at shuck-fall. Only 2 consecutive applications of DMI fungicides can be made during this period.
	Brown Rot (Blossom phase) (<i>Monilinia fruticola</i>)	Qld, NSW, Tas, SA, WA only				

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Stone Fruit (<i>cont'd</i>)	Brown Rot (Fruit phase)	Qld, NSW Vic, Tas, SA, WA only	-	<p>Dilute spraying: 25 mL/100L</p> <p>Concentrate Spraying: Refer to the Mixing /Application section</p>	1 day	<p>Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Apply 3 weeks and 1 week before harvest.</p> <p>Only 2 consecutive applications of DMI fungicides can be made during this period. The last Blossom Blight spray and the first Brown Rot (fruit phase) spray should be regarded as consecutive applications.</p> <p>For varieties with extended harvesting periods, a third spray during the picking period may be applied if conditions are favourable for disease development.</p>
Sugar Cane	Pineapple Disease (<i>Ceratocystis paradoxa</i>)	Qld, NSW, WA only	-	20 mL/100L	-	Ensure thorough coverage of the cut ends of sugar cane setts.

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	Qld, NSW, Vic, SA, WA only	250 or 500 mL	-	28 days	Spray between joining and end of flowering when 10 to 20% of leaves are infected. A repeat spray 21 to 28 days later may be required. Use higher rate under high infection pressure or where longer residual protection is required.
		Tas only				Spray when 10 to 20% of leaves are infected. A repeat spray 21 to 28 days later may be required. Use higher rate under high infection pressure or when longer residual protection is required.
	Powdery Mildew (<i>Blumeria graminis</i>)	All states	150 to 500 mL	Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.		

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Wheat (cont'd)	Stem Rust (<i>Puccinia graminis</i>)	All states	500 mL	-	28 days	Apply at the first sign of the disease and before there are more than 2 pustules per tiller. Ensure thorough coverage of stems and leaves.
	Septoria Nodorum Blotch (<i>Phaeosphaeria nodorum</i>)		150 to 500 mL			Apply after flag leaves are around 70% emerged if infection averages 10% on the flag-2 leaf. The high rate of application gives a longer period of protection than the lower rates. Use higher rates on higher potential crops when conditions favour severe disease development during flowering. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
			Per Hectare	High Volume		
Wheat (cont'd)	White Leaf Rust (<i>Puccinia triticina</i>)	All States	150 to 500 mL	-	28 days	Apply after flag leaf is 70% emerged and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5 to 10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Yellow Spot (<i>Pyrenophora tritici-repentis</i>)		250 to 500 mL			Apply after flag leaves are around 70% emerged if infection averages 10% on the flag-2 leaf. Higher rates provide longer protection. Use higher rates on higher potential crops when conditions favour severe disease development. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Septoria Tritici Blotch (<i>Mycosphaerella graminicola</i>)					

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

PERENNIAL RYEGRASS:	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION
PEPPERMINT, SPEARMINT:	DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION
POPPIES:	DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION
BARLEY, OATS, WHEAT:	DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION
PEANUTS:	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION
APRICOTS, BANANAS, STONE FRUIT:	DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION
PINEAPPLES, SUGAR CANE:	DO NOT HARVEST FOR 1 DAY AFTER APPLICATION
	WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

DO NOT apply more than 6 consecutive sprays per season.

NOTE: For cereals the flag 2 leaf is the 3rd last fully emerged leaf i.e. the second leaf below flag.

Wheat Stripe Rust susceptible varieties apply when 10% leaves infected.

Wheat Stripe Rust moderately susceptible varieties apply when 15 to 20% leaves infected.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

Imtrade Cracker 250 EC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Imtrade Cracker 250 EC Fungicide is a Group 3 Fungicide. Some naturally occurring individual fungi resistant to Imtrade Cracker 250 EC Fungicide and other Group 3 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi populations if these fungicides are used repeatedly. These resistant fungi will not be controlled by Imtrade Cracker 250 EC Fungicide and other Group 3 Fungicides thus resulting in a reduction in efficacy and possible

yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Cracker 250 EC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

MIXING

Shake well before use. Fill the spray tank and then add concentrate. Mix well.

Pineapples Pre-Plant Dip

Add the required amount of Imtrade Cracker 250 EC Fungicide directly to the dip and mix well. Avoid excessive contamination of the dip with the organic matter.

APPLICATION

Imtrade Cracker 250 EC Fungicide may be applied by ground rig, high or low volume, or by air.

Cereals: May be applied by boom spray or aircraft. Ensure complete coverage of all leaves and stems are obtained. The objective of spraying is to control disease on the upper 2 to 3 leaves during grain filling.

With aircraft, as a guide, apply to 10 to 20L/ha with the lower rate being used when applications are made with a cross wind of not less than 5 knots. Use the higher rate when applying to dense crops.

Apricots, Plums and other Stone Fruits: Apply by high volume (dilute) sprayer or concentrate sprayer.

Bananas: Apply by misting machine, air blast sprayer or aircraft. Use a minimum of 30L of water if applying by air.

Dilute Spraying

Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Direction for Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set-up and operation may also need to be changed as the crop grows.

Concentrate Spraying

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute Spraying above) for the

crop canopy. This is needed to calculate the concentrate-mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only

1. Dilute spray volume as determined above: for example 1500L/ha.
2. Your chosen concentrate spray volume: for example 500L/ha.
3. The concentration factor in this example is: $3 \times$ (i.e. 1500L divided by 500L = 3).
4. If the dilute label rate is 10 mL/100L, then the concentrate rate becomes 3×10 that is 30 mL/100L of concentrate spray.

The chosen spray volume amount of product per 100L of water and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying uses are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

COMPATIBILITY

Imtrade Cracker 250 EC Fungicide may be mixed with any one of the registered products containing azinphos-methyl, chlorothalonil, demeton-s-methyl, diazinon, methomyl liquid, metalaxyl, propargite, parathion, dimethoate, copper oxychloride, mancozeb and zineb.

PRECAUTION

Re-entry Period

DO NOT enter treated area until spray has dried.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

Low hazard to bees. No special precautions are required.

STORAGE AND DISPOSAL

Store in the closed, original container in a locked, well-ventilated area as cool as possible and away from children, animals, food, feedstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale vapour. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC, nitrile or neoprene gloves, goggles and disposable fume mask. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC, nitrile or neoprene gloves and goggles. If product is in eyes, wash it out immediately with water. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Ph Australia 13 11 26). If swallowed **DO NOT** induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet which is available from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITION OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgement in purchasing or using the same and every person dealing with this product does so at their own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

<p>For Specialist Advice In An Emergency Dial 1800 033 111 24 Hours Australia Wide</p>
<p>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</p>

APVMA Approval No: 63820/0909