

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Flufol



IN FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

*For the control of certain fungal diseases
on wheat, barley and canola when mixed
with fertiliser or applied as a foliar spray
as per the Directions for Use Table.*

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E, 75-85 O'Riordan Street, Alexandria NSW 2015
T: 02 8338 9473 F: 02 9313 4352 W: www.farmalinx.com

CONTENTS
5 Litres

FARMALINX

Flufol

IN FURROW & FOLIAR FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Flufol In Furrow & Foliar Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

APVMA Approval No.: 64291/1-5L/0809

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



IN A TRANSPORT
EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

SPECIALIST ADVICE IN AN EMERGENCY ONLY

1800 024 973

ALL HOURS – AUSTRALIA WIDE

BARCODE

Batch No.

Date of Manufacture

farmalinx

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Flufol



IN FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E, 75-85 O'Riordan Street, Alexandria NSW 2015
T: 02 8338 9473 F: 02 9313 4352 W: www.farmalinx.com

CONTENTS
20 Litres



FARMALINX

Flufol

IN FURROW & FOLIAR FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

For Non-Refillable Containers: Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs FARMALINX Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Flufol In Furrow & Foliar Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

APVMA Approval No.: 64291/10-1000L/0809

	IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 024 973 ALL HOURS – AUSTRALIA WIDE

BARCODE

Batch No.
Date of Manufacture



FARMALINX

Flufol

IN FURROW & FOLIAR FUNGICIDE

DIRECTIONS FOR USE:

A. FARMALINX FLUFOL APPLIED IN-FURROW AS A FERTILISER TREATMENT

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS																												
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400 mL	4 weeks grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 400 mL of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333																
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																															
60	667*																															
70	571																															
80	500																															
100	400																															
120	333																															
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	200 or 400 mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> <th>200 mL rate</th> <th>400 mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td><td></td></tr> <tr><td>60</td><td>333</td><td>667*</td><td></td></tr> <tr><td>70</td><td>286</td><td>571</td><td></td></tr> <tr><td>80</td><td>250#</td><td>500</td><td></td></tr> <tr><td>100</td><td>200#</td><td>400</td><td></td></tr> <tr><td>120</td><td>167#</td><td>333</td><td></td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	200 mL rate	400 mL rate	50	400	800*		60	333	667*		70	286	571		80	250#	500		100	200#	400		120	167#	333	
	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	200 mL rate	400 mL rate																												
50	400	800*																														
60	333	667*																														
70	286	571																														
80	250#	500																														
100	200#	400																														
120	167#	333																														
	Septoria Tritici Blotch (<i>S. tritici</i>), Take-all (<i>Gaeumannomyces graminis</i>)	400 mL																														
Barley	Powdery Mildew (<i>Erysiphe graminis</i>), Scald (<i>Rhynchosporium secalis</i>)	200 or 400 mL		* Application volumes over 600 mL per 100 kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. # Where application volume is low, product may be diluted with water to improve coverage.																												

B. FARMALINX FLUFOL APPLIED AS A FOLIAR SPRAY

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	250-500 mL Ground Application Apply in 100-150 L water plus 200 mL FARMALINX WetDrop per	7 weeks Harvest or Grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf Rust (<i>P. recondita</i>), Septoria Nodorum Blotch (<i>Septoria nodorum</i>), Septoria Tritici Blotch (<i>S. tritici</i>)	100 L of spray solution. Aerial Application Apply in a minimum of 15 L water plus 200 mL FARMALINX WetDrop per		Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
Barley	Powdery Mildew (<i>Erysiphe graminis</i>)	100 L of spray solution.	10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3-4 weeks apart at the low rate, may be required.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

APPLIED AS A FOLIAR SPRAY

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY – 10 WEEKS

WHEAT – 7 WEEKS

DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY – 10 WEEKS

WHEAT – 7 WEEKS

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

FARMALINX Flufol In-Furrow & Foliar Fungicide is a member of the DMI group of fungicides. For fungicide resistance management FARMALINX Flufol In-Furrow & Foliar Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled this product and other Group 3 fungicides thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

GROUP 3 FUNGICIDE

APPLICATION

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Flufol In Furrow & Foliar Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245

South Sydney Corporate Park, Unit W2E,

75-85 O'Riordan Street, Alexandria NSW 2015

T: 02 8338 9473 F: 02 9313 4352

Website: www.farmalinx.com

APVMA Approval No.: 64291/0809

Foliar Application

Check the drop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

The method of disposal of the container depends on the container type. Read the storage and disposal instructions on the label that is attached to the container.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Flufol

IN FURROW & FOLIAR FUNGICIDE
ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E,
75-85 O'Riordan Street, Alexandria NSW 2015

farmalinx