

# POISON

**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

FARMALINX

# Quiz



## HERBICIDE

**ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL**  
**SOLVENT: 802 g/L HYDROCARBON LIQUID**

**GROUP A HERBICIDE**

*For the control of certain grass  
weeds in broadleaf crops  
as per Directions for Use table.*

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING**

### STORAGE AND DISPOSAL

DO NOT store tank-mixes of FARMALINX Quiz Herbicide with other materials. When storing, keep from contact with fertilisers, fungicide and seeds. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

**For Non-Refillable Containers:** Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For Refillable Containers:** Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs FARMALINX Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do not induce vomiting. Give a glass of water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Quiz Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

APVMA Approval No.: 64307/5-1000/0809

	IN A TRANSPORT EMERGENCY <b>DIAL 000</b> POLICE OR FIRE BRIGADE
	SPECIALIST ADVICE IN AN EMERGENCY ONLY <b>1800 024 973</b> ALL HOURS – AUSTRALIA WIDE

BARCODE	Batch No.
	Date of Manufacture

## farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245  
South Sydney Corporate Park, Unit W2E, 75-85 O'Riordan Street, Alexandria NSW 2015  
T: 02 8338 9473 F: 02 9313 4352 W: www.farmalinx.com

CONTENTS  
**20 Litres**

## farmalinx

FARMALINX

**Quiz**

HERBICIDE

**DIRECTIONS FOR USE:**

**RESTRAINTS:** DO NOT apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).  
DO NOT apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing.  
Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.  
DO NOT apply under conditions of high temperature and low humidity.  
DO NOT apply if rainfall is expected within 3 hours.  
DO NOT apply less than 50 L prepared spray/ha.

**BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES**

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Canola Carrots Cauliflower Chickpeas Clover & Sub Clover Pastures Cucumbers• Faba Beans Field Peas Honey Dew Melons• Lentils Lucerne Lupins Medic Pasture Mung Beans Navy Beans Onions Potatoes Pumpkins• Radish Tomatoes Vetch	Annual Ryegrass ( <i>Lolium rigidum</i> ), Brome Grasses ( <i>Bromus</i> spp.)	3 leaf to early tillering	300 mL or 375 mL	14 days 9 weeks 11 weeks 10 weeks 14 days 12 weeks	— 28 days — — — 28 days 14 days	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.*
	Barley Grass ( <i>Hordeum</i> spp.), Volunteer Barley, Volunteer Wheat	3 leaf to early tillering	250 mL	14 days 12 weeks 9 weeks 28 days	— — 28 days 28 days	Apply when weeds are actively growing.*
	Wild Oats ( <i>Avena</i> spp.)	3 leaf to mid-tillering	125 mL or 250 mL	9 weeks 12 weeks — 6 weeks — 12 weeks 12 weeks 18 weeks 10 weeks 9 weeks 21 days 4 weeks 12 weeks	— 28 days 28 days 14 days 28 days — — — — — — — 28 days	Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control.*

**\*ALWAYS ADD A SURFACTANT/WETTING AGENT AT THE RECOMMENDED RATE – SEE SECTION ON "USE OF SURFACTANT/WETTING AGENT" FOR DETAILS.**  
**• IMPORTANT: DO NOT APPLY FARMALINX QUIZ HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.**

**BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES**

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Carrots Cucumbers• Green Beans Honey Dew Melons• Lucerne Mung Beans Navy Beans Onions Peanuts Potatoes Pineapples Pumpkins• Radish Soybeans Sunflower Tomatoes	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ), Barnyard Grass ( <i>Echinochloa crusgalli</i> )	3 leaf to early tillering	500 mL or 750 mL	14 days 9 weeks 14 weeks 10 weeks 14 days	— — — — — 28 days 28 days 28 days	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.*
	Couch Grass ( <i>Cynodon dactylon</i> )	2-4 leaf seedling	500 mL	5 weeks 9 weeks — 12 weeks	28 days 28 days 28 days	Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed.*
	Dinebra ( <i>Dinebra retroflexa</i> )	3 leaf to early tillering	250 mL or 375 mL	12 weeks 18 weeks 11 weeks 10 weeks 7 days 9 weeks 21 days	— 11 weeks — — — — 28 days	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.*
	Crowsfoot Grass ( <i>Eleusine indica</i> )	2 leaf to early tillering	500 mL	9 weeks 21 days	— — 28 days	Apply when weeds are actively growing.*
	Foxtail Millet ( <i>Setaria italica</i> )	2 leaf to early tillering	250 mL or 500 mL	12 weeks 9 weeks 4 weeks	28 days 11 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.*
	Columbus Grass ( <i>Sorghum x alnum</i> ), Johnson Grass ( <i>Sorghum halepense</i> ), Liverseed Grass ( <i>Urochloa panicoides</i> )	2 leaf to 3 tiller	500 mL or 750 mL	—	—	—

**\*ALWAYS ADD A SURFACTANT/WETTING AGENT AT THE RECOMMENDED RATE – SEE SECTION ON "USE OF SURFACTANT/WETTING AGENT" FOR DETAILS.**  
**• IMPORTANT: DO NOT APPLY FARMALINX QUIZ HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.**

**BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES**

**BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES**

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

**PRECAUTION**

**SPRAYER CLEANUP**

**GROUND SPRAYING**

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**STORAGE AND DISPOSAL**

**COMPATIBILITIES**

**USE OF A SURFACTANT/WETTING AGENT**

**RESISTANT WEEDS WARNING**

**GENERAL INSTRUCTIONS**

**WITHOLDING PERIODS:**

**GRAZING:**

**SAFETY DIRECTIONS**

**FIRST AID**

**MATERIAL SAFETY DATA SHEET**

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

\* Other trademarks

**farmalinx**

FARMALINX Pty Ltd ABN 95 134 353 245

South Sydney Corporate Park, Unit W2E,

75-85 O'Riordan Street, Alexandria NSW 2015

T: 02 8338 9473 F: 02 9313 4352

Website: www.farmalinx.com

APVMA Approval No.: 64307/0809

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**Quiz**  
FARMALINX

HERBICIDE

ACTIVE CONSTITUENT: 99.5 g/L QUIZALOFOP-P-ETHYL  
SOLVENT: 802 g/L HYDROCARBON LIQUID

GROUP A HERBICIDE

For the control of certain grass weeds in broadleaf crops  
as per Directions for Use table.IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE  
CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCTFARMALINX Pty Ltd ABN 95 134 353 245  
South Sydney Corporate Park, Unit W2E,  
75-85 O'Riordan Street, Alexandria NSW 2015**farmalinx****BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES - continued**

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Carrots Cucumbers• Green Beans Honey Dew Melons• Lucerne Mung Beans Navy Beans Onions Peanuts Potatoes Pineapples Pumpkins• Soybeans Sunflower Tomatoes	Rhodes Grass ( <i>Chloris gayana</i> ), Stink Grass ( <i>Eragrostis cilianensis</i> )	3 leaf to early tillering	250 mL or 500 mL	14 days 9 weeks 14 weeks 10 weeks 14 days 5 weeks 9 weeks	— — — — 28 days — —	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.*
	Queenland Blue Grass ( <i>Dichanthium sericeum</i> )	3 leaf to early tillering	500 mL	14 days 5 weeks 9 weeks	— 28 days —	Apply when weeds are actively growing.*
	Summer Grass ( <i>Digitaria ciliaris</i> )	3 leaf to early tillering	500 mL or 1000 mL	— 12 weeks 12 weeks 18 weeks 11 weeks 10 weeks 7 days 9 weeks 21 days 12 weeks 9 weeks 4 weeks	28 days 28 days 28 days 11 weeks — — — — 28 days 11 weeks —	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.*

**\*ALWAYS ADD A SURFACTANT/WETTING AGENT AT THE RECOMMENDED RATE – SEE SECTION ON "USE OF SURFACTANT/WETTING AGENT" FOR DETAILS.**  
**• IMPORTANT: DO NOT APPLY FARMALINX QUIZ HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.**

**HAND GUN SPRAYING: ALL STATES**

CROP	WEEDS CONTROLLED	WEEDS STAGE	RATE/L	WITHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Carrots Cucumbers• Faba Beans Field Peas Grapes Green Beans Honey Dew Melons• Lentils Lucerne Lupins Medic Pasture Mung Beans Navy Beans Non crop Onions Peanuts Pineapples Potatoes Pumpkins• Radish Soybeans Sunflowers Tomatoes Vetch	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ), Crowsfoot Grass ( <i>Eleusine indica</i> )	3 leaf to early tillering	125 mL	14 days 9 weeks 14 days 11 weeks 10 weeks 12 weeks	— — 28 days — — 28 days 14 days	Apply when weeds are actively growing. Use a minimum of 800 L of prepared spray/ha. Thoroughly wet target weeds (especially well established clumps) without causing run-off. Repeat spraying may be necessary for perennial grass species.
	Paspalum ( <i>Paspalum dilatatum</i> )	Seedling to established clumps	—	14 days 12 weeks 9 weeks	— 28 days 28 days	Always add a surfactant/wetting agent at the recommended rate – see section on "Use of Surfactant/Wetting agent" for details.
	Johnson Grass ( <i>Sorghum halepense</i> )	Seedling 2 leaf to early tillering	—	12 weeks 9 weeks	28 days 28 days	—
	Rhizome 2 leaf to early tillering	—	250 mL	5 weeks 9 weeks 12 weeks	28 days 28 days 28 days	—
	Kikuyu ( <i>Pennisetum landestinum</i> )	Seedling to established plants	—	6 weeks 12 weeks 12 weeks	28 days 28 days 28 days	—

**• IMPORTANT DO NOT APPLY FARMALINX QUIZ HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.**

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

www.farmalinx.com

CONTINUED OVERLEAF

**COMPATIBILITIES**

**USE OF A SURFACTANT/WETTING AGENT**

**RESISTANT WEEDS WARNING**

**GENERAL INSTRUCTIONS**

**WITHOLDING PERIODS:**

**GRAZING:**

**SAFETY DIRECTIONS**

**FIRST AID**

**MATERIAL SAFETY DATA SHEET**

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

\* Other trademarks

RECEIVED

By vscarlet at 2:05 pm, Aug 27, 2009