

150 mm

140 mm

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**★ RYGEL**

# RYGREX

## Selective Herbicide

**ACTIVE CONSTITUENTS:** 250 g/L MCPA present as the ethyl hexyl ester,  
25 g/L DIFLUFENICAN  
**SOLVENTS:** 325 g/L LIQUID HYDROCARBONS,  
150 g/L N-METHYL-2-PYRROLIDONE

GROUP **FI** HERBICIDE

For the control of certain broadleaf weeds in winter cereals  
and clover as specified in Directions for Use Table..

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING**  
APVMA Approval No: 64747/5L/0310



**5 LITRES**

**★ RYGEL**  
AUSTRALIA PTY LTD

194 mm

170 mm

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



★ **RYGEL**

# RYGREX

## Selective Herbicide

**ACTIVE CONSTITUENTS:** 250 g/L MCPA present as the ethyl hexyl ester,  
25 g/L DIFLUFENICAN  
**SOLVENTS:** 325 g/L LIQUID HYDROCARBONS,  
150 g/L N-METHYL-2-PYRROLIDONE

GROUP **FI** HERBICIDE

For the control of broadleaf weeds in winter cereals and clover as specified in Directions for Use Table.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING**

APVMA Approval No: 64747/20L/0310



# 20 LITRES

★ **RYGEL**  
AUSTRALIA PTY LTD

194 mm

170 mm

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



★ **RYGEL**

# RYGREX

## Selective Herbicide

**ACTIVE CONSTITUENTS:** 250 g/L MCPA present as the ethyl hexyl ester,  
25 g/L DIFLUFENICAN  
**SOLVENTS:** 325 g/L LIQUID HYDROCARBONS,  
150 g/L N-METHYL-2-PYRROLIDONE

GROUP **FI** HERBICIDE

For the control of broadleaf weeds in winter cereals and clover as specified in Directions for Use Table.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING**

APVMA Approval No: 64747/100L/0310



# 100 LITRES

★ **RYGEL**  
AUSTRALIA PTY LTD

194 mm

170 mm

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



★ **RYGEL**

# RYGREX

## Selective Herbicide

ACTIVE CONSTITUENTS: 250 g/L MCPA present as the ethyl hexyl ester,  
25 g/L DIFLUFENICAN  
SOLVENTS: 325 g/L LIQUID HYDROCARBONS,  
150 g/L N-METHYL-2-PYRROLIDONE

GROUP **FI** HERBICIDE

For the control of broadleaf weeds in winter cereals and clover as specified in Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING

APVMA Approval No: 64747/110L/0310



# 110 LITRES

★ **RYGEL**  
AUSTRALIA PTY LTD

194 mm

170 mm

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



★ **RYGEL**

# RYGREX

## Selective Herbicide

ACTIVE CONSTITUENTS: 250 g/L MCPA present as the ethyl hexyl ester,  
25 g/L DIFLUFENICAN  
SOLVENTS: 325 g/L LIQUID HYDROCARBONS,  
150 g/L N-METHYL-2-PYRROLIDONE

GROUP **FI** HERBICIDE

For the control of broadleaf weeds in winter cereals and clover as specified in Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING

APVMA Approval No: 64747/1000L/0310



**1000 LITRES**

★ **RYGEL**  
AUSTRALIA PTY LTD

130 mm

140 mm

**Rygel Rygrex Selective Herbicide**

**DIRECTIONS FOR USE**

For complete directions for use please read attached booklet.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

**5L, 20L containers only**

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and setup for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

**Re-Fillable Containers**

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured.

Empty product as required into application equipment. Do not attempted to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that equipment used in transfer of the product is disconnected, triple rinsed and drained after each use.

When the container is empty, close all caps and valves and return the container to the point of purchase.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and

**PEEL HERE** ▶

before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of Rygel Rygrex Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Rygel Australia Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Rygel Australia Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

APVMA Approval No: 64747/0310

Batch No: } see container

DOM:

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**



**Rygel Australia Pty Ltd**  
ACN: 106 839 007  
103 Ordish Rd, Dandenong South, Vic 3175  
Tel: 03 9768 2803 Fax: 03 9768 2804

Top layer - rear label (sticks to bottom layer)

110mm 600mm  
Folded to 110mm x 100mm  
Page 1  
Section 1

**WEED LIST**

<b>WEED</b>	<b>Common Name</b>	<b>Scientific Name</b>
	Canola (Rape)	Brassica napus
	Capeweed	Arctotheca calendula
	Charlock	Sinapis arvensis
	Chickweed	Stellaria media
	Common Sowthistle (Milk Thistle)	Sonchus oleraceus
	Corn Gromwell	Buglossoides arvensis
	Cowvine	Ipomoea lonchophylla
	Crassula	Crassula spp.
	Deadnettle	Lamium amplexicaule
	Dense-flower Fumitory	Fumaria densiflora
	Dock	Rumex spp.
	Doublegee (Spiny Emex)	Emex australis
	Fat Hen	Chenopodium album
	Fireweed	Senecio spp.
	Fumitory	Fumitory spp.
	Hedge Mustard	Sisymbrium officinale
	Hexham Scent (King Island Melilot)	Melilotus indicus
	Horehound	Marrubium vulgare
	Hyssop Loosestrife	Lythrum hyssopifolia
	Iceplant	Mesembryanthemum spp.
	Indian Hedge Mustard	Sisymbrium orientale
	London Rocket	Sisymbrium irio
	Long Storksbill	Erodium botrys
	Marshmallow	Malva parviflora
	Mouse-eared Chickweed	Cerastium glomeratum
	Night-scented Stock	Matthiola longipetala
	Paterson's Curse	Echium plantagineum
	Peppercress	Lepidium spp.
	Prickly Lettuce	Lactuca scariola
	Purple Goosefoot	Scleroblitum atriplicinum
	Rough Poppy	Papaver hybridum
	Saffron Thistle	Carthamus lanatus
	Scarlet Pimpernel	Anagallis arvensis
	Shepherd's Purse	Capsella bursa-pastoris
	Skeleton Weed	Chondrilla juncea
	Sorrel	Rumex spp.
	Stemless Thistle	Onopordum acaulon
	Toad Rush	Juncus bufonius
	Tree Hogweed	Polygonum patulum
	Turnip Weed	Rapistrum rugosum
	Variogated Thistle	Silybum marianum
	Vetch (Tares)	Vicia sativa
	Volunteer Lupins	Lupinus spp.
	Ward's Weed	Carrichtera annua
	Wild Radish	Raphanus raphanistrum
	Wild Turnip	Brassica tournefortii
	Wireweed (Hogweed)	Polygonum aviculare

**DIRECTIONS FOR USE**

**RESTRAINTS**

DO NOT apply to crops or weeds which are stressed due to dry or excessively moist conditions.  
 DO NOT apply to crops under stress due to disease or insect damage.  
 DO NOT apply to frost affected crops or if frosts are imminent.  
 DO NOT apply when heavy rain is expected within 4 hours.

Crop	Weeds Controlled	Stage of Weed Growth	State	Rate /ha	Critical Comments
<b>CEREALS</b> Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)	Wild Radish	Up to the 2 leaf stage and not more than 60mm in diameter	WA only	250mL	<b>CROP STAGE</b> <b>Cereals</b> <b>Up to 750mL</b> (3 leaf to late tillering stage - Z13 to 30) <b>Over 750mL</b> (5 leaf to late tillering stage - Z15 to 30) Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing).  <b>WA only:</b> <b>DO NOT</b> apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).  <b>Warning:</b> Rygrex may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to ' <b>Crop Tolerance</b> ' section of <b>General Instructions</b> ).  Application is recommended prior to the eighth trifoliate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or Lucerne.  <b>Warning:</b> Rygrex may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to ' <b>Crop Tolerance</b> ' section of <b>General Instructions</b> ).  <b>WEED STAGE</b> Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: • high crop and weed density • late season germinations • abnormal weed growth (including early flowering), high rate of product (up to the maximum rate of application specified for that weed) may be required. Rygrex will not effectively control: • regrowth of suppressed weeds; • transplanted weeds; • regrowth from rhizomes or roots; • weeds growing under stress from previous herbicide applications.  <b>GRAZING</b> Efficacy on larger weeds will be improved by grazing with normal levels of stock after 7 day withholding period. Refer to 'Protection of Livestock' for grazing precautions.  <b>APPLICATION</b> Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased.  <b>WILD RADISH</b> Rygrex will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where: • rates lower than 1.0L/ha are used; • dry conditions prevail; • poor coverage of the soil surface is achieved; • crop is planted in non-wetting sand; • soils contain a high content of organic matter. Optimum results will be obtained if good soil moisture exists at and after application.
		Up to the 4 leaf stage and not more than 120mm in diameter	All States	500mL	
		Up to the 6 leaf stage and not more than 150mm in diameter		750mL	
		Up to the 8 leaf stage and not more than 180mm in diameter		1.0L	
	Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip	Up to the 2 leaf stage and not more than 60mm in diameter		500mL	
		Up to the 4 leaf stage and not more than 120mm in diameter		750mL	
		Up to the 6 leaf stage and not more than 150mm in diameter		1.0L	
	London Rocket	Up to the 5 leaf stage and not more than 120mm in diameter	Qld only	750mL	
	Ward's Weed		SA only		
	Capeweed	Up to the 2 leaf stage and not more than 60mm in diameter	All States	500mL	
		Up to the 4 leaf stage and not more than 120mm in diameter		1.0L	
	Crassula	Up to the 2 leaf stage		500mL	
		Up to the 4 leaf stage		750mL	
	Prickly Lettuce	Up to the 2 leaf stage		500mL	
		Up to the 4 leaf stage		750mL	
Up to the 6 leaf stage			1.0L		
Dense-flower Fumitory	Up to the 2 leaf stage		750mL		
Corn Gromwell, Saffron Thistle, Toad Rush	Up to the 2 leaf stage		1.0L		
Deadnettle		NSW, Vic, SA only	1.0L		
Sorrel	Up to the 2 leaf stage	Vic only	1.0L		
Canola (Rape)	Up to the 4 leaf stage	All States	500mL		
Purple Goosefoot	Up to the 6 leaf stage	Qld only	500mL		
Turnip Weed Wild Turnip	Cotyledon to 2 leaf	NSW only (West of Newell Hwy.) SA only (Eyre peninsula north of the line between Venus Bay and Cowell)	350mL		
<b>CEREALS</b> Wheat, Barley, Oats, Triticale, Cereal Rye	Fumitory	2 - 6 leaf stage	All States	500mL + 200mL terbutylm (500g/L)	



**SUPPRESSION OF THE FOLLOWING WEEDS**

Crop	Weeds Controlled	Stage of Weed Growth	State	Rate /ha	Critical Comments
<b>CEREALS</b> Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)  <b>PASTURE</b> Newly Sown and Established Clover-based pasture, Clover for Hay and Seed Production	Saffron Thistle	Up to the 6 leaf stage	All States	1.0L	<b>Refer to critical comments in the previous section.</b>
	Chickweed, Fireweed, Hexham Scent (King Island Melilot), Iceplant, Mouse-eared	Up to the 4 leaf stage		750mL	
	Chickweed, Night-scented Stock, Paterson's Curse, Peppergrass, Skeleton Weed, Long Storksbill, Volunteer Lupins.				
	Wireweed (hogweed)			750mL	
	Common Sowthistle (Milk Thistle), Cowvine, Dock, Doublegee, (Spiny Emex), Fat Hen, Horehound, Hyssop, Loosestrife, Marshmallow, Rough Poppy, Scarlet Pimpernel, Stemless Thistle, Tree Hogweed, Varigated Thistle, Vetch (Tares).	Up to the 2 leaf stage		1.0L	

Crop	Weeds Controlled	Stage of Weed Growth	State	Rate /ha	Critical Comments
<b>CEREALS</b> Wheat, Barley, Oats, Triticale, Cereal Rye	Wild Radish	Up to the 4 leaf stage and not more than 120mm in diameter	All States	350mL plus 200mL L.V.E. MCPA (500g/L)	<b>Refer also to all Critical Comments relating to weed stage, grazing, application and wild radish above.</b> *Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180mm in diameter. DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics. <b>Crop Stage</b> Rygrex 350mL + L.V.E. MCPA 200mL: Apply from 3 leaf to fully tillered (Zadoks Z13 to Z30). Rygrex 500mL + L.V.E. MCPA 200mL: Apply from 3 leaf to fully tillered (Zadoks Z13 to Z30). Rygrex 500mL + L.V.E. MCPA 400mL: Apply from 5 leaf stage to fully tillered (Zadoks Z15 to Z30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post-sowing). <b>WA only: DO NOT</b> apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). <b>Warning:</b> Rygrex may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to 'Crop Tolerance' section of General Instructions). Observe instructions also on L.V.E. MCPA product label.
		Up to the 6 leaf stage and not more than 150mm in diameter		500mL plus 200mL L.V.E. MCPA (500g/L)	
		Up to the 8 leaf stage and not more than 180mm in diameter*		500mL plus 400mL L.V.E. MCPA (500g/L)	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.**

**All Crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**GENERAL INSTRUCTIONS  
 RESISTANT WEEDS WARNING**

**GROUP F HERBICIDE**

Rygrex is a member of the phenoxy and nicotinanilide groups of herbicides and acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth. For weed resistance management Rygrex is both a Group F and a Group I herbicide. Some naturally occurring weed biotypes resistant to Rygrex and other Group F and Group I herbicides may exist through normal genetic variability in and weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Rygrex or other Group F or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Rygel Australia Pty Ltd accepts no liability for any losses that may result from the failure of Rygrex to control resistant weeds.

**Tolerance**

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Sub-clover is particularly sensitive.

**Cereals**

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

**Warning (Oats)**

The tolerance of varieties Esk and Nile (the two main varieties grown in Tasmania) to Rygrex has not been tested. Test a small area of crop before using Rygrex over large areas. Consult your local Rygel Australia Pty Ltd representative for advice on specific varieties.

**Pasture**

The tolerance of clover varieties to Rygrex can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

**Warning**

Rygrex may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500mL/ha and in areas of double spray. For this reason application is recommended prior to the 8<sup>th</sup> trifoliolate leaf stage. However, at the lower rates (500mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Rygrex applied at 500mL/ha:

**Arrowleaf:** Zulu

**Berseem:** Sacramonte

**White:** Haifa

**Balansa:** Paradana

**Persian:** Kyambro, Lupers, Maral

**Subterranean Clover:** Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup. The effects of Rygrex on clover seed yield have been tested on the following varieties. Under normal growing conditions they show levels of tolerance to Rygrex applied at 500mL/ha.

**Subterranean Clover:** Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

**Warning**

Rose and Strawberry clover have shown increased sensitivity to Rygrex. Rygrex may affect the seed of yield of subterranean variety Woogenellup. Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of Rygrex. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward. Varieties not listed should be tested before using Rygrex over large areas. Consult your local Rygel Australia Limited representative for advice on specific varieties.

**Subsequent Crops**

To reduce effect on susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

**Mixing**

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing Rygrex should not be left to stand overnight. Prolonged periods of exposure to cold temperature could result in settling out of the product in the mixture.

**Warning**

The rubber components present in some spraying units may be affected by exposure to the solvents in Rygrex and some other agricultural products. To reduce the risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

**Application**

**Boom Sprayer**

A minimum of 50L of water per hectare should be used, however, for optimum results water rates of 70-100L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

**Selective Herbicide**

**Aircraft (NSW, Vic, SA only)**

Apply in a minimum of 30L water per hectare. Effective control will only be achieved where good coverage of leaf surface is achieved.

**Compatibility**

The following products are physically compatible with Rygrex as a two-way mixture in the spray tank but should only be used for the crops specified:

Crop	Rygrex	Compatible Product
Wheat, triticale and cereal rye only	Up to 750mL/ha	Hoegrass (also barley), Puma Progress (wild oats only), Tristar Advance (1.5L rate only), Wildcat (wild oats only)
Cereals (including undersown)	All rates	Chlorpyrifos (500g/L), dimethoate, endosulfan
Cereals (not undersown)	Up to 500mL/ha	Rygel Metsulfuron, Rygel Chlorsulfuron, L.V.E. MCPA, Rygel triasulfuron
	All rates	Bromoxynil (200g/L), Amine 625, clopyralid, Tordon 50-D, dicamba (900g/kg) (up to 115g only), Eclipse
Wheat, Barley, Triticale, and Cereal Rye only (not undersown)		Rygel Tralkoxydim
Wheat only (not undersown)		Rygel Clodinafop
Clover	Up to 750mL/ha	Targa, Fusilade
Sub clover		Simazine Liquid, Simazine Liquid + Nuquat mixture
	Up to 1.0L/ha	2,4-DB amine (500g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Hoegrass, Wildcat, Puma Progress or Tristar Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with dicamba dry a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties. Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label. DO NOT add surfactant when mixing Rygrex and Rygel Metsulfuron.

**Simazine:** Refer to the simazine label for correct application rates, especially with regard to soil types. This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on the product: Hallmark 50EC, Dominex 100EC, Karate, Decis Forte EC, and Talstar.

**Warning**

DO NOT use crop oils with Rygrex or Rygrex mixtures with other products in cereals. As formulations of other manufacturer's products are beyond the control of Rygel Australia Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.**

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

**PROTECTION OF LIVESTOCK**

**Grazing Precaution**

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where capeweed, Paterson's curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's curse. If in doubt, contact your nearest Department of Agriculture.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. The method of disposal of the container depends on the container type. Read the 'Storage and Disposal' statement on the label that is attached to the container.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of Rygel Rygrex Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Rygel Australia Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Rygel Australia Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

**In an Emergency Dial 000 Police or Fire Brigade**

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**  
**Rygel**

**Rygrex**  
**SELECTIVE HERBICIDE**

ACTIVE CONSTITUENTS: 250 g/L MCPA present as the ethyl hexyl ester  
25 g/L DIFLUFENICAN  
SOLVENTS: 325 g/L LIQUID HYDROCARBONS,  
150 g/L N-METHYL-2-PYRROLIDONE

**GROUP FI HERBICIDE**

**For the control of certain broadleaf weeds in winter cereals and  
clover as specified in Directions for Use Table.**

**IMPORTANT: READ THIS BOOKLET BEFORE USE**

APVMA Approval No.: XXXXX/0310



Rygel Australia Pty Ltd ACN: 106 839 007  
103 Ordish Road, Dandenong South, Vic 3175  
Tel: 03 9768 2803 Fax: 03 9768 2804

130 mm

140 mm

**Rygel Rygrex Selective Herbicide**

**DIRECTIONS FOR USE**

For complete directions for use please read attached booklet.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

**5L, 20L containers only**

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and setup for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

**Re-Fillable Containers**

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty product as required into application equipment. Do not attempted to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that equipment used in transfer of the product is disconnected, triple rinsed and drained after each use.

When the container is empty, close all caps and valves and return the container to the point of purchase.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray

wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of Rygel Rygrex Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Rygel Australia Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Rygel Australia Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

APVMA Approval No: 64747/0310

Batch No: } see container  
DOM:

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**



**Rygel Australia Pty Ltd**  
ACN: 106 839 007  
103 Ordish Rd, Dandenong South, Vic 3175  
Tel: 03 9768 2803 Fax: 03 9768 2804

Bottom layer - rear label (sticks to container)

12 mm



18 mm

15 mm





15 mm

12-mm strip NOT  
peelable overglass



15mm x 15mm  
peel tabs from  
backing sheet



Adhesive side of 'top' label  
this side sticks to the 'bottom' label