

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Tufman

Infopest



Verified

HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON

GROUP B HERBICIDE

For the control of annual ryegrass, paradoxa grass and certain broadleaf weeds in cereal crops as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245

South Sydney Corporate Park, Unit W2E, 75-85 O'Riordan Street, Alexandria NSW 2015

T: 02 8338 9473 F: 02 9313 4352 W: www.farmalinx.com

NET
CONTENTS
1 kg



**APPROVED
LABEL**

TUFMAN 1KG LABEL – Size: 135 mm wide x 110mm deep
COLOURS: BLUE 285, PURPLE, GREEN 354

FILENAME: 64775_49241_1KG Label_MPL_V1
SCALE SIZE: PRINT AT 100% ON A4/A3 PAPER
DATE: 19/12/09

FARMALINX

 **Tufman**

HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Tufman Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

APVMA Approval No.: 64775/1kg/0110



IN A TRANSPORT
EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

SPECIALIST ADVICE IN AN EMERGENCY ONLY

1800 024 973

ALL HOURS – AUSTRALIA WIDE

Batch No.

Date of Manufacture

farmalinx



DIRECTIONS FOR USE

RESTRAINTS:
DO NOT store FARMALINX Tufman Herbicide whilst in suspension for more than 2 days or significant breakdown will occur.
DO NOT store tank mixes of FARMALINX Tufman Herbicide and other products for more than 24 hours to avoid possible product breakdown.
DO NOT spray weeds for post emergent control if rainfall is expected within 4 hours.
DO NOT apply to plants suffering stress.

METHOD OF USE: PRE-SOWING INCORPORATED BY SOWING

Table with 5 columns: CROP/SITUATION, WEEDS CONTROLLED, STATE, RATE g/ha (SOIL TYPE, Light to Medium, Heavy), CRITICAL COMMENTS. Includes rows for Wheat and Triticale only.

Main table with 5 columns: CROP/SITUATION, WEEDS CONTROLLED, STATE, RATE g/ha, CRITICAL COMMENTS. Lists various crops and weeds like Annual Ryegrass, Black Bindweed, etc.

METHOD OF USE: POST CROP AND WEED EMERGENCE

Table with 5 columns: CROP/SITUATION, WEEDS CONTROLLED, STATE, RATE g/ha (SOIL TYPE, Light to Medium, Heavy), CRITICAL COMMENTS. Includes rows for Wheat, Barley, Oats, etc.

Main table with 5 columns: CROP/SITUATION, WEEDS CONTROLLED, STATE, RATE g/ha, CRITICAL COMMENTS. Lists various crops and weeds like Corn Gromwell, Black Bindweed, etc.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

www.farmalinx.com

CONTINUED OVERLEAF

STORAGE AND DISPOSAL: Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.
PROTECTION OF WILDLIFE: DO NOT apply to areas used for grazing or to areas where there are nesting birds.
PROTECTION OF HUMANS AND OTHER ANIMALS: DO NOT apply to areas where there are children or other animals.
COMPARABILITY: FARMALINX Tufman Herbicide can be tank mixed with paraquat and glyphosate.
MIXING: FARMALINX Tufman Herbicide is a water dispersible granule to be mixed with water and applied as a spray.

Table with 5 columns: Soil pH, Minimum Interval for Recropping (in Months from Application), Cereals, Barley, Oats, etc. Includes notes on soil pH and recropping intervals.

SAFETY DIRECTIONS: Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET: Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Tufman Herbicide is available from FARMALINX Pty Ltd on request.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use.

Other registered trademarks

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E,
75-85 O'Riordan Street, Alexandria NSW 2015
T: 02 8338 9473 F: 02 9313 4352
Website: www.farmalinx.com

APVMA Approval No.: 64775/0110

CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FARMALINX
Tufman
HERBICIDE
ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON
GROUP B HERBICIDE
For the control of annual ryegrass, paradoxa grass and certain broadleaf weeds in cereal crops as specified in the Directions for Use table.
IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

GENERAL INSTRUCTIONS: FARMALINX Tufman Herbicide is a systemic herbicide which acts by inhibiting cell division and growth in the tips of roots and shoots of sensitive plants.

PRE-SOWING INCORPORATED BY SOWING: WA ONLY: Avoid applying to dry, sandy soils. Rapid leaching is possible with early season rainfall.

CONVENTIONALLY SOWN CROPS: On soils with pH less than 7 application should occur just before sowing or with the actual sowing operation. On soils with pH above 7 timing the application just prior to sowing is critical.

POST CROP AND WEED EMERGENCE: Where treatment is delayed or where weeds are not actively growing due to adverse conditions results may be slow to appear and weeds may be only stunted or suppressed.

Wheat, Triticale and Cereal Rye - apply after crop emergence when weeds are small and actively growing (Annual Ryegrass no more than 3 leaves, Zadok's Code 13). Broadleaved weeds no more than 5 cm in height or diameter (for Black Bindweed refer to specific recommendations).

Barley and Oats - apply between 2 leaf stage of crop (3 leaf stage - SA only) and early tillering when weeds are small and actively growing (Annual Ryegrass no more than 3 leaves, Zadok's Code 13). Broadleaved weeds no more than 5 cm in height or diameter (for Black Bindweed refer to specific recommendations).

ALL CROPS: DO NOT use more than once each season in any one crop.

RESISTANT WEEDS WARNING: FARMALINX Tufman Herbicide is a member of the sulfonylurea group of herbicides. FARMALINX Tufman Herbicide has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action.

GROUP B HERBICIDE: FARMALINX Tufman Herbicide is a Group B herbicide. Some naturally occurring weed biotypes resistant to FARMALINX Tufman Herbicide and other Group B herbicides (including Annual Ryegrass and some broadleaf weeds) are known to exist.

The following table applies to all states. Soil resistance (grom through to maturity) should be tested on small scale strips before sowing larger areas. FARMALINX Tufman Herbicide treated soils may be sown to any of the crops listed but only after the intervals indicated have been observed.

THE FOLLOWING TABLES REFER SPECIFICALLY TO NSW & VIC ONLY: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.

THE FOLLOWING TABLES APPLY SPECIFICALLY TO QLD, SA, WA AND TAS ONLY: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.

THE FOLLOWING TABLES REFER SPECIFICALLY TO NT: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.

THE FOLLOWING TABLES REFER SPECIFICALLY TO WA & VIC ONLY: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.

THE FOLLOWING TABLES REFER SPECIFICALLY TO QLD, SA, WA AND TAS ONLY: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.

THE FOLLOWING TABLES REFER SPECIFICALLY TO NT: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.

THE FOLLOWING TABLES REFER SPECIFICALLY TO WA & VIC ONLY: FARMALINX Tufman Herbicide is NOT recommended for use on soils of pH 8.6 and above.