

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Conquest

BROM M

SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 200g/L BROMOXYNIL present as the N-OCTANOYL ESTER
200g/L MCPA present as the ETHYL HEXYL ESTER
SOLVENT: 371g/L LIQUID HYDROCARBON

GROUP	C I	HERBICIDE
-------	------------	-----------

For the control of broadleaf weeds in wheat, oats, barley, cereal rye and triticale, linseed, grass pastures and turf as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS:

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
76 Walters Drive
Osborne Park WA 6017

Telephone (08) 9347 0500
Facsimile: (08) 9347 0551

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

For Non-Refillable Containers: Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs, Conquest Crop Protection Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS:

Product is poisonous if inhaled or swallowed. Attacks the eyes and will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

In a Transport Emergency Dial 000 Police, or Fire Brigade	FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 ALL HOURS AUSTRALIA WIDE
------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------

Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932

76 Walters Drive

Osborne Park WA 6017

Telephone (08) 9347 0500

Facsimile: (08) 9347 0551

Batch No:

DOM

APVMA Approval No: 64809 | 52837 B

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Conquest **BROM M** SELECTIVE HERBICIDE

**ACTIVE CONSTITUENTS: 200g/L BROMOXYNIL present as the N-OCTANOYL ESTER
200 g/L MCPA present as the ETHYL HEXYL ESTER
SOLVENT: 371g/L LIQUID HYDROCARBON**

GROUP	CI	HERBICIDE
--------------	-----------	------------------

For the control of broadleaf weeds in wheat, oats, barley, cereal rye and triticale, linseed, grass pastures and turf as specified in the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
76 Walters Drive
Osborne Park WA 6017

Telephone (08) 9347 0500
Facsimile: (08) 9347 0551

DIRECTIONS FOR USE:**RESTRAINTS:**

DO NOT spray if weeds are wilted by dry or cold weather or if crops are at a susceptible stage, or if rain is likely within 4 hours.

DO NOT apply if frosts are imminent.

DO NOT apply to diseased or frost affected crops or turf.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
Wheat, Barley, Cereal Rye, Oats and Triticale, Linseed; Grass Pastures	Amsinckia (<i>Amsinckia</i> spp.), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Charlock (<i>Sinapis arvensis</i>), Corn Gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Hexham Scent (King Island Melilot) (<i>Melilotus indica</i>), Horned Poppy (<i>Giaucium flavum</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mexican Poppy (<i>Argemone ochroleuca</i>), Mintweed (<i>Salvia reflexa</i>), Patersons Curse (Salvation Jane) (<i>Echium plantagineum</i>), Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella bursapastoris</i>), Slender Thistle (<i>Carduus tenuiflorus</i> , <i>C. pycnocephalus</i>), Tree Hogweed (<i>Polygonum patulum</i>), Turnip Weed (Giant Mustard) (<i>Rapistrum rugosum</i>), Wild Radish (<i>Raphanus raphanistrum</i>) Wild Turnip (<i>Brassica tournefortii</i>)	Plants up to the 6 leaf stage but not more than 50 mm in diameter	WA only	750 mL/ha	<p>CROP STAGE: Wheat, Barley, Cereal Rye, Oats and Triticale: 3 leaf to fully tillered stage (Z13-30) for rates of 750 mL to 1.4 L/ha; 5 leaf to fully tillered stage (Z15-30) for rate of 2.0 L/ha (NSW, ACT, Vic, Tas, SA, WA only); or early tillering to fully tillered stage (Z21-30) (Qld only).</p> <p>Linseed: 50 to 150 mm high. DO NOT apply to linseed by aircraft.</p> <ul style="list-style-type: none"> If spraying Black Bindweed when wheat, barley, cereal rye, oats, triticale or linseed is more than 300 mm high, increase spray volume to 200 L/ha. <p>For optimal control, apply only when the weeds are actively growing. In the early stages following spraying, temporary wilting of crop and pasture may occur. Crop scorch may occur, particularly on oats if treatment is delayed to later stages of growth. Refer to the General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. DO NOT apply to crops or pastures undersown with lucerne, clover or medics unless some damage is acceptable.</p>
			Qld, NSW, ACT, Vic, Tas, SA, NT only	1.4 L/ha	
		Plants up to the 8 leaf stage but not more than 75 mm in diameter	2.0 L/ha		
			WA only	1.4 L/ha	
	Black Bindweed* (Climbing Buckwehat) (<i>Fallopia convolvulus</i>), Common Peppergrass (<i>Lepidium africanum</i>), Mustards (<i>Sisymbrium</i> spp), Purple Calandrinia (<i>Calandrinia menziesii</i>)	Plants up to the 4 leaf stage but not more than 35 mm in diameter		1.0 L/ha	

Wheat, Barley, Cereal Rye, Oats and Triticale; Linseed; Grass Pastures - continued		Plants up to the 6 leaf stage but not more than 50 mm in diameter	All States	1.4 L/ha		
		Plants up to the 8 leaf stage but not more than 75 mm in diameter		2.0 L/ha		
Fumitory (<i>Fumaria spp</i>), Threecornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>)		Plants up to the 3 leaf stage but not more than 50 mm in diameter	Qld, NSW, ACT, Vic, Tas, SA only	1.4 L/ha	CROP STAGE: Wheat, Barley, Cereal Rye, Oats and Triticale: 3 leaf to fully tillered stage (Z13-30) for rates of 750 mL to 1.4 L/ha; 5 leaf to fully tillered stage (Z15-30) for rate of 2.0 L/ha (NSW, ACT, Vic, SA, WA, Tas only); or early tillering to fully tillered stage (Z21-30) (Qld only). Linseed: 50 to 150 mm high. DO NOT apply to linseed by aircraft. For optimal control, apply only when the weeds are actively growing. In the early stages following spraying, temporary wilting of crop and pasture may occur. Crop Scorch may occur, particularly on oats if treatment is delayed to later stages of growth. Refer to the General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. ** Where Wireweed (Hogweed) occurs in red soils of low fertility, it has been found to be less susceptible. In these circumstances use 2.0 L/ha.	
		Plants up to the 4 leaf stage but not more than 60 mm in diameter	WA only	750 mL/ha		
		Plants up to the 5 leaf stage but not more than 75 mm in diameter	Qld, NSW, ACT, Vic, Tas, SA only	2.0 L/ha		
		Plants up to the 6 leaf stage but not more than 85 mm in diameter	WA only	1.4 L/ha		
	Variegated Thistle (<i>Silybum marianum</i>), Wireweed (Hogweed)** (<i>Polygonum aviculare</i>)		Plants up to the 4 leaf stage but not more than 35 mm in diameter	Qld, NSW, ACT, Vic, Tas, SA only		750 mL/ha
			Plants up to the 6 leaf stage but not more than 50 mm in diameter			2.0 L/ha
				WA only		1.4 L/ha
	Mountain Sorrel (<i>Oxalis acetosella</i>), Three-horned Bedstraw (<i>Galium tricomutum</i>)		Plants up to the 6 leaf stage			750 mL/ha
			Plants up to the 8 leaf stage			1.4 L/ha
	Cleavers (<i>Galium aparine</i>)		2-4 stem stage and 1-3 whorls of leave per stem			750 mL/ha
Wheat, Barley, Cereal Rye, Oats, Triticale	Common Sowthistle (<i>Sonchus oleraceus</i>)				Apply under good growing conditions. Avoid spraying when soil is dry. DO NOT spray under frosty conditions. In early stages following spraying,	

					temporary wilting of crop may occur. Avoid application when maximum daily temperatures above 20°C or are likely to occur for a few days after application. DO NOT apply to crops undersown with lucerne, clover or medics unless some damage is acceptable.
		Up to 4 leaves	Northern NSW and Qld only	1.4 L/ha	CROP STAGES: Wheat, Barley, Oats, Triticale: 3 leaf to full tillering (Z13-30). Use as early as possible to avoid shading of weeds by the crop. Strictly for use on small weeds early in crop before shading out. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron apply 2.1 L/ha. Use a minimum spray volume of 70 L/ha.
		4 to 8 leaves		2.1 L/ha	CROP STAGES: Wheat, Barley, Oats and Triticale: 3 leaf to full tillering (Z13-30). Use as early as possible to avoid shading of weeds by the crop.
Turf, established Couch (incl. Queensland Blue), Kikuyu, Paspalum, Buffalo, Fescue, Bent & Rye lawns	Bindy-eye, Jo-Jo (Onehunga), Cudweed, Dandelion, Catsear, Plantains, Thistle, Oxalis, Capeweed, Creeping Buttercup, Fleabane, Clover (not Tas)	-	All States	300 mL/100L water or 60 mL/100m ²	Apply evenly over the area to be treated. DO NOT mow turf for 2 days after treatment.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions.

Mixing Information

To ensure even mixing, half fill spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION

DO NOT use rain-drop or air-induction nozzles as coverage of weeds may not be adequate.

Boom Sprayer

Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed.

Cereals

Apply 50 to 200 litres of water per hectare. Use higher water volume (min 100 L/ha) where the weed infestation is heavy, the crop cover is dense or when tank mixing with diclofop-methyl. Complete coverage of weeds is essential.

Linseed

Apply a minimum of 220 litres of water per hectare.

Turf

Apply a minimum of 500 litres of water per hectare.

Aerial Application

Apply a minimum of 22 litres of water per hectare or 30 litres of water per hectare if tank mixing with diclofop-methyl. Markers should always move upwind before aircraft passes overhead. Application by micronair equipment may give variable results.

Equipment Maintenance

After use wash equipment thoroughly with Spraymate Tank & Equipment Cleaner and rinse several times.

Compatibility

This product is physically compatible in the spray tank with one of the following herbicides or insecticides in wheat and barley only.

Conquest Metsulfuron 600 (600 g/L), Conquest Diclofop (500 g/L), Conquest CamQuesta 500 (500 g/L), MCPA, Dimethoate, Omethoate, Endosulfan or Chlorpyrifos.

When mixed with metsulfuron-methyl or dicamba, DO NOT apply to cereals undersown with lucerne. When mixed with diclofop-methyl, DO NOT exceed 1.4 litres of this product per hectare. Observe all instructions on the mixture partner label. As formulations of other manufactures products are beyond the control of Conquest Crop Protection Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

Resistant Weeds Warning

GROUP	C I	HERBICIDE
-------	-----	-----------

Conquest BROM M Selective Herbicide is a member of both the nitriles and the phenoxy groups of herbicides. Conquest BROM M Selective Herbicide has both the inhibitors of photosynthesis at photosystem II and disruptors of plant cell growth mode of action. For weed resistance management, the product is a Group C and Group I herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group C and Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group C or Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT apply by misting machines.

NOTE: Maximum daily temperatures at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. Do not contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

Grazing Precaution

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where Capeweed, Paterson's Curse and Variegated Thistle predominate in the pasture. Avoid grazing with young or breeding stock.

DO NOT graze horses or pigs on Paterson's Curse.

If in doubt, contact your nearest Department of Agriculture.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

For Non-Refillable Containers: Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs, Conquest Crop Protection Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS:

Product is poisonous if inhaled or swallowed. Attacks the eyes and will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves face shield or goggles and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product.

The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

<p>In a Transport Emergency Dial 000 Police, or Fire Brigade</p>	<p>SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE</p>
---------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------

Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932

76 Walters Drive

Osborne Park WA 6017

Telephone (08) 9347 0500

Facsimile: (08) 9347 0551

APVMA Approval No:

6 4809 | 52837B