

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST



RADICATE

Selective Herbicide

ACTIVE CONSTITUENT: 250 g/L MCPA (present as the ethyl hexyl ester)
25 g/L DIFLUFENICAN
SOLVENTS: 150 g/L N-METHYL-2-PYRROLIDONE

GROUP **F I** HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE.

Contents: 110L, 20L, 10L



RLP
Approved

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
76 Walters Drive
Osborne Park WA 6017

Telephone: (08) 9347 0500
Facsimile: (08) 9347 0551

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Returnable containers (110L only): Empty contents fully into application equipment. Close all valves. Return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and face-shield or goggles. If products in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill or judgment in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

For Specialist advise in an Emergency Dial
1800 033 1111
24 hours Australia wide.

Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932

76 Walters Drive

Osborne Park WA 6017

Telephone (08) 9244 1268

Facsimile: (08) 9204 2831

Batch No.

DOM

APVMA Approval No: 65139/50249

DrumMuster Logo

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

RADICATE

Selective Herbicide

ACTIVE CONSTITUENT: 250 g/L MCPA (present as the ethyl hexyl ester)
25 g/L DIFLUFENICAN
SOLVENTS: 150 g/L N-METHYL-2-PYRROLIDONE

GROUP **F I** HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE.

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
76 Walters Drive
Osborne Park WA 6017

Telephone: (08) 9347 0500
Facsimile: (08) 9347 0551

DIRECTIONS FOR USE Restraints

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.
 DO NOT apply to crops under stress due to disease or insect damage. DO NOT apply to frost-affected crops or if frosts are imminent.
 DO NOT apply when heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ HA	CRITICAL COMMENTS	
CEREALS Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover)	Wild radish	Up to the 2 leaf stage and not more than 60 mm in diameter	WA only	250 mL	CROP STAGE Cereals Up to 750 mL (3 leaf to fully tillered stage - Z13 to 30) Over 750 mL (5 leaf to fully tillered stage - Z15 to 30) Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: Radicate may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions) Clover Application is recommended prior to the eighth trifoliate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or lucerne. Warning: Radicate may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to "Crop Tolerance" section of General Instructions). WEED STAGE Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: * high crop and weed density * late season germinations * abnormal weed growth (including early flowering), Higher rates of product (up to the maximum rate of application specified for that weed) may be required. Radicate will not effectively control: * regrowth of suppressed weeds; * transplanted weeds; * regrowth from rhizomes or roots; * weeds growing under stress from previous herbicide applications	
		Up to the 4 leaf stage and not more than 120 mm in diameter	All States	500 mL		
		Up to the 6 leaf stage and not more than 150 mm in diameter		750 mL		
		Up to the 8 leaf stage and not more than 180 mm in diameter		1.0 L		
	PASTURE Newly sown and established clover-based pasture, clover for hay and seed production	Charlock, hedge mustard, Indian hedge mustard, shepherd's purse, turnip weed, wild turnip	Up to the 2 leaf stage and not more than 60 mm in diameter	Qld only SA only		500 mL
			Up to the 4 leaf stage and not more than 120 mm in diameter			750 mL
			Up to the 6 leaf stage and not more than 150 mm in diameter			1.0 L
		London rocket	Up to the 5 leaf stage and not more than 120 mm in diameter	All States		750 mL
		Ward's weed	500 mL			
		Capeweed	Up to the 2 leaf stage and not more than 60 mm in diameter	All States		1.0 L
			Up to the 4 leaf stage and not more than 120 mm in diameter			500 mL
		Crassula	Up to the 2 leaf stage	All States		750 mL
Up to the 4 leaf stage			500 mL			
Prickly lettuce		Up to the 2 leaf stage	All States	750 mL		
	Up to the 4 leaf stage	1.0 L				
	Up to the 6 leaf stage	750 mL				
Dense-flower fumitory	Up to the 2 leaf stage	NSW, Vic, SA only	1.0 L			
Corn gromwell, saffron thistle, toad rush						
Deadnettle						

CONTINUED OVERLEAF

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover) PASTURE Newly sown and established clover based pasture, clover for hay and seed production	Sorrel	Up to the 2 leaf stage	Vic only	1.0 L	CONTINUED FROM PREVIOUS PAGE GRAZING Efficacy on larger weeds will be improved by grazing with normal levels of stock after the 7 day withholding period. Refer to 'Protection of Livestock' for grazing precautions. APPLICATION Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased. WILD RADISH Radicate will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where: <ul style="list-style-type: none"> * rates lower than 1.0 L/ha are used; * dry conditions prevail; * poor coverage of the soil surface is achieved; * crop is planted in non-wetting sand; * soils contain a high content of organic matter. Optimum results will be obtained if good soil moisture exists at and after application.
	Canola (rape)	Up to the 4 leaf stage	All States	500 mL	
	Purple goosefoot	Up to the 6 leaf stage	Qld only	500 mL	
	Turnip weed, wild turnip	Cotyledon to 2 leaf stage	NSW only (West of Newell Hwy.) SA only (Eyre peninsula north of the line between Venus Bay and Cowell)	350 mL	
CEREALS Wheat, barley, oats, triticale, cereal rye	Fumitory	2 - 6 leaf stage	All States	500 mL + 200 mL terbutryn (500 g/L)	
CEREALS Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover) PASTURE Newly sown and established clover based pasture, clover for hay and seed production	SUPPRESSION OF THE FOLLOWING WEEDS				
	Saffron thistle	Up to the 6 leaf stage	All States	1.0 L	
	Chickweed, fireweed, hexham scent (King Island mellilot), iceplant, mouse-eared chickweed, night-scented stock, Paterson's curse, peppergrass, skeleton weed, long storksbill, volunteer lupins	Up to the 4 leaf stage			
	Wireweed (hogweed)				
Common sowthistle (milk thistle), cowvine, dock, doublegee (spiny emex), fat hen, horehound, hyssop loosestrife, marshmallow, rough poppy, scarlet pimpernel, stemless thistle, tree hogweed, variegated thistle, vetch (tares)	Up to the 2 leaf stage	1.0 L			

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ HA	CRITICAL COMMENTS
CEREALS Wheat, barley, oats, triticale, cereal rye	Wild radish	Up to the 4 leaf stage and not more than 120 mm in diameter	All States	350 mL plus 200 mL MCPA LVE (500 g/L)	<p>Refer also to all Critical Comments relating to weed stage, grazing, application and wild radish above.</p> <p>* Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180 mm in diameter.</p> <p>DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics.</p> <p><u>Crop Stage</u> Radicate 350 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Radicate 500 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Radicate 500 mL + MCPA LVE 400 mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30).</p> <p>Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: Radicate may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions)</p> <p>Observe instructions also on MCPA LVE product label.</p>
		Up to the 6 leaf stage and not more than 150 mm in diameter		500 mL plus 200 mL MCPA LVE (500 g/L)	
		Up to the 8 leaf stage and not more than 180 mm in diameter*		500 mL plus 400 mL MCPA LVE (500 g/L)	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

All crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS:

RESISTANT WEEDS WARNING

GROUP F I HERBICIDE

Conquest Radicate Selective Herbicide is a member of the phenoxy and nicotinanilide group of herbicides and acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth. For weed resistance management, Conquest Radicate is both a Group F and a Group I Herbicide. Some naturally occurring weed biotypes resistant to Conquest Radicate and other Group F and I Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group F and I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

TOLERANCE

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Subterranean clover is particularly sensitive.

Cereals

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

Warning (Oats)

The tolerance of oat varieties Esk and Nile (the two main varieties grown in Tasmania) to Conquest Radicate has not been tested. Test a small area of crop before using Conquest Radicate over large areas. Consult your local Conquest representative for advice on specific varieties.

Pasture

The tolerance of clover varieties to Conquest Radicate can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

Warning

Conquest Radicate may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500 mL/ha and in areas of double spray. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, at the lower rates (500 mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Conquest Radicate applied at 500 mL/ha.

ArrowLeaf: Zulu

Balansa: Paradana

Berseem: Sacromonte

Persian: Kyambro, Lupers, Maral

White: Haifa

Subterranean clover: Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The effects of Conquest Radicate on clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to Conquest Radicate applied at 500 mL/ha:

Subterranean clover: Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

Warning

Rose and Strawberry clover have shown increased sensitivity to Conquest Radicate. Conquest Radicate may affect the seed yield of subterranean clover variety Woogenellup.

Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of Conquest Radicate.

Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward.

Varieties not listed should be tested before using Conquest Radicate over large areas. Consult your Conquest representative for advice on specific varieties.

Subsequent Crops

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixture containing Conquest Radicate should not be left to stand over night. Prolonged periods of exposure to cold temperatures could result in settling out of the product in the mixture.

Warning

The rubber components present in some spraying units may be affected by exposure to the solvents in Conquest Radicate and some other agricultural products. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray manufacturer to determine the suitability of the rubber components for use with agricultural products.

APPLICATION**Boom Sprayer**

A minimum of 50 L water / ha should be used, however, for optimum results water rates of 70-100 L / ha are recommended. Increase the water volume where weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential.

Aircraft (NSW, Vic, SA only)

Apply in a minimum of 30 L water per hectare. Effective weed control will only be achieved where good coverage of leaf surface is achieved.

COMPATIBILITY

The following products are physically compatible with Conquest Radicate as two way mixtures in the spray tank, but should only be used for the crops specified:

Crop	Conquest Radicate	Compatible Product
Wheat, triticale and cereal rye only	Up to 750 mL / ha	Conquest Diclofop 375 (barley also), Puma® Progress (wild oats only), Tristar® Advance (1.5 L rate only), Wildcat® (wild oats only)
Cereals (including undersown)	All rates	Chlorpyrifos (500 g/L), dimethoate, Thiodan®
Cereals (not undersown)	Up to 500 mL / ha	Conquest Metsulfuron 600, Conquest Chlorsulfuron 750, MCPA LVE, Conquest Rival 750
	All Rates	Bromoxynil 200 g/L, 2,4-D Amine 500 herbicide, Conquest Corsair, Conquest Buckwheat, Cadence® (up to 115 g only), Eclipse®
Wheat, barley, triticale and cereal rye only (not undersown)		Conquest Crossfire
Wheat only (not undersown)		Conquest Snooker 200 EC
Clover	Up to 750 mL / ha	Conquest Atomic 100, Fusilade®
Subterranean clover		Simazine (500 g/L), Simazine (500 g/L) + Paraquat (200 g/L) mixture
	Up to 1.0 L / ha	2,4-D amine (500 g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Conquest Diclofop 375, Wildcat, Puma Progress or Tristar Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with Cadence a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties.

Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plant back restrictions on that label. DO NOT add surfactant when mixing Conquest Radicate and Conquest Metsulfuron 600.

Simazine: Refer to the simazine label for correct application rates, especially with regard to soil types.

This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on this product: Hallmark® 50EC, Dominex® 100EC, Karate®, Decis Options® and Talstar®.

Warning

DO NOT use crop oils with Conquest Radicate or Conquest Radicate tank mixtures with other products in cereals.

As formulations of other manufacturers' products are beyond the control of Conquest Crop Protection, all mixtures should be tested prior to mixing commercial quantities.

WEEDS LIST

WEED (Common Name)	(Scientific name)
Canola (rape)	<i>Brassica napus</i>
Capeweed	<i>Arctotheca calendula</i>
Charlock	<i>Sinapis arvensis</i>
Chickweed	<i>Stellaria media</i>
Common sowthistle (milk thistle)	<i>Sonchus oleraceus</i>
Corn gromwell	<i>Buglossoides arvensis</i>
Cowvine	<i>Ipomoea lonchopylla</i>
Crassula	<i>Crassula spp.</i>
Deadnettle	<i>Lamium amplexicaule</i>
Dense-flower fumitory	<i>Fumaria densiflora</i>
Dock	<i>Rumex spp.</i>
Doublegee (spiny emex)	<i>Emex australis</i>
Fat hen	<i>Chenopodium album</i>
Fireweed	<i>Senecio spp.</i>
Fumitory	<i>Fumaria spp.</i>
Hedge mustard	<i>Sisymbrium orientale</i>
Hexham scent (Kind Island mellilot)	<i>Mellilotus indicus</i>
Horehound	<i>Marrubium vulgare</i>
Hyssop loosestrife	<i>Lythrum hyssopifolia</i>
Iceplant	<i>Mesembryanthemum spp.</i>
Indian hedge mustard	<i>Sisymbrium orientale</i>
London rocket	<i>Sisymbrium irio</i>
Long storksbill	<i>Erodium botrys</i>
Marshmallow	<i>Malva parviflora</i>
Mouse-eared chickweed	<i>Cerastium glomeratum</i>
Night-scented stock	<i>Matthiola longipetala</i>
Paterson's curse	<i>Echium plantagineum.</i>
Peppergrass	<i>Lepidium spp.</i>
Prickly lettuce	<i>Lactuca serriola</i>
Purple goosefoot	<i>Scleroblitum atriplicinum</i>
Rough poppy	<i>Papaver hybridum</i>
Saffron thistle	<i>Carthamus lanatus</i>
Scarlet pimpernel	<i>Anagallis arvensis</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>
Skeleton weed	<i>Chondrilla juncea</i>
Sorrel	<i>Rumex spp.</i>
Stemless thistle	<i>Onopordum acaulon</i>
Toad rush	<i>Juncos bufonius</i>
Tree hogweed	<i>Polygonum patulum</i>
Turnip weed	<i>Rapistrum rugosum</i>
Variegated thistle	<i>Silybum marianum</i>
Vetch (tares)	<i>Vicia sativa</i>
Volunteer lupins	<i>Lupinus spp.</i>
Ward's weed	<i>Carrichtera annua</i>
Wild radish	<i>Raphanus raphanistrum</i>
Wild turnip	<i>Brassica tournefortii</i>
Wireweed (hogweed)	<i>Polygonum aviculare</i>

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK**Grazing Precaution**

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where Capeweed, Paterson's curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

Returnable containers (110L only): Empty contents fully into application equipment. Close all valves. Return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale vapours. When opening the container, preparing spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and face-shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26). If swallowed, **DO NOT** induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

For Specialist advise in an Emergency Dial
1800 033 1111
24 hours Australia wide.

Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932

76 Walters Drive

Osborne Park WA 6017

Telephone: (08) 9347 0500

Facsimile: (08) 9347 0551

Batch No.

DOM

APVMA Approval No: 65139/50249

DrumMuster Logo