

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**CONQUEST**

# **TRICON MAXI 600**

## **SELECTIVE HERBICIDE**

Active constituent: 600 g/L TRIFLURALIN  
Solvent: 9.5 g/L LIQUID HYDROCARBON

<b>GROUP</b>	<b>D</b>	<b>HERBICIDE</b>
--------------	----------	------------------

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as listed in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY  
BEFORE USING THIS PRODUCT.**

**Contents:**  
**20L**  
(100L, 110L, 200L, 1000L)



**Conquest Crop Protection Pty Ltd.**  
ABN 84 098 814 932  
76 Walters Drive  
Osborne Park, WA 6017

Telephone (08) 9347 0500  
Facsimile: (08) 9347 0551

 **CONQUEST**  
CROP PROTECTION

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near food, feedstuffs, fertilisers or seed.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or preferably pressure rinse containers for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SPILL AND LEAK MANAGEMENT

Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways and drains. **Small spills/leaks:** Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal.

## SAFETY DIRECTIONS

Harmful if swallowed. Poisonous if inhaled. Will damage eyes. Will irritate the skin. May irritate nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. If product or spray on skin, immediately wash area with soap and water. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist, a washable hat and elbow length chemical resistant gloves, goggles and half face piece respirator. If product or spray in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

## CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

**For specialist advice in an emergency dial  
1 800 033 111  
24 hours Australia wide.**

## Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932  
76 Walters Drive  
Osborne Park WA 6017

Telephone (08) 9347 0500

Facsimile: (08) 9347 0551

Batch Number:

Date of Manufacture:

APVMA Approval No: 65526 / 58930

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**CONQUEST**

# **TRICON MAXI 600**

**SELECTIVE HERBICIDE**

Active constituent: 600 g/L TRIFLURALIN  
Solvent: 9.5 g/L LIQUID HYDROCARBON

<b>GROUP</b>	<b>D</b>	<b>HERBICIDE</b>
--------------	----------	------------------

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as listed in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY  
BEFORE USING THIS PRODUCT.**

**Conquest Crop Protection Pty Ltd.**  
ABN 84 098 814 932  
76 Walters Drive  
Osborne Park, WA 6017

Telephone (08) 9347 0500  
Facsimile: (08) 9347 0551

 **CONQUEST**  
CROP PROTECTION

## DIRECTIONS FOR USE

### 1. FIELD CROPS

Situation & Crop	Weeds	State	Rate L/Ha soil type			Critical Comments
			Light	Medium	Heavy	
Chickpeas	Annual Ryegrass, Paradoxa Grass (Canary Grass), Wireweed (Hogweed) Black Pigweed, Suppression of Climbing Buckwheat (Black Bindweed), soil surface Wild Oats	Qld only	1-1.4L/ha	1-1.4 L/ ha	1 - 1.4L/ha	Use 1L/ha when applying immediately prior to sowing. Use 1.4L/ha when applying to dry soil before the planting rain.
	Annual Ryegrass, Wireweed (hogweed), Deadnettles	Vic only	640 mL/ha plus 1L/ha Tri-Allate (500g/L)			Incorporate as per recommendations for wheat, barley and triticale.
	Red & White Fumitory, Rough Poppy, Wireweed, Annual Ryegrass, Barley Grass, Canary Grass, Sand Fescue, Suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burrweed, Brome Grass, Cereal Oats and soil surface Wild Oats	SA only	1L / ha	1 L / ha	1 L / ha	Apply to level seedbed 0 to 4 weeks before sowing. Incorporate as per Incorporation Table 6 for method of incorporation.
	Amaranthus, Annual Ryegrass, Barnyard Grass, Caltrop, Crab Grass,	NSW, ACT only	960mL - 1.2L/ha	1.2L/ ha	1.35L/ha	Apply from 4 weeks up to just prior to sowing. See Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Adzuki Beans, Cowpeas, Lablab, Mung Beans, Borlotti Beans, Red Kidney Beans	Paradoxa Grass (Canary Grass), Pigweed, soil surface Wild Oats, Winter Grass, Wireweed, Suppression of Fumitory	NSW, ACT, Qld only	960mL - 1.2L/ha	1.2L/ ha	1.35L/ ha	
Faba Beans	Annual Ryegrass, Barley Grass, Capeweed, Corn Gromwell (Sheepweed), Fumitories, Geranium, Ivy Leaf, Speedwell, Mustards, Turnips, Wireweed, Suppression of Brome Grass, Soursob and soil surface Wild Oats	SA & WA only	640mL/ha plus 1.1 kg/ha Conquest Simaquest 900 WG Herbicide			Apply to bare moist soil and grate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to ridged or excessively cloddy soil. For full reliable results, significant rainfall (20 or 30mm) is necessary within 2-3 weeks of application.
Pigeon Peas	Amaranthus, Barnyard Grass, Canary Grass, Crowsfoot Grass, Pigweed, Spiny Burrgrass, Summer Grass, soil surface Wild Oats, Wireweed (hogweed), Suppression of Yellow Vine (Caltrop), From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass	NSW, ACT only	960mL/ha	1.2L/ ha	1.35L/ha	Apply between 4 weeks and just before sowing. Refer to Incorporation Table 3, 4 or 6 for suitable method of incorporation.
Lentils	Annual Phalaris, Annual Ryegrass, Wild Oats, Wireweed	NSW, ACT only	640 mL/ ha	960mL/ ha	960mL/ha	Apply 1 to 4 weeks before sowing
	Fumitory - Red and White, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue	SA only	1L/ha	1L/ha	1L/ha	Apply 1 to 4 weeks before sowing.

Navy Beans	Annual Ryegrass, Barnyard Grass,	All States	960mL/ha	1.2L/ha	1.35L/ha	Spray between 4 weeks and just before sowing takes place. See Incorporation Table directions 3, 4 or 5.
Soybeans	Canary Grass, Caltrop (Bullhead/Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot ( <i>Amaranthus</i> ), Redshank (Prince of Wales Feather), Summer Grass, soil surface Wild Oats, Winter Grass, Wireweed (Hogweed), From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass ( <i>Urochloa</i> )		960mL/ha	1.2L/ha	1.85L/ha	
Vetch	Annual Ryegrass, Deadnettle, Wireweed, soil surface Wild Oats, suppression of Brome Grass, Rough Poppy, Speedwell, Three Cornered Jack, Yellow Burr Weed, Sheepweed	SA & WA only	1.35L/ha	1.35L/ha	1.35L/ha	Apply to level seedbed 0 to 4 weeks before sowing. Refer to incorporation Table 6 for method of incorporation.
Cotton	Annual Ryegrass, Barnyard Grass, Canary Grass, Caltrop (Bullhead/Yellow Vine), Crab grass,	Qld, NSW, WA only	960mL/ha	1.35L/ha	1.85L/ha	Spray between 6 weeks and just before sowing takes place. Refer incorporation Table 1 & 2 for method of incorporation.
Legume Seed Crop Establishment	Fumitory, Mossman River Grass (Innocent Weed), Phalaris spp., Pigweed, Redroot ( <i>Amaranthus</i> ), Redshank (Prince of Wales Feather), Summer Grass, soil surface Wild Oats, Winter Grass, Wireweed (Hogweed), Black Pigweed (Qld only), <i>Phalaris spp.</i> Fumitory From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass ( <i>Urochloa</i> )	NSW, ACT, SA, WA, Vic, Tas only	960mL/ha	960mL/ha	1.35L/ha	<b>Autumn Sowing</b> - Apply from 4 weeks to 7 days before sowing takes place. Refer Incorporation Table 6 for method of incorporation.  <b>Spring Sowing</b> - Apply between 4 weeks and 3 days before sowing takes place. Refer Incorporation Table 6 for method of incorporation.  In both cases seedling disease, cold weather, excessive moisture, high salt concentrations and drought could weaken crop seedlings and damage
- Annual Medics - Clover (Berseem, Red, Strawberry Sub & White) Lucerne - Lucerne for hay and seed crops		All States	960mL/ha	1.35 L/ha	1.35 L/ha	
Linseed		NSW, ACT, SA, WA, Vic only	960mL/ha	1.2L/ha	1.35L/ha	Spray 2-4 weeks before sowing. Sowing depth should be 1.3 to 2.5cm. Deeper sowing may result in some stand reduction. Refer incorporation Table 6 for method of incorporation
Peanuts		WA, Qld only				Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Peas		All States				Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 or 11 for method of incorporation.
Canola, Safflower						
Sugarcane - Early season Late season		Qld, NSW only	2.4L/ha	2.4L/ha	2.4L/ha	Apply to plant cane after emergence to "out of hand" stage. Apply to ratoon cane immediately after harvest. Refer Incorporation Table 3, 4 or 10 for suitable method of incorporation
Sunflowers		All States	1.85L/ha	1.85L/ha	1.85L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for method of incorporation
Lupins		All States	960mL/ha	1.2L/ha	1.35L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 for method of incorporation.  <b>WA Only:</b> Use higher rate for heavier stubble coverage. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. Refer Table 13 for method of incorporation.

	Annual Grasses and Broadleaf Weeds	NSW, ACT, Vic, Qld only	640 mL/ha plus 1.7 kg/ha Conquest Simaquest 900 WG Herbicide			Use a low volume boom applying 50-100 litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing the crop. Incorporate within 4 hours of application. Do not spray to a ridged soil.
	Capeweed, Turnip, Radish, Doublegee (Spiny Emex), Suppression of Annual Ryegrass and Wild Oats	WA only	1L/ha plus 560 - 830 g/ha Conquest Simaquest 900 WG Herbicide			Rate for Yellow Sands. See Incorporation Table Directions 11, 12 or 13.
	As above plus suppression of Brome Grass		1L/ha plus 1.1 kg/ha Conquest Simaquest 900 WG Herbicide			Rate for all other soil types. Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to rigid or excessively cloddy soil. For Simazine to be effective sufficient rainfall (20 to 30 mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Results with Simazine can be variable if seasonal conditions are dry prior to sowing and Lupins are sown into dry or low moisture seedbeds.
	Capeweed, Doublegee, Wild Radish, Wild Turnip plus suppression of Annual Ryegrass, soil surface Wild Oats and Brome Grass		1L/ha plus 1.1 kg/ha of Conquest Diuron 900 WG Herbicide			DO NOT use on white or grey sands as severe crop damage may result. Use tank mix of Diuron & Trifluralin where Annual Ryegrass is present. Apply at pre-sowing stage. See the incorporation method in Table 13. For Post-sowing Pre-emergent application, ensure seed is adequately covered with soil. Refer to incorporation Table 12.
	Red & White Fumitory, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue, suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr Weed, Brome Grass, Cereal Oats, soil surface Wild Oats	SA only	1 - 1.4L/ ha	1 - 1.4L/ ha	1 - 1.4L/ ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 for method of incorporation.
	Above weeds plus Capeweed, Common Fumitory, Geranium, Indian Hedge Mustard, Sheepweed, Shepherd Purse, Toadrush, Turnips, suppression of Ice Plant & Soursob		1L/ha to 1.4L/ha plus 1.1 - 2.2 kg/ha Conquest Simazine 900 WG Herbicide			Use a low volume boom applying 50-100 litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing the crop. Incorporate within 4 hours of application. Do not apply to ridged soil.
Tobacco	Summer Grass, Crowsfoot Grass, Red Natal Grass, Love Grass, Button Grass, Rhodes Grass, Pigweed	Qld only	640 mL/ha	960mL/ ha	-	Apply to soil 3-4 weeks prior to transplanting. The longer period to be used for applications made during June & July. Incorporate to a depth of 10 cm.
	Crowsfoot Grass	NSW, ACT, only			960mL/ha	Apply to light sandy soil 14 to 21 days before transplanting. Do not incorporate to a depth greater than 6 cm. Apply to loam (medium soil) 14 to 21 days before transplanting. DO NOT incorporate to a depth of greater than 6 cm.
Wheat, Barley & Triticale	Annual Ryegrass, Wireweed (Hogweed), Phalaris spp.	NSW, ACT WA & Vic only	640 mL/ha	640 mL/ha	640 mL/ ha	Apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Incorporation Table 6 for method of incorporation.
	Fumitory	WA only				
	Canary Grass	Vic only				
	1.Pre-Sowing Only	As above except for Fumitory	Vic only			
2.Pre-Sowing & Post-sowing (self mulching soils)					Pre - sowing - Apply more than 4 weeks before sowing to prevent crop damage. Post sowing- Apply within 2 days after sowing to well prepared seedbed. Refer Incorporation Table 6 for suitable method of incorporation.	

Wheat	Annual Ryegrass, Paradoxa Grass (Canary Grass), soil surface Wild Oats, Wireweed (Hogweed)	Qld only	640 mL/ ha	640 mL/ha	640 mL/ ha	On non-self mulching soils apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. On self mulching soils as above except apply more than 4 weeks before sowing to prevent crop damage. Refer incorporation Table 6 for method of incorporation.
Barley						
Wheat, Triticale, Rye	Annual Ryegrass, Red & White Fumitory, Phalaris spp, Wireweed, (Hogweed), suppression of Deadnettle, Rough Poppy, Yellow Burr Weed	SA only	1L/ha	1L/ha	1L/ha	Apply 1 - 4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer incorporation Table 6 for method of incorporation.
Barley						
Wheat & Triticale only	Annual Phalaris	NSW, ACT only	640mL/ha plus 20g/ha of 750 g/kg Chlorsulfuron			If possible spray and incorporate into the soil in one operation. If this is not possible incorporation should take place within 4 hours of spraying. Delay may cause inferior weed control.

## 2. FIELD CROPS: PRE SOWING OR INCORPORATED BY SOWING (IBS) (Using Incorporation Table 13)

Situation & Crop	Weeds	State	Rate	Critical Comments
Wheat, Barley, Triticale	Annual Ryegrass, Wireweed, Phalaris spp, Fumitory	WA, SA, Vic, NSW only	1.2 – 1.6L/ha	Use higher rate on lighter sandy and sandy loam soils. DO NOT use on heavy soils. Use with knife/blade point sowing equipment. Use higher rate for heavier stubble coverage. Stubble coverage above 40 – 50% ground cover can reduce weed control below acceptable levels. Refer to table 13 for methods of incorporation.
Chickpeas		WA only	1 – 1.4L/ha plus 1.1kg/ha of Conquest Simazine 900 WG Herbicide	Incorporate as per incorporation table 13.

## 3. VEGETABLES, ORCHARDS AND VINEYARDS

Situation & Crop	Weeds	State	Rate L/ha soil type			Critical Comments
			Light	Medium	Heavy	
Transplants only Broccoli, Cabbage, Cauliflowers, Tomatoes	Annual Ryegrass, Barnyard Grass, Canary Grass, Caltrop, (Bullhead, Yellow Vine), Crab	All states	1L/ha	1.4L/ha	1.85L/ ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Direct Seeded Only Broccoli, Brussels Sprouts, Cabbage	Grass, Mossman River Grass (Innocent Grass) Pigweed, Redroot ( <i>Amaranthus</i> )					
Cauliflower	Redshank (Prince of Wales Feather),					
Carrots	Summer Grass, Wild					
Chicory	Oats, Winter					
Green Beans	Grass, Wireweed (Hogweed)					



Orchards & Vineyards	From seed only:	Qld, SA, Tas, Vic, WA only	1L/ha	1.4L/ha	1.85L/ha	Apply to new planting during pre plant cultivation. Apply to established crops in spring after weeds and green manure crop has been ploughed into ground. Refer Incorporation Table 8 or 9 for suitable method of incorporation.
	Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	Qld, SA, WA, Vic, Tas only				

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

#### GENERAL INSTRUCTIONS

**THIS PRODUCT MUST BE INCORPORATED INTO THE SOIL WITHIN 4 HOURS OF APPLICATION EXCEPT WHERE THE CROP IS SOWN WITH MINIMUM TILLAGE SOWING EQUIPMENT (FITTED WITH KNIFE POINTS OR BLADES LESS THAN 12 MM WIDE, USUALLY WITH PRESS WHEELS) WHERE APPLICATION MAY OCCUR UP TO 24 HOURS PRIOR TO THE SOWING PROCESS.**

#### INCORPORATION TABLE

1. **Prior to furrowing out:** 2 workings at an angle required using Offset or Tandem disc harrows.
2. **After furrowing out:** 2 workings required using Go-Devil discs or Lilliston cultivators set at 10 cm depth.
3. **Rotary Hoe:** 1 working required at 5-7.5 cm depth. Sugar Cane: 7.5-13 cm depth.
4. **Offset or Tandem Disc Harrows:** (preferably with spiked harrows in tandem) 2 workings at an angle required at 7.5-15 cm depth at 6.5-10 km per hour.
5. **Heavy Diamond or Stump Jump Harrows** - (weighted 20-30 kg per section) at 10-13 km/h speed. Then cross work with offset or tandem disc harrows set to 7.5-15 cm depth at speed 6.5-10 km per hour.
6. **Weighted Heavy Diamond or Stump Jump Harrows** (weighted with 20-30 kg per section) at 10-13 km per hour. Cross work with combine at 5-7.5 cm depth at speed of 10-13 km/hr.
7. **Disc Ratoon Cultivator:** 2 workings needed with discs and cultivator set at 7.5-13 cm depth.
8. **Offset or Tandem Disc Harrows:** set at 7.5-15 cm depth. A second discing is required working in opposite direction with discs set to throw treated soil into tree or vine row.
9. **Rotary Hoe:** 1 working needed at 5-10 cm depth.
10. **Offset Discs (Bumpers):** 2 workings needed at depth of 7.5-13 cm.
11. Incorporation by sowing (IBS) on suitably prepared seedbed with heavy diamond harrows trailing or as a separate operation.
12. **Post-sowing/Pre-emergence:** Use heavy diamond harrows cross working at right angles to the direction of sowing. Do not attempt this method of incorporation on poorly prepared, clumpy or cloddy soils.
13. Incorporation by sowing (IBS) with knife or blade points. Use press wheels to avoid dragging treated soil back into the seed furrow. Maintain slow to moderate speed to ensure that soil throw is not into adjacent furrows. Note: knife or blade point systems can result in poor weed control in the seed furrow as chemical displacement from this zone occurs. Stubble coverage above 40 – 50% ground cover can reduce weed control below acceptable levels. A knife or blade point is 12 mm or less, has no wings, inverted T or blade and is generally placed on a minimum 200 mm tyne spacing.

#### MIXING

This product is an emulsifiable concentrate, which mixes readily with water. Add the recommended amount to the spray tank during filing operation and apply 70-450L of water/ha (broadcast basis) dependent on soil type and stubble coverage level. For minimum tillage/stubble retention seeding systems use of the higher water volumes may help to reduce the impact of stubble.

**Ensure adequate agitation is continued throughout the operation.** Leaving the made up mixture for long periods of time without agitation is not recommended.

Under hot conditions or where possible, spray and incorporate into the soil in one operation. Delay may cause inferior weed control. Use properly calibrated boom sprayer with nozzles that provide medium through to very coarse spray droplets.



## Integrated Weed Management

The use of Integrated Weed Management (IWM) techniques in conjunction with Conquest Tricon Maxi 600 are always recommended. Agronomic practices that reduce the weed seed bank in the soil prior to the use of Conquest Tricon Maxi 600 will result in higher weed control levels from Conquest Tricon Maxi 600. Failure to use agronomic and Integrated Weed Management practices that reduce the weed seed bank in the soil will result in higher weed seed populations. Paddocks with excessively high weed seed banks may have sufficient weed numbers surviving such that final weed control may be considered below a commercially acceptable level and additional herbicide treatments may be necessary. The use of Integrated Weed Management techniques will also reduce the potential for the development or survival of Group D herbicide resistant biotypes.

## WILD OATS

Germinating wild oat seeds lying on soil surface will not be controlled. Therefore, specific wild oat control is only possible with shallow cultivation. Poor control will occur on self-mulching soils and all soil types where deep cultivation is practiced.

## COMPATIBILITY

This product may be mixed in the spray tank with: **Herbicides:** Broadstrike, Spinnaker, Chlorsulfuron, cyanazine, metribuzin, paraquat, Conquest Diuron 900 WG, Conquest Simaquest 900 WG, triasulfuron, Tri-Allate, Metribuzin. **Insecticides:** Conquest Chlorpyrifos 500 EC, endosulfan.

### NOTE:

1. As formulations of other manufacturer's products are beyond the control of Conquest Crop Protection, all mixtures should be tested on a small scale in the spray tank.

2. Tank mixing instructions.

Fill the spray tank  $\frac{1}{4}$  full of water and agitate. Add wettable powders and water dispersible granules first. Agitate until these are uniformly dispersed, meanwhile adding water until the tank is 90% full. Add suspension concentrates (flowables) then soluble concentrates. Emulsifiable concentrates go in last. Top off the tank with water and continue agitation until all the ingredients are properly mixed. Observe any mixing sequence instructions for tank mix products.

## EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible; otherwise, spray tanks, pumps, lines and nozzles should be thoroughly rinsed several times with clean water following application. Spraymate Tank & Equipment Cleaner is suitable for this purpose and will also remove Trifluralin stains.

## RESISTANT WEEDS WARNING

<b>GROUP</b>	<b>D</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Conquest Tricon Maxi 600 Selective Herbicide is a member of the Dinitroaniline group of herbicides. Conquest Tricon Maxi 600 Selective Herbicide has the inhibitors or tubulin formation mode of action. For weed resistance management Conquest Tricon Maxi 600 Selective Herbicide is a Group D herbicide.

Some naturally occurring weed biotypes resistant to Conquest Tricon Maxi 600 Selective Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Tricon Maxi 600 Selective Herbicide or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use Conquest Crop Protection Pty Ltd accept no liability for any losses that may result from the failure of Conquest Tricon Maxi 600 Selective Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Conquest representative.

## PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT use in high winds

DO NOT exceed rates specified, to avoid crop damage.

DO NOT plant sensitive grasses such as oats, sorghum, millets, phalaris spp, ryegrass or wheat for 12 months following the use of this product except where wheat follows wheat or other winter crops.

DO NOT plant oilseed poppies when a detectable residue of Trifluralin is present in the soil. Levels as low as 0.02 ppm may interact with other unfavourable factors (moisture, stress disease etc) to reduce poppy growth and vigour.

DO NOT apply to orchards and vineyards after first flush of growth or when residues can lodge on or in fruit.

Reduced germination of wheat and barley may occur due to combination of following circumstances and use of this product:

- Short Coleoptile cultivars
- Use of seed dressings (except Vitavax)
- Shallow or uneven seedling depth.

**Drift Warning:** DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

#### **RE-ENTRY/RE-HANDLING STATEMENT**

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

#### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Do not store below 5°C. Do not store near food, feedstuffs, fertilisers or seed. The method of disposal of the container depends on the container type. Read the STORAGE and Disposal instructions on the label that is attached to the container.

#### **SPILL AND LEAK MANAGEMENT**

Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways and drains. **Small spills/leaks:** Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal.

#### **SAFETY DIRECTIONS**

Harmful if swallowed. Poisonous if inhaled. Will damage eyes. Will irritate the skin. May irritate nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. If product or spray on skin, immediately wash area with soap and water. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist, a washable hat and elbow length chemical resistant gloves, goggles and half face piece respirator. If product or spray in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

#### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (Tel: 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

#### **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

#### **CONDITIONS OF SALE**

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

**For specialist advice in an emergency dial**

**1 800 033 111**

**24 hours Australia wide.**

**Conquest Crop Protection Pty Ltd.**

ABN 84 098 814 932

76 Walters Drive

Osborne Park WA 6017

Telephone (08) 9347 0500

Facsimile: (08) 9347 0551

APVMA Approval No: 65526 / 58930

