

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BUZZ ULTRA 750 WG FUNGICIDE

ACTIVE CONSTITUENT: 750 g/kg TEBUCONAZOLE

For the control of various diseases of bananas, peanuts, grapes, cereal crops, vegetables and other crops as specified in the Directions for Use Table.

GROUP **3** FUNGICIDE



CONTENTS: 1kg, 5 kg- 30kg

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

Sulphur Mills Australia Pty Ltd
Level 15 Adelaide Street
Brisbane Qld 4000
Tel (07)30030000

APVMA Approval number: 65600/61942
Date of manufacture:
Batch number:

DIRECTIONS FOR USE**DO NOT apply by aerial application methods to wheat, oats, barley****DO NOT apply by equipment carried on back of user (backpack/knapsack) to wheat, oats, barley**

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Bananas (Qld, NSW, WA, NT only)	Leaf Spot (Yellow Sigatoka), Leaf Speckle, Black Sigatoka	130g/ha Add 3-6L water miscible oil/ha	H 1 day	<p>Maintain good deleafing practices to reduce disease inoculum. Very old leaves, and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of Buzz Ultra 750 WG Fungicide.</p> <p>Tropical areas (e.g. North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a maximum of 2 consecutive Buzz Ultra 750 WG Fungicide sprays at 14 day intervals. do NOT apply more than 6 Buzz Ultra 750 WG Fungicide sprays in any 12 month period. do NOT apply any Buzz Ultra 750 WG Fungicide sprays in the months of July, August, September and October.</p> <p>Sub-tropical areas (e.g. South Qld, NSW): Commence spraying with Buzz Ultra 750 WG Fungicide at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a maximum of 2 consecutive Buzz Ultra 750 WG Fungicide sprays. do NOT apply more than 5 Buzz Ultra 750 WG Fungicide sprays in any 12 month period. Buzz Ultra 750 WG Fungicide is approved for use in banana plantations interplanted with avocados.</p> <p>Ground application: Apply by misting machine or airblast sprayer in a convenient volume of water.</p>
Grapevines	Powdery Mildew	Dilute Spraying 17 g/100 L Concentrate spraying Refer to	H 8 weeks	Apply thoroughly as part of the following 5 spray program: 1. when shoots 10-20 cm long. 2. pre-flowering. 3. flowering. 4. after fruit set. 5. before bunches close. This use is subject to a Croplife Australia Fungicide Resistance Management Strategy. Do NOT apply more than two

		'Special instructions for grapevines' in the Application section		consecutive sprays of Buzz Ultra 750 WG Fungicide. Do NOT apply more than three sprays of Buzz Ultra 750 WG Fungicide per season. Do NOT use Buzz Ultra 750 WG Fungicide curatively. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use in equipment that requires concentrate rates greater than 150ml/100 L of water (5x). Do NOT apply in volumes less than 250 L/ha. Add a non-ionic wetting agent at 10mL/100 L regardless of whether applying by dilute or concentrate spraying.
Green beans	Rust	200 g/ha + Agridex* 1 L/ha	G 3 days H 3 days	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high. Applied by ground rig only.
Lettuce	Sclerotinia Rot	200 g/ha	H 5 weeks	Apply by boom spray. Apply only during the early stages of plant development (note the 5 week WHP). Apply a maximum of 2 applications at 7-10 day intervals in rotation with other chemicals registered for this use. Apply only to field crops. DO NOT apply in greenhouse (i.e. protected cropping) situations or hydroponically grown lettuce crops. Control of Sclerotinia in lettuce should not be solely reliant on fungicides. Other control measures such as crop rotations, resistant varieties and planting techniques should be employed.
Onions (TAS only)	White Root Rot	1 g/100 m of row mixed with 145-218 g lime super/100 m of row	-	Before sowing, apply Buzz Ultra 750 WG Fungicide onto lime super. Ensure good coverage of all lime super particles. Apply Buzz Ultra 750 WG Fungicide treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a band width of 2 cm. Ensure that the correct rate of Buzz Ultra 750 WG Fungicide is used otherwise some delay in emergence and reduced stands of seedlings may occur
Pawpaw	Black Spot	170 g/ha	H 3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of Buzz Ultra 750 WG Fungicide. Apply Buzz

				Ultra 750 WG Fungicide at 14 day intervals. Ensure thorough coverage of leaves and fruit. Alternate Buzz Ultra 750 WG Fungicide with sprays of a protectant fungicide. DO NOT apply more than 6 sprays of Buzz Ultra 750 WG Fungicide (or any Group 3 fungicide) on any block in any 12 month period.	
Peanuts (South Qld, NSW only)	Early leaf Spot, late leaf Spot, Rust, Net Blotch	Low disease 100 g/ha + Agridex 1 L/ha	H 21 days weeks G 21 days	Regularly check high risk areas in the crop for disease (eg lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found then either; – spray immediately after the last disease conducive weather (eg rain or heavy dews) OR – spray before the next disease conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat before or just after the next disease conducive weather. Ground application: Apply in at least 100 L of water/ha. Use of Buzz Ultra 750 WG Fungicide on peanuts is subject to a CropLife Australia Fungicide Resistance Management Strategy. Apply no more than 3 consecutive Group 3 sprays alone (eg. Buzz Ultra 750 WG Fungicide) before changing to a fungicide of a different activity group. Apply a maximum of 5 Group 3 sprays per season	
		High disease OR wet weather 166 g/ha + Agridex 1 L/ha			
		High disease AND wet weather 250 g/ha + Agridex 1 L/ha			
Peanuts (North Qld, WA, NT only)	Early leaf Spot, late leaf Spot, Rust	Low disease 132 g/ha + Agridex 1 L/ha	H 21 days G 21 days	Begin spraying at 3-4 weeks after planting. If band spraying, apply half the appropriate rate as a 45 cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14 day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days. Ground application: Apply in at least 100 L of water/ha. Use of Buzz Ultra 750 WG Fungicide on peanuts is subject to a CropLife Australia Fungicide Resistance Management Strategy. Apply no more than 3 consecutive Group 3 sprays alone (eg. Buzz Ultra 750 WG Fungicide) before changing to a fungicide of a different activity group. Apply a maximum of 5 Group 3 sprays per season	
		Moderate disease 166 g/ha + Agridex 1 L/ha			
		Severe disease 250 g/ha + Agridex 1 L/ha			
	Net Blotch	Low disease 166 g/ha + Agridex 1 L/ha			Apply at 14 day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. Ground application: Apply in at least 100 L of water/ha. Use of Buzz Ultra 750 WG Fungicide on peanuts is subject to a CropLife Australia Fungicide Resistance Management Strategy. Apply no more than 3 consecutive Group 3 sprays alone (eg. Buzz Ultra 750 WG Fungicide) before changing to a fungicide of a different activity group.
		Moderate - severe disease 250 g/ha +			

		Agridex 1 L/ha		Apply a maximum of 5 Group 3 sprays per season.
Peas	Powdery Mildew	83 g/ha	H 3 days G 3 days	Apply at flowering or at first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions. Ground application: Apply in at least 50 L of water/ha.
Pyrethrum	Sclerotinia sclerotiorum	200 g/ha	-	Apply twice, in rotation with other control measures, at 7-10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf Rust, Stem Rust	166 g/ha	-	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or reinfection may require a further application 3-4 weeks later. Apply in at least 100 L of water/ha. Ensure thorough coverage, and use higher water volumes in dense or advanced crops.
Sugar cane (variety Q124 only)	Orange Rust	166 g/ha + Agridex 1 L/ha	H 4 weeks G 4 weeks	Even low levels of orange rust suppress yields so it is important to apply Buzz Ultra 750 WG Fungicide early in the development of the disease epidemic. Begin monitoring disease levels early. Check crops at least weekly when climatic conditions favour the development of disease. Apply as a foliar spray when disease begins to escalate rapidly. Repeat application after 14 days if conditions remain favourable to orange rust spore germination. DO NOT apply more than 2 Buzz Ultra 750 WG Fungicide sprays per season. DO NOT apply if heavy rains or storms that are likely to cause surface runoff are forecast with greater than 50% probability within 24 hours (48 hours if possible) of application. Ground application: Use droppers and directed sprays and sufficient water volume to ensure thorough coverage.
Wheat	Leaf Rust, Stripe Rust, Septoria Nodorum Blotch, Yellow leaf Spot	83 or 166 g/ha	H 5 weeks G 14 days	Use higher rate when longer disease control is required. Stripe Rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil (e.g. D-C-Trate or equivalent) at 1%, may improve performance of Buzz Ultra 750 WG Fungicide on wheat, oats and barley. Ground application: Apply in at least 50 L of water/ha.
	Septoria Tritici Blotch	166 g/ha		
Oats	Crown Rust	83 or 166 g/ha		
Barley	Scald	83 g/ha	H 5 weeks G 14 days	Apply at late tillering to early jointing. Ground application: Apply in at least 50 L of water/ha
	Powdery Mildew	83 or 166 g/ha		Apply when 5% of the leaf area is infected. Use higher rate when longer

				disease control is required. Ground Application: Apply in at least 50 L of water/ha.
Wheat, oats	Stem rust		H 5 weeks G 14 days	Apply if more than 5% of stems become infected between full flag leaf emergence to late flowering. Where stem rust is the major disease, yield responses are usually optimised by delaying application until full head emergence, and using the higher rate. In severe cases, if a majority of stems are infected prior to full head emergence, apply at 83 g/ha as soon as possible and if necessary, repeat after 3 weeks when heads are fully emerged. Ground application: Apply in at least 50 L of water/ha.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS H = harvest, G = grazing

Beans, Peas: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 DAYS AFTER APPLICATION.

Cereals, Lettuce: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION. Cereals: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

Grapevines: DO NOT HARVEST FOR 8 WEEKS AFTER APPLICATION.

Onions: NOT REQUIRED WHEN USED AS DIRECTED.

Pawpaw: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

Peanuts: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

Sugar cane: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

Bananas: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

GENERAL INSTRUCTIONS

SPECIAL WARNING- BANANAS

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking. Never include adjuvants other than water miscible oils with Buzz Ultra sprays. The inclusion of wetting agents is known to cause phytotoxicity to young fruit.

Foliar diseases on cereal crops

DO NOT apply to cereal crops more than once per season (166 g/ha rate) or more than twice per season (83 g/ha rate). Treatment will give approximately three weeks disease suppression. Economic responses may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application.

Spray timings for Stripe Rust control

Obtain advisory literature from department of Agriculture for classification of resistant and susceptible varieties. The following spray program is suggested:

Seedling infections: When approximately 20 out of 100 leaves show first signs of infection during tillering to jointing, apply a spray within one week.

Adult infections - susceptible varieties: When approximately 10 out of 100 leaves show first sign of infection, apply a spray within one week. DO NOT delay.

Adult infections - moderately susceptible varieties: When approximately 15 to 20 leaves out of 100 leaves show first sign of infection, apply a spray within one week. DO NOT delay.

Adult infections - moderately resistant and resistant varieties: Monitor carefully. If rust appears and spreads, spray within one week.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Buzz Ultra 750 WG Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, SULPHUR MILLS AUSTRALIA PTY LIMITED accepts no liability for any losses that result from failure of this product to control resistant fungi.

EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Buzz Ultra 750 WG Fungicide. If you are growing edible produce for export, please

check with SULPHUR MILLS AUSTRALIA PTY LIMITED for the latest information on MRLs and import tolerances before using Buzz Ultra 750 WG Fungicide.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of Buzz Ultra 750 WG Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of non-ionic wetting agent and mix thoroughly.

APPLICATION

Aircraft should fly as low as possible under the prevailing conditions to minimise drift

SPECIAL INSTRUCTIONS FOR GRAPEVINES

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- determine an appropriate dilute spray volume (See dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. dilute spray volume as determined above: For example 1500 L/ha .
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 3 X (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$)

4. As the dilute label rate is 30ml/100 L, then the concentrate rate becomes 3 x 30, that is 90 mL/100 L of concentrate spray.

- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- DO NOT use a concentrate rate higher than that specified in the Critical Comments.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

PRECAUTIONS

Re-entry Period

DO NOT allow entry into treated areas for 18 days when applying to wheat, oats, barley unless wearing cotton overalls buttoned to the neck and wrist and chemical resistant gloves. Clothing must be laundered after each day's use

DO NOT allow entry into treated areas for 5 days when applying to grapevines unless wearing cotton overalls buttoned to the neck and wrist and chemical resistant gloves. Clothing must be laundered after each day's use

Aerial application

DO NOT apply by aerial application methods to wheat, oats, barley

DO NOT apply by equipment carried on back of the user to wheat, oats, barley

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, drains or waterways with the chemical or used containers. A spraydrift minimisation strategy should be employed at all times. Spray drift may occur under adverse meteorological conditions or from certain spraying equipment.

DO NOT allow spray to drift onto sensitive areas including, but not limited to, susceptible plants/crops, cropping land, pasture, natural streams, rivers, wetlands, waterways or human dwellings.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, puncture or shred and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes and skin. When opening the container and preparing the product for use, wear cotton overalls over normal clothing, buttoned to the neck and wrist and elbow-length chemical resistant gloves and face shield or goggles. When using the prepared spray, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, a washable hat and elbow-length chemical resistant gloves. If product on skin, immediately wash area with soap and water and if product in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from Sulphur Mills Australia Pty Ltd.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so Sulphur Mills Australia Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.