

POISON

KEEP OUT OF REACH OF CHILDREN.
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide

ACTIVE CONSTITUENT: 200g/L QUIZALOFOP-P-ETHYL
SOLVENTS: 592g/L HYDROCARBON LIQUID
150g/L N-METHYL-PYRROLIDONE



GROUP **A** HERBICIDE

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.**

CONTENTS	APVMA Approval Number
[] 10L	65753/51795
[] 20L	65753/51795
[] 110L	65753/51795
[] 200L	65753/51795
[] 1000L	65753/51795

Distributed by
Corona Industries Pty.Ltd
ACN 099 280 529
84 Ocean Rd, Spearwood, WA, 6163,
Tel: (08) 9434 2400 Fax: (08) 9434 2400

Batch Number:
D.O.M:



PRECAUTION: RE-ENTRY

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL**(20 and 200L only)**

DO NOT store tank-mixes of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. When storing, keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

If the container is a returnable/refillable type, when empty, or contents no longer required, return the container to the point of purchase. This type of container remains the property of Shandong Rainbow International Co., Ltd.

Refillable Containers:**(1000L only)**

Store in closed, original container in a cool, well – ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container:

Store the original sealed ENVIRODRUM in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with Micro Matic valve or security seal. DO NOT contaminate the ENVIRODRUM with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the ENVIRODRUM have been used, please return the empty ENVIRODRUM to the point of purchase. The ENVIRODRUM remains the property of Shandong Rainbow International Co., Ltd.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length PVC gloves. If product on skin immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

For further information refer to Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

**THIS PRODUCT IS CLASSIFIED AS A DANGEROUS
GOOD UNDER THE AUSTRALIAN CODE FOR THE
TRANSPORT OF DANGEROUS GOODS BY ROAD AND RAIL.**

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

**FOR SPECIALIST ADVICE,
CONTACT EMERGENCY
RESPONSE COORDINATOR:
0403 017 534
ENVIRONMENTALLY HAZARDOUS SUBSTANCE LIQUID
N.O.S (CONTAINS QUIZALOFOP-P-ETHYL)
UN 3082**



POISON

**KEEP OUT OF REACH OF CHILDREN.
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide

ACTIVE CONSTITUENT: 200g/L QUIZALOFOP-P-ETHYL

SOLVENTS: 592g/L HYDROCARBON LIQUID

150g/L N-METHYL-PYRROLIDONE

GROUP **A HERBICIDE**

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

**IMPORTANT: READ COMPLETE DIRECTIONS FOR
USE BEFORE USING THIS PRODUCT.**

APVMA Approval No: 65753/51795

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

Distributed by
Corona Industries Pty.Ltd
ACN 099 280 529
84 Ocean Rd, Spearwood, WA, 6163,
Tel: (08) 9434 2400 Fax: (08) 9434 2400

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).

DO NOT apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.

DO NOT apply under conditions of high temperature and low humidity.

DO NOT apply if rainfall is expected within 3 hours.

DO NOT apply less than 50L prepared spray/ha – see section "Ground Spraying".

BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES.

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/ ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Canola Carrots Cauliflower Chickpeas Clover & subclover pastures Cucumbers•	Annual Ryegrass (<i>Lolium rigidum</i>) Brome Grasses (<i>Bromus</i> spp.)	3 leaf to early tillering	150mL or 190mL	14 days 9 weeks 11 weeks 10 weeks 14 days 12 weeks -	-- -- 4 weeks -- -- 4 weeks 14 days	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
Faba beans Field peas Honey dew melons• Lentils Lucerne Lupins	Barley grass (<i>Hordeum</i> spp) Volunteer Barley Volunteer Wheat	3 leaf to early tillering	125mL	14 days 12 weeks 9 weeks 9 weeks 12 weeks -	-- 4 weeks 4 weeks -- 4 weeks 4 weeks 4 weeks	Apply when weeds are actively growing*
Medic pasture Mung beans Navy beans Onions Potatoes Pumpkins• Radish Tomatoes Vetch	Wild Oats (<i>Avena</i> spp)	3 leaf to mid tillering	65mL or 125mL	- 12 weeks 12 weeks 18 weeks 10 weeks 9 weeks 21 days 4 weeks 12 weeks	14 days 4 weeks 4 weeks -- -- -- -- -- 4 weeks	Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control*

*Always add a surfactant/wetting agent at the recommended rate. See section on "use of surfactant/wetting agent" for details.

•**IMPORTANT: DO NOT** apply Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES.

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/ ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Carrots Cucumbers* Green beans Honey dew melons* Lucerne Mung beans Navy beans Onions Peanuts Potatoes Pineapples Pumpkins* Radish Soybeans Sunflower Tomatoes	Awnless Barnyard Grass (<i>Echinochloa colona</i>) Barnyard Grass (<i>Echinochloa crusgalli</i>)	3 leaf to early tillering	250mL or 375mL	14 days 9 weeks 14 weeks 10 weeks 14 days 5 weeks	-- -- -- -- -- 4 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Couch grass (<i>Cynodon dactylon</i>)	2 – 4 leaf seedling	250mL	9 weeks - 12 weeks 12 weeks 18 weeks 11 weeks	-- 4 weeks 4 weeks 4 weeks -- 11 weeks	Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed*.
	Dinebra (<i>Dinebra retroflexa</i>)	3 leaf to early tillering	125mL or 190mL	10 weeks 7 days 9 weeks 21 days 12 weeks 9 weeks 4 weeks	-- -- -- -- 4 weeks 11 weeks --	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Crowsfoot grass (<i>Eleusine indica</i>)	2 leaf to early tillering	250mL	4 weeks	--	Apply when weeds are actively growing*.
	Foxtail Millet (<i>Setaria italica</i>)	2 leaf to early tillering	125mL or 250mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*
	Columbus grass (<i>Sorghum x almum</i>) Johnson grass (<i>Sorghum halepense</i>) Liverseed grass (<i>Urochloa panicoides</i>)	2 leaf to 3 tiller	250mL or 375mL			
	Rhodes grass (<i>Chloris gayana</i>) Stink grass (<i>Eragrostis ciliaransis</i>)	3 leaf to early tillering	125mL or 250mL			

	Queensland Blue grass (<i>Dichanthium sericeum</i>)	3 leaf to early tillering	250mL			Apply when weeds are actively growing*.
	Summer grass (<i>Digitaria ciliaris</i>)	3 leaf to early tillering	250mL or 500mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*

*Always add a surfactant/wetting agent at the recommended rate. See section on "use of surfactant/wetting agent" for details.

•**IMPORTANT: DO NOT** apply Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

HAND GUN SPRAYING: ALL STATES

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/h a	WITHHOLDING PERIOD		CRITICAL COMMENTS	
				HARVEST	GRAZING		
Beetroot	Awnless	3 leaf to early tillering	65 mL	14 days	--	Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha. Thoroughly wet target weeds (especially well established clumps) without causing runoff. Repeat spraying may be necessary for perennial grass species.	
Cabbage	Barnyard			9 weeks	--		
Cauliflower	Grass			14 days	--		
Canola	(<i>Echinochloa colona</i>)			11 weeks	4 weeks		
Carrots				10 weeks	--		
Chickpeas	Crowsfoot			12 weeks	4 weeks		
Clover & subclover pastures	grass (<i>Eleusine indica</i>)			--	14 days		
Cucumbers*	Paspalum			Seedling to	14 days		--
Faba beans	(<i>Paspalum dilatatum</i>)			Established Clumps	12 weeks		4 weeks
Field peas					9 weeks		4 weeks
Grapes	Johnson grass (<i>Sorghum halepense</i>)	Seedling 2 leaf to early tillering	--	--	Always add a surfactant/wetting agent at the recommended		
Green beans		Rhizome 2 leaf to early tillering	5 weeks	4 weeks			
Honey dew melons*			9 weeks	--			
Lentils			125 mL	12 weeks	4 weeks		

Lucerne	Kikuyu (<i>Pennisetum Clandestinum</i>)	Seedling to Established plants	--	4 weeks	rate – see section on “Use of Surfactant/Wetting agent” for details.
Lupins			6 weeks	4 weeks	
Medic pasture			--	14 days	
Mung beans			12 weeks	4 weeks	
Navy beans			12 weeks	4 weeks	
Non crop			--	--	
Onions			18 weeks	--	
Peanuts			11 weeks	11 weeks	
Pineapples			7 days	--	
Potatoes			10 weeks	--	
Pumpkins•			9 weeks	--	
Radish			21 days	--	
Soy beans			12 weeks	4 weeks	
Sunflowers			9 weeks	11 weeks	
Tomatoes			4 weeks	--	
Vetch			12 weeks	4 weeks	

•**IMPORTANT: DO NOT** apply Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

HARVEST

BEETROOT, CAULIFLOWER, CUCUMBERS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

CABBAGE, FIELD PEAS, HONEY DEW MELON, PUMPKINS, SUNFLOWERS: DO NOT HARVEST FOR 9 WEEKS AFTER APPLICATION.

CARROTS, POTATOES: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

CHICKPEAS, FABA BEANS, LENTILS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

CANOLA, PEANUTS: DO NOT HARVEST FOR 11 WEEKS AFTER APPLICATION.

GREEN BEANS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

GRAPES: NOT REQUIRED WHEN USED AS DIRECTED.

LUPINS: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

ONIONS: DO NOT HARVEST FOR 18 WEEKS AFTER APPLICATION.

PINEAPPLES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

RADISH: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

TOMATOES: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

GRAZING

CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, GREEN BEANS, LENTILS, LUCERNE, LUPINS, MUNG BEANS, NAVY BEANS, SOYBEANS,

VETCH: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.

CLOVER, MEDIC PASTURES (EXCEPT LUCERNE): DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.

PEANUTS, SUNFLOWERS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops.

Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates from the treated foliage to the root system and growing points of the plant. Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

Time of Application

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to curcubit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and / or grazing or cutting for stockfeed.

IMPORTANT: DO NOT apply Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide to curcubit crops before the five true leaf stage.

Spray Preparation

If spray equipment has been previously used for the application of sulfonylurea herbicides eg (Metsulfuron methyl), chlorsulfuron, thifensulfuron methyl, triasulfuron, ensure that the recommended sprayer equipment procedure for the product previously applied has been followed, before using the equipment for the application of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide. Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the required amount of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide directly to the spray tank and mix well. If mixing with mancozeb or carbendazim, add these to the spray tank and mix thoroughly PRIOR to adding Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide. If spray solution has been left standing for more than 3 hours ensure thorough re-agitation before commencement of spraying. DO NOT allow spray mixture to stand overnight, as re-suspension may be difficult.

Use of a Surfactant/Wetting Agent

Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide may be used with either of the following adjuvant system:

- 1) A methylated and/or ethylated seed crop oil (eg Hasten#) at 1L/100L of final spray volume (i.e. 1% v/v).
- 2) A non-ionic surfactant (1000 gai/L) at 200 mL/100L of final spray volume (i.e. 0.2% v/v).

DO NOT use Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide with mineral based crop oils (eg D-C-Trate#) or non-methylated spray oils of crop origin (eg Codacide# oil, Synetrol# oil) as spray adjuvants.

Compatibilities

Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is compatible with the herbicides sethoxydim and clethodim. The recommended surfactant/wetting agent is the methylated and/or ethylated seed crop oil adjuvant system. Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is compatible with the insecticides deltamethrin, omethoate, dimethoate, methomyl, endosulfan, the fungicides carbendazim and mancozeb and the herbicide clopyralid when using the methylated and/or ethylated seed crop oil adjuvant system. Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is compatible with the herbicide diflufenican when using the non-ionic surfactant only adjuvant system. DO NOT add crop oil to tank mixtures of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide /diflufenican. Application of Rainbow Quizalofop-P-Ethyl 200 EC Selective

Herbicide /diflufenican tank mix may result in crop effects, which usually appear as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected. To minimise crop effects apply Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide a minimum of 5 days prior to diflufenican application or a minimum 10 days after diflufenican application. (14-21 days if weeds have been stressed).

Ground Spraying

Use between 50 and 100 litres of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150 litres prepared spray/ha.

Use sprayers with flat fan or hollow cone nozzles at 210-350 kPa to give an even distribution of droplets with no less than 50% of the spray volume in the 150-300 micron range. 1100 flat fan nozzles are preferred. For proper spray coverage adjust the boom and nozzle height according to the specification listed by the nozzle manufacturers. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained. Avoid overlapping boomspray widths and shut off spray booms while starting, slowing or stopping.

Sprayer cleanup

Thoroughly clean all traces of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops. If spray equipment has been previously used for the application of sulfonyleurea herbicides eg (Metsulfuron methyl), thifensulfuron methyl, chlorsulfuron, triasulfuron ensure that the recommended sprayer cleanup procedure of the product previously sprayed has been followed before using the equipment for the application of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide.

RESISTANT WEEDS WARNING

GROUP A HERBICIDE

Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is a member of the "Fop" (aryloxyphenoxy propionate) group of herbicides. Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Shandong Rainbow International Co., Ltd accepts no liability for any losses that may result from failure of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, Rainbow's Representative, local Department of Agriculture or Primary Industries.

PRECAUTION: RE-ENTRY

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

(20 and 200L only)

DO NOT store tank-mixes of Rainbow Quizalofop-P-Ethyl 200 EC Selective Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. When storing, keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for this purposes clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

If the container is a returnable/refillable type, when empty, or contents no longer required, return the container to the point of purchase. This type of container remains the property of Shandong Rainbow International Co., Ltd.

Refillable Containers:

(1000L only)

Store in closed, original container in a cool, well – ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container:

Store the original sealed ENVIRODRUM in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with Micro Matic valve or security seal. DO NOT contaminate the ENVIRODRUM with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the ENVIRODRUM have been used, please return the empty ENVIRODRUM to the point of purchase. The ENVIRODRUM remains the property of Shandong Rainbow International Co., Ltd.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length PVC gloves. If product on skin immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

For further information refer to Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

Not a trademark of Shandong Rainbow International Co., Ltd.