

FLOSUL 800 g/L SC FUNGICIDE and MITICIDE**ACTIVE CONSTITUENT: 800g/L SULPHUR (S)**

For the control of certain fungal diseases in grape vines, ornamentals, pome and stone fruit and vegetables and for control of mites in citrus, grape vines, and vegetables in home garden situations

CONTENTS: 250 mL**IMPORTANT: READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.**

SULPHUR MILLS LIMITED AUSTRALIA
Level 15, 15 Adelaide St Brisbane Qld 4000
Tel 07 300 300 00

APVMA Approval number: 66160/60299

Batch number

DOM



HOW TO USE

DO NOT spray in hot weather

DO NOT apply 21 days before or after an oil spray

DO NOT mix with an oil spray

DO NOT spray cucumbers, rockmelons, apricots or other sulphur-sensitive plants.

CROP	DISEASE / PEST	RATE mL/10L water	HOW TO APPLY
Citrus	Brown Citrus Rust Mite	20 to 50	Spray July to August with thorough coverage. Use the higher rate for heavy infestations.
	Citrus Bud Mite, Citrus rust mite	40 to 50	
	White Louse Scale		Either spray at petal fall with copper or spray on its own during spring or autumn. Use the higher rate for heavy infestations
Grapes Vines	Grape leaf Blister Mite Grape leaf Rust Mite	20-35	Apply at 2-4 week intervals depending on the incidence of pests. Use the higher rate for heavy infestations
	Powdery Mildew		Apply when shoots are 20-35 cm long and repeat at intervals of 10-14 days during humid weather. Use the higher rate for heavy infestations
Ornamentals	Powdery Mildew, Two-spotted Mite	20-30	Apply when pest or disease first seen and repeat as necessary.
Pome fruit (e.g. apples and pears)	Powdery Mildew Black Spot	20-35	Apply at pink bud petal fall and at intervals of 2-3 weeks. Some varieties are sulphur sensitive and care should be taken when spraying these varieties.
Vegetables (except cucumbers and rockmelons)	Two-spotted Mite Powdery Mildew, Rust, Tomato Russet Mite, Bean Spider Mite	20-35	Apply when pest or disease is first seen and repeat as necessary. Use the higher rate in crops with a large leaf cover or when conditions favour disease and/or mite development.
Stone Fruit (except apricot)	Brown Rot	20-35	Apply 4 weeks after petal fall and then as cover sprays
	Rust		Apply after petal fall and then at intervals. Use the higher rate when conditions favour diseases

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

**HOW TO PREPARE
MIXING- SHAKE THE CONTAINER WELL BEFORE USE**

Place half the water in the spray pack or backpack sprayer, and add the required amount of Flosul, then add the remaining water to the correct level. Ensure that the Flosul has mixed thoroughly with the water. After spraying, clean the sprayer thoroughly and flush with clean water.

HOW TO APPLY

Ensure both sides of leaves are sprayed to point of run-off when applying the prepared product

PRECAUTIONS

Avoid application under extremes of climate eg rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated.

STORAGE and DISPOSAL

Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

CAUTION

Do not spray cucumbers, rockmelons, apricots, raspberries and other sulphur sensitive plants. Reduce the dosage to avoid scorching during the hot summer months.

DO NOT allow chemical or used containers or spray to get into drains, sewers, streams or ponds.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier

CONDITIONS OF SALE

Sulphur Mills Limited Australia will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Sulphur Mills Limited Australia is limited to the replacement of the goods or (at the option of Sulphur Mills Limited Australia) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Sulphur Mills Limited Australia within thirty days of delivery.



FLOSUL 800 g/L SC FUNGICIDE and MITICIDE

ACTIVE CONSTITUENT: 800g/L SULPHUR (S)

GROUP M2 FUNGICIDE

For the control of certain fungal diseases in grape vines, ornamentals, pome and stone fruit and vegetables and for control of mites in citrus, grape vines, and vegetables

CONTENTS: 500mL; 1-20L; 110L; 500L: 1000L


IMPORTANT: READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

SULPHUR MILLS LIMITED AUSTRALIA
Level 15, 15 Adelaide St Brisbane Qld 4000
Tel 07 300 300 00

APVMA Approval number: **66160/60299**

Batch number

DOM



DIRECTIONS FOR USE FOR USE IN ALL STATES

Restrictions:

DO NOT spray in hot weather

DO NOT apply 21 days before or after an oil spray

DO NOT mix with an oil spray

CROP	DISEASE / PEST	RATE mL/100L water	CRITICAL COMMENTS
Citrus	Brown Citrus Rust Mite	200 to 500	Spray July to August with thorough coverage. Use the higher rate for heavy infestations.
	Citrus Bud Mite, Citrus rust mite	400 to 500	
	White Louse Scale		Either spray at petal fall with copper or spray on its own during spring or autumn. Use the higher rate for heavy infestations
Grapes Vines	Grapeleaf Blister Mite Grapeleaf Rust Mite	200-350	Apply at 2-4 week intervals depending on the incidence of pests. Use the higher rate for heavy infestations
	Powdery Mildew		Apply when shoots are 20-35 cm long and repeat at intervals of 10-14 days during humid weather. Use the higher rate for heavy infestations
Ornamentals	Powdery Mildew, Two-spotted Mite	200-300	Apply when pest or disease first seen and repeat as necessary.
Pome fruit	Powdery Mildew Black Spot	200-350	Apply at pink bud petal fall and at intervals of 2-3 weeks. <u>Some varieties are sulphur sensitive and care should be taken when spraying these varieties. Never spray when hot or with full dose rate.</u>
Vegetables (except cucumbers and rockmelons)	Two-spotted Mite Powdery Mildew, Rust, Tomato Russet Mite, Bean Spider Mite	200-350	Apply when pest or disease is first seen and repeat as necessary. Use the higher rate in crops with a large leaf cover or when conditions favour disease and/or mite development.
Stone Fruit (except apricot)	Brown Rot	200-350	Apply 4 weeks after petal fall and then as cover sprays
	Rust		Apply after petal fall and then at intervals. Use the higher rate when conditions favour diseases

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

MIXING- SHAKE THE CONTAINER WELL BEFORE USE

Place half the water in the spray tank, add the required amount of Flosul whilst agitating, then add the remaining water to the correct level. Maintain agitation until spraying is completed. The addition of Wetter 1000 at 0.1% of water volume to improve coverage, spray retention and rain fastness is strongly recommended. After spraying empty all filters, spray lines etc of dilute spray and clean the sprayer tank thoroughly. Flush tank with clean water and pump through booms to clean all parts of the sprayer

TANK MIXING COMPATIBILITY

Flosul is compatible with a wide range of other crop protection chemicals and micronutrients provided vigorous agitation is maintained throughout the mixing and spraying operation. When mixing copper products, make sure Flosul is fully dispersed before adding copper. It is also compatible with dinocop 50% EC formulations and dimethoate formulations, provided the sulphur is added to the other diluted pesticide and agitation is maintained throughout the mixing and spraying operation. The safe spraying period for both products must be observed. Do not mix with deltamethrin, DSM, mineral oils and high N fertilisers. Consult Sulphur Mills Limited Australia if in doubt.

PROTECTION OF CROPS

Do not spray cucumbers, rockmelons, apricots, raspberries and other sulphur sensitive plants. Reduce the dosage to avoid scorching during the hot summer months.

FUNGICIDE RESISTANCE WARNING

GROUP M2 FUNGICIDE

FLOSUL 800 g/L SC FUNGICIDE AND MITICIDE is a member of the multi-site activity group of fungicides. For fungicide resistance management FLOSUL 800 g/L SC FUNGICIDE AND MITICIDE is a GROUP M2 fungicide. Some naturally occurring individual fungi resistant to FLOSUL 800 g/L SC FUNGICIDE AND MITICIDE and other GROUP M2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by FLOSUL 800 g/L SC FUNGICIDE AND MITICIDE and other Group M2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sulphur Mills Limited Australia accepts no liability for any losses that may result from the failure of FLOSUL 800 g/L SC FUNGICIDE AND MITICIDE to control resistant fungi.

PRECAUTIONS

Avoid application under extremes of climate eg rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated.

STORAGE and DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a well-ventilated area as cool as possible. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

REFILLABLE CONTAINERS

Empty containers fully into application equipment. Close all valves and return to point of supply for refill or storage.

SPILLS MANAGEMENT

Hose small spillage (less than 1 L of concentrate) away with a large excess of water. Large spillage should not be directed to drains. Large spillage should be absorbed in sawdust or sand and the absorbent material containing the product disposed of in a sealed container at an approved disposal site

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with this product or the used containers.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier

CONDITIONS OF SALE

Sulphur Mills Limited Australia will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Sulphur Mills Limited Australia is limited to the replacement of the goods or (at the option of Sulphur Mills Limited Australia) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Sulphur Mills Limited Australia within thirty days of delivery.