



Product Name: CYRO-FLY 60 SPRAY-ON SHEEP BLOWFLY TREATMENT

APVMA Approval No.: 66441/120538

Label Name:	CYRO-FLY 60 SPRAY-ON SHEEP BLOWFLY TREATMENT
-------------	--

Signal Headings:	FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	60 g/L CYROMAZINE
-------------------------	-------------------

Claims:	Will protect long wool sheep for up to eleven weeks against fly strike by cyromazine-susceptible blowfly (<i>Lucilia cuprina</i>) when used as directed. INDICATIONS Cyro-Fly 60 Spray-On Sheep Blowfly Treatment will protect long wool sheep for up to eleven weeks against fly strike by cyromazine-susceptible blowfly (<i>Lucilia cuprina</i>) when used as directed.
---------	--

Net Contents:	1L 5L 10L 20L
---------------	------------------------

Directions for Use:	READ DIRECTIONS FOR USE BEFORE USING THIS PRODUCT
---------------------	---

Restrains:	DO NOT USE on sheep which are producing, or may in future produce, milk or milk products for human consumption.
------------	---

Contraindications:	This product should not be used on sheep with less than 6 weeks wool as a reduced period of protection may result. This product is contraindicated for use on open wounds, e.g. those resulting from marking or mulesing operations. The product should not be applied during gusty or windy weather as spray drift will reduce the period of protection.
--------------------	---

Precautions:	
--------------	--

Side Effects:	
Dosage and Administration:	<p>[DOSE TABLE]</p> <p>[DIAGRAM SHOWING: Positions of bands on sheep for prevention of body, breech and poll strike]</p> <p>CRITICAL COMMENTS</p> <ol style="list-style-type: none"> 1. Use only a Jurox-recommended Spray-On applicator. 2. Ensure the applicator gun is clean. 3. Calibrate applicator gun using an accurate volume measure. 4. Ensure all areas to be protected are treated. 5. When preventing body strike ensure bands overlap along the midline - see diagram 6. This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. Use a diazinon knockdown product as a dressing for existing strike. 7. Heavy rain following the application of this product could diminish the period of protection. 8. Cyro-Fly 60 Spray-On is coloured for band identification. This scourable dye will disappear shortly after treatment. <p>This section contains file attachment.</p>
General Directions:	<p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p> <p>DO NOT DILUTE. DO NOT mix with any other product.</p> <p>Apply directly from the container using a Jurox recommended Spray-On applicator gun. Except for small lambs under 15 kg, the dose volume is determined by the number of months since shearing. Apply Cyro-Fly 60 Spray-On according to the diagrams and tables in the Directions for Use. Cyro-Fly 60 Spray-On may be applied to all breeds of sheep, providing protection against fly strike. Cyro-Fly 60 Spray-On is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25 cm above the fleece, to achieve a 15 cm wide band at each pass.</p>
Withholding Periods:	<p>MEAT: DO NOT USE less than 7 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE on sheep which are producing, or may in future produce milk or milk products for human consumption.</p> <p>WOOL HARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer or the APVMA website (www.apvma.gov.au/residues/).</p> <p>Use of this product may result in wool residues that may not comply with European Union environmental quality standards.</p> <p>Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.</p>

Safety Directions:	Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	
Additional User Safety:	<p>SHEEP REHANDLING INTERVAL: After treatment wait until sheep are dry before rehandling.</p> <p>Additional information is listed in the Safety Data Sheet, available by contacting 1800 023 312.</p>
Environmental Statements:	Protection of Wildlife, Fish, Crustaceans and the Environment: Do not contaminate rivers or streams with the chemical, discarded sump solution or used containers.
Disposal:	<p>[5 L, 10 L, 20 L pack sizes] Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>[alternate statement for 1 L pack] Dispose of empty container by wrapping with paper and putting in garbage.</p>
Storage:	KEEP OUT OF REACH OF CHILDREN. Store below 30°C (room temperature), store in original container, tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

Dose table

MONTHS SINCE SHEARING	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NUMBER OF BANDS	NUMBER OF SHEEP TREATED PER 1L	NUMBER OF SHEEP TREATED PER 5L	NUMBER OF SHEEP TREATED PER 10L	NUMBER OF SHEEP TREATED PER 20L
FOR THE PREVENTION OF BODY STRIKE							
Up to 6	34	17	2	29	147	294	588
7	38	19	2	26	131	263	526
8	44	22	2	22	113	227	454
9	50	25	2	20	100	200	400
10	56	28	2	17	89	178	357
Lambs less than 15kg	17	17	1	58	294	588	1176
FOR THE PREVENTION OF BODY STRIKE AND CRUTCH STRIKE							
Up to 6	51	17	3	19	98	196	392
7	57	19	3	17	87	175	350
8	66	22	3	15	75	151	303
9	75	25	3	13	66	133	266
10	84	28	3	11	59	119	238
Lambs less than 15kg	34	17	2	29	147	294	588
FOR THE PREVENTION OF POLL STRIKE: Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of Cyro-Fly 60 Spray-On to the areas requiring protection, around the poll and horns.							

