

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**EuroChem<sup>®</sup>**  
**DOMICLO<sup>®</sup> 400 WP**  
**Fungicide**

**ACTIVE CONSTITUENT: 400g/kg MYCLOBUTANIL**

<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------



For the Control of Black Spot on Apples and Pears and  
Powdery Mildew on Apples and Strawberries as per the  
Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET THOROUGHLY BEFORE USE**

**Contains 10 x 60g measure packs  
which it is illegal to sell  
separately**

**Contents : 600 gram**

**EuroChem<sup>®</sup>**  
**TGAC Australia Pty Ltd**  
**ACN 134 570 700**  
**9 Heales Rd, Lara, VIC, 3212**  
**Tel : 03-5274 2500, Fax: 03-4206 7012**

### STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

- May irritate the eyes and nose and throat and skin.
- Avoid contact with eyes and skin.
- When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat.
- Wash hands after use.
- After each day's use, wash gloves and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.  
**Phone Australia 131 126.**

### MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet available from your supplier.

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

### MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

TGAC Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. TGAC Australia Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

*® Registered trademark.*

APVMA Approval Number : 66932 / 54713

Batch No : \_\_\_\_\_

Date of Manufacture : \_\_\_\_\_

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**EuroChem<sup>®</sup>**  
**DOMICLO<sup>®</sup> 400 WP**  
**Fungicide**

**ACTIVE CONSTITUENT: 400g/kg MYCLOBUTANIL**

<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

For the Control of Black Spot on Apples and Pears and  
Powdery Mildew on Apples and Strawberries as per the  
Directions for Use Table.

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

**APVMA Approval No : 66932 / 54713**

**EuroChem<sup>®</sup>**  
**TGAC Australia Pty Ltd**  
**ACN 134 570 700**  
**9 Heales Rd, Lara, VIC, 3212**  
**Tel : 03-5274 2500, Fax: 03-4206 7012**

**DIRECTIONS FOR USE****TREE AND VINE CROPS**

<b>RATE</b>			<b>CRITICAL COMMENTS</b>
In the following table, all rates are given for Dilute Spraying. For Concentrate Spraying, refer to the <b>Application Section</b>			For all uses in this table apply by Dilute or Concentrate Spraying equipment. <b>Apply the same total amount of product to the target crop whether applying this product by Dilute or Concentrate Spraying methods</b>
<b>CROPS</b>	<b>DISEASE</b>	<b>RATE per 100L</b>	
<b>Apples Pears</b>	Black spot (scab) <b>Protective treatment</b>	<b>Dilute spraying</b>  12 g	Apply at green tip, or at spur burst following a recommended green tip fungicide spray. Repeat at intervals of 7 to 10 days or as indicated in district spray schedules. After petal fall, and whenever spray intervals may exceed 10 days, add a protectant fungicide. A protectant should always be added to the last EuroChem® DOMICLO® 400 WP Fungicide spray.  The use of EuroChem® DOMICLO® 400 WP Fungicide for black spot (scab) control is subject to a Croplife Resistance Management Strategy.
	Black spot (scab) <b>Curative treatment</b>		Apply within 5 days after the commencement of an infection period.
<b>Apples</b>	Powdery mildew		Apply at early pink stage, and repeat at intervals of 10 to 14 days until December or as indicated in district spray schedules, and depending upon varieties and susceptibility.

**NON TREE AND VINE CROPS**

CROPS	DISEASE	RATE	CRITICAL COMMENTS
Strawberries	Powdery mildew	120 g/ha Use where spray volume is less than 1000 L/ha. Do not apply volumes less than 500L/ha	Reduce background levels of disease by removing plant debris and rotted fruit. Commence spraying before disease appears. Plants should be thoroughly and evenly sprayed to run-off on both sides of the leaf. If conditions favour powdery mildew development, apply <b>EuroChem® DOMICLO® 400 WP Fungicide</b> at 7 to 10 day intervals. To prevent or delay the onset of resistance to Group 3 fungicides, do not apply more than two successive Group C sprays before switching to a fungicide of a different group.
		Where spray volume exceeds 1000L/ha use 12g/100L	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

APPLES and PEARS : DO NOT APPLY LATER THAN 21 DAYS BEFORE HARVEST  
STRAWBERRIES : NOT REQUIRED WHEN USED AS DIRECTED

**GENERAL INSTRUCTIONS****FUNGICIDE RESISTANCE WARNING**

GROUP	3	FUNGICIDE
-------	---	-----------

For fungicide resistance management **EuroChem® DOMICLO® 400 WP Fungicide** is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to **EuroChem® DOMICLO® 400 WP Fungicide** and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by **EuroChem® DOMICLO® 400 WP Fungicide** or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, TGAC Australia Pty Ltd accepts no liability for any losses that may result from the failure of **EuroChem® DOMICLO® 400 WP Fungicide** to control resistant fungi.

## MIXING/APPLICATION:

### Mixing

**EuroChem® DOMICLO® 400 WP Fungicide** is a Wettable Powder (WP) packed in water soluble sachets. Add the required number of water soluble sachets direct to the vat when about one third full with agitators in motion. Add other products after **EuroChem® DOMICLO® 400 WP Fungicide** has dispersed. If premixing, use at least 2 litres of water for each sachet. Do not mix with water with high free chlorine content, or with borax, as it may affect solubility of sachets.

### Application

**EuroChem® DOMICLO® 400 WP Fungicide** may be applied with hand wands or conventional drive pest orchard sprayers.

### Application

#### Dilute spraying – Apples and Pears

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Direction for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

#### Concentrate spraying – Apples and Pears

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. Consult your local advisor, agronomist or Department of Agriculture to determine this volume. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

#### **Concentrate Spraying Example only**

1. Dilute spray volume as determined above: for example 1,500 L/ha
2. Your chosen concentrate spray volume: for example 500 L/ha
3. The concentration factor in this example is: 3 x (ie 1,500 L ÷ 500 L = 3)

4. If the dilute label rate is 12 g/100 L, then the concentrate rate becomes 3 x 12; i.e. 36 g/ 100 L of concentrate spray.

- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

#### **EQUIPMENT MAINTENANCE AND CLEANING**

Ensure spray equipment is well maintained and calibrated before using. Check the pump, nozzles and hose lines for leaks or wear. After use, flush lines thoroughly with clean water.

#### **COMPATABILITY**

**EuroChem® DOMICLO® 400 WP Fungicide** may be mixed with most commonly used insecticides and fungicides. As formulations of other manufacturers' products are beyond the control of TGAC Australia Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

#### **EXPORT OF TREATED PRODUCE:**

**EuroChem® DOMICLO® 400 WP Fungicide** may leave detectable chemical residues in harvested strawberries. Overseas markets may not have appropriate residue tolerances in place or may have established residue tolerances that are lower than Australian maximum residue limit. If you are using this product on destined for export, please contact TGAC Australia Pty Limited or the appropriate industry body or regulatory authority for up to date information.

#### **RE-ENTRY (STRAWBERRIES)**

Do not allow entry into treated area until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

#### **PROTECTION OF LIVESTOCK**

**EuroChem® DOMICLO® 400 WP Fungicide** presents a low hazard to bees. May be applied to plants at any time.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with this product or used container.

#### **STORAGE AND DISPOSAL**

**KEEP OUT OF REACH OF CHILDREN.** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **SAFETY DIRECTIONS**

- May irritate the eyes and nose and throat and skin.
- Avoid contact with eyes and skin.
- When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat.
- Wash hands after use.
- After each day's use, wash gloves and contaminated clothing.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre.

**Phone Australia 131 126.**

### **MATERIAL SAFETY DATA SHEET**

If additional hazard information is required refer to the Material Safety Data Sheet available from your supplier.

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

### **MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

TGAC Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. TGAC Australia Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® *Registered trademark.*



**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**EuroChem<sup>®</sup>**  
**DOMICLO<sup>®</sup> 400 WP**  
**Fungicide**

**ACTIVE CONSTITUENT: 400g/kg MYCLOBUTANIL**

**Contents : 60g**

**Not to be sold separately.**

**Before use read all directions on the outer pack.**  
**Water soluble packaging. Keep Dry.**

**APVMA Number: 66932 / 54713**