

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Kenso Agcare

THREE-WAY

Selective Herbicide



**Active Constituents: 140 g/L BROMOXYNIL present as the octanoate ester
280 g/L MCPA present as the 2-ethylhexyl ester
40 g/L DICAMBA present as the acid
Solvents: 256 g/L HYDROCARBON LIQUID**

GROUP C HERBICIDE

For the control of certain broadleaf weeds in turf and wheat, barley, commercial rye, oats, triticale as per the Directions for Use table.

CONTENTS: 1L, 2.5L, 5L, 10L, 20L, 200L and 1000L

APVMA Approval No: 67012 | 54961



**Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 32179788**

Kenso Agcare Three-way Selective Herbicide

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. Do not inhale vapour or spray mist. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves, face shield and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with soap and water.

After each day's use wash gloves, face shield and goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 32179788

Batch No:
Date of Manufacture:

Kenso Agcare Three-way Selective Herbicide

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Kenso Agcare

THREE-WAY

Selective Herbicide

Active Constituents: 140 g/L BROMOXYNIL present as the octanoate ester
280 g/L MCPA present as the 2-ethylhexyl ester
40 g/L DICAMBA present as the acid
Solvents: 256 g/L HYDROCARBON LIQUID

GROUP **C** | **I** HERBICIDE

For the control of certain broadleaf weeds in turf and wheat, barley, commercial rye, oats, triticale as per the Directions for Use table.

CONTENTS: 1L, 2.5L, 5L, 10L, 20L, 200L and 1000L

APVMA Approval No: 67012 | 54961



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 32179788

Kenso Agcare Three-way Selective Herbicide

DIRECTIONS FOR USE:

RESTRAINTS: DO NOT spray when rain seems likely to occur within 3 hours.

DO NOT spray when weeds are wet with dew or rain, or under stress from drought, low soil fertility, extreme cold or water-logging. DO NOT spray when crop is under stress from drought, excessively wet conditions or low soil fertility. DO NOT apply to crops undersown with legumes.

CROP	CROP STAGE	WEEDS	STATE	RATE	CRITICAL COMMENTS
Recreational Turf, Established Couch (including Queensland Blue), Kikuyu, Paspalum, Fescue, Bent and Rye Lawns		Broadleaf weeds including: Bindy-Eye (<i>Calotis hispidula</i>), Capeweed (<i>Arctotheca calendula</i>), Carrot Weed (<i>Cotula australis</i>), Catsear (<i>Hypochoeris radicata</i>), Caustic weed (<i>Euphorbia drummondii</i>), Clovers (<i>Trifolium</i> spp.), Common Plantain (<i>Plantago</i> spp.), Creeping buttercup (<i>Ranunculus repens</i>), Cudweed (<i>Gnaphalium</i> spp.), Dandelion (<i>Taraxacum officinale</i>), Fleabane (<i>Ictonyza</i> spp.), Funnel Weed (<i>Cotula turbinata</i>), Jo-Jo (Onehunga) (<i>Salvia pterosperma</i>), Thistles, Three cornered Jack (Doublegee) (<i>Emex australis</i>), Wood Sorrel (<i>Oxalis</i> spp.)	NSW, Qld, SA, Vic, WA, ACT only	4-5L in 150L of water per hectare	Apply to actively growing weeds. A slight browning of turf, especially Paspalum, may occur under hot conditions. Use the lower rate for weeds up to the 4 leaf stage. DO NOT mow turf for 2 days after treatment. DO NOT apply to Buffalo Grass. DO NOT feed turf/flawn clippings from treated areas to poultry or livestock.
Wheat, Barley, Oats, Triticale, Commercial Rye	3 leaf to fully tillered from Zadocks (13 to 30)	Group A**: Ball Mustard (<i>Neslia paniculata</i>), Capeweed (<i>Arctotheca calendula</i>), Docks, seedlings only (<i>Rumex</i> sp.), Doudle gee (Spiny emex, Three cornered Jack) (<i>Emex australis</i>), Hedge Mustard (<i>Sisymbrium officinale</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Wild Mustard (<i>Sisymbrium</i> sp), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed) <i>Polygonum aviculare</i> , Yellow burr Weed (<i>Amsinkia</i> sp)	NSW, ACT, Vic, SA, WA only	750mL/ha	3 to 4 leaf stage but no more than 3cm in diameter.
				1L/ha	4 to 6 leaf stage but not more than 5cm in diameter.
	5 leaf to fully tillered (from Zadocks 15 to 30)	Turnip Weed (<i>Rapistrum rugosum</i>), Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>) Deadnettle (<i>Lamium amplexicaule</i>), Vetch (<i>Vicia</i> spp.) Charlock (<i>Sinapsis arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Lesser Swinecress (Bitter cress) (<i>Coronopus didymus</i>)	WA only	1.4L/ha	6 to 8 leaf stage but not more than 10cm in diameter.
			Vic only		
			Tas only		
	3 leaf to fully tillered (from Zadocks 15 to 30cm)	Group B** Climbing Buckwheat (Black bindweed) (<i>Polygonum convolvulus</i>), Common Iceplant (<i>Mesembryanthemum crystallinum</i>), Dense Flowered Fumitory – Red (<i>Fumaria densiflora</i>), Hoary Cress (<i>Cardaria draba</i>), Small Flowered Fumitory – White (<i>Fumaria parviflora</i>), Tree Hogweed (<i>Polygonum patulum</i>), White Iron weed, (Sheepweed, Corn Gromwell (<i>Buglossoides arvensis</i>)	NSW, ACT, Vic, SA, WA only	1L/ha	2 to 4 leaf stage but no more than 3cm in diameter.
1.4L/ha				6 to 8 leaf stage but not more than 10cm in diameter	
5 leaf to fully tillered (from Zadocks 15 to 30)	Slender Thistle (<i>Carduus</i> spp.), Spear Thistle (<i>Cirsium vulgare</i>), Variegated Thistle (<i>Silybum marianum</i>) Wall Fumitory (Pink weed) (<i>Fumaria muralis</i>)	Tas only	1.4L/ha	4 to 6 leaf stage but not more than 100-120mm in diameter. Wall Fumitory is severely stunted by 1.4L/ha.	

**Where GROUP A & B weeds occur together, use the highest application rate.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT CUT OR GRAZE FOR STOCKFOOD FOR 8 WEEKS AFTER APPLICATION.

Kenso Agcare Three-way Selective Herbicide

General Instructions

Note: A temporary wilting may be evident in crops after application. Under some conditions, regrowth and successive germination of weeds may occur following treatment and may require follow-up spraying.

Crop Safety:

DO NOT spray outside crop growth stages recommended as crop damage may result. DO NOT apply to recently released cereal varieties until advice has been sought from the manufacturer or Local Department of Agriculture.

Mixing:

Add the required amount of the product directly to the water in tank and mix well.

Application:

DO NOT apply by misting machines.

For Boom application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare.

For Aerial Application: Apply at least 25 litres of total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

Resistant Weeds Warning

GROUP C I HERBICIDE

Kenso Agcare Three-way Selective Herbicide is a member of the nitriles, phenoxy and benzoic acids groups of herbicides. Kenso Agcare Three-way Selective Herbicide has the inhibitors of photosynthesis at photosystem II and disruptors of plant cell growth mode of action. For weed resistance management, Kenso Agcare Three-way Selective Herbicide is a Group C and Group I herbicide. Some naturally occurring weed biotypes resistant to Kenso Agcare Three-way Selective Herbicide and other Group C and Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Kenso Agcare Three-way Selective Herbicide or other Group C and Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Corporation (M) Sdn. Bhd. accepts no liability for any losses that may result from the failure of Kenso Agcare Three-way Selective Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops.

DO NOT use in close proximity to vines, tomatoes and other susceptible crops including flowers, vegetables, ornamentals, lupins and fruit trees. To prevent damage, DO NOT permit drift or vapour or spray to come into contact with fruit and ornamental trees, cotton, vines, vegetables, flowers, tobacco, lupins or any other field crop. Minimise spray drift by using low pressure and nozzles which do not give a fine droplet size.

DO NOT spray in high winds. DO NOT use this product on or near desirable plants or trees or in locations where the chemical may be washed into contact with their roots.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT use container for any other purpose. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. Do not inhale vapour or spray mist. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves, face shield and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with soap and water.

After each day's use wash gloves, face shield and goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

Kenso Agcare Three-way Selective Herbicide

CONDITIONS OF SALE

Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

In a Transport Emergency
Dial **000**
Police or Fire Brigade



Kenso Corporation (M) Sdn Bhd
3C / 59 Oxford Street
Bulimba QLD 4171
Phone (07) 32179788

APVMA Approval No: 67012 | 54961