

Product Name: BARMAC TRIPLET HERBICIDE
APVMA Approval No: 67538/RV2024



Label Name	BARMAC TRIPLET HERBICIDE
Signal Headings	DANGEROUS POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statement:	ACTIVE CONSTITUENT: 280 g/L MCPA present as the ISO-OCTYL ESTER 140 g/L BROMOXYNIL present as the OCTANOATE ESTER 40 g/L DICAMBA present as the ACID SOLVENT: 289 g/L HYDROCARBON LIQUID
Mode of Action:	GROUP 6 4 HERBICIDE
Statement of Claims:	For the control of certain broadleaf weeds in wheat, barley, cereal rye, oats, triticale and turf as per the Directions of Use.
Net Contents:	
Restrains:	DO NOT spray when rain seems likely to occur within 3 hours. DO NOT spray when weeds are wet with dew or rain, or when under stress from drought, low soil fertility, extreme cold or water logging. DO NOT spray when crop or turf is under stress from drought, frost (extreme cold), excessively wet conditions or low soil fertility. DO NOT apply to crops undersown with legumes.
Directions For Use:	
Other Limitations:	NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
Withholding Period:	DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.
Trade Advice:	
General Instructions:	GENERAL INSTRUCTIONS Note: A temporary wilting may be evident in crops after application. Under some conditions, regrowth and successive germination of weeds may occur following treatment and may require follow-up spraying. CROPSAFETY DO NOT spray outside crop growth stages recommended as crop damage may result. DO NOT apply to recently released cereal varieties until advice has been sought from BARMAC Pty Ltd or the local Department of Agriculture. MIXING Add the required amount of product directly to the water in the tank and mix well.

	<p>APPLICATION DO NOT apply my misting machines. Ground Application: Apply with a properly calibrated boom spray using not less than 50 L water per hectare. Aerial Application: Apply at least 25 L total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.</p>
--	---

<p>Resistance Warning:</p>	<p>GROUP 6, 4 HERBICIDE BARMAC TRIPLET HERBICIDE is a combination of the Nitriles, Phenoxys and Benzoic acid groups of herbicides. BARMAC TRIPLET HERBICIDE has the inhibitors of photo-synthesis at photosystem II and the disruptors of plant cell growth modes of action. For weed resistance management BARMAC TRIPLET HERBICIDE is a Group 6 and Group 4 herbicide. Some naturally occurring weed biotypes resistant to BARMAC TRIPLET HERBICIDE and other Group 6 and Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by BARMAC TRIPLET HERBICIDE or other Group 6 and Group 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, BARMAC Pty Ltd accepts no liability for any losses that may result from the failure of BARMAC TRIPLET HERBICIDE to control resistant weeds.</p>
-----------------------------------	---

Precautions:	
---------------------	--

<p>Protection Statements:</p>	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS Spray equipment must be calibrated accurately prior to working on crops. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT use in close proximity to vines, tomatoes and other susceptible crops including flowers, vegetables, ornamentals, lupins and fruit trees. To prevent damage. DO NOT permit drift or vapour or spray mist to come into contact with fruit and ornamental trees, cotton, vines, vegetables, flowers, tobacco, lupins or any other field crop. Minimise spray drift by using low pressure and nozzles which do not give a fine droplet size. DO NOT use in high wind. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used container.</p>
--------------------------------------	---

<p>Storage and Disposal:</p>	<p>This product must be stored in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.</p> <p>For Non-Refillable Containers: Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with Local, State or Territory government regulations. DO NOT burn empty containers or product.</p>
-------------------------------------	--

	For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs BARMAC Pty Ltd should be advised immediately. Empty containers fully into application equipment. Close all valves and return to point of supply for refill or storage.
--	--

Safety Directions:	Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour or spray mist. Repeated exposure may cause allergic disorders. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and a face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves. If products on eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.
---------------------------	---

First Aid Instructions:	If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting.
--------------------------------	---

First Aid Warnings:	
----------------------------	--

DIRECTIONS FOR USE

CROP	CROP STAGE	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Oats, Triticale, Cereal Rye	3 leaf to fully tillered (from Zadoks 13 to 30)	GROUP A** Ball Mustard (<i>Neslia paniculata</i>), Capeweed (<i>Arctotheca calendula</i>), Docks (seedlings only) (<i>Rumex</i> spp.), Doublegree (Spiny Emex, Three- cornered Jack) (<i>Emex</i> <i>australis</i>), Hedge Mustard (<i>Sisymbrium officinale</i>), Indian Hedge Mustard (<i>Sisymbrium</i> <i>orientale</i>), Wild Mustard (<i>Sisymbrium</i> spp.), Wild Radish (<i>Raphanus raphanistrum</i>), Wireweed (Hogweed) (<i>Polygonum</i> <i>aviculare</i>), Yellow Burr Weed (<i>Amsinckia</i>) (<i>Amsinckia</i> spp.)	QLD, NSW, VIC, SA, WA only	750 mL/ha	3 to 4 leaf stage but not more than 3 cm in diameter
				1 L/ha	4 to 6 leaf stage but not more than 5 cm in diameter
	5 leaf to fully tillered (from Zadoks 13 to 30)	Turnip Weed (<i>Rapistrum rugosum</i>), Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>)	WA only	1.4 L/ha	6 to 8 leaf stage but not more than 10 cm in diameter
		Vetch (<i>Vicia</i> spp.)	VIC only		
	Charlock (<i>Sinapis arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Lesser Swinecress (Bittercress) (<i>Coronopus didymus</i>)	TAS only			
	3 leaf to fully tillered (from Zadoks 13 to 30)	GROUP B** Climbing Buckwheat (Black Bindweed) (<i>Polygonum</i> <i>convolvulus</i>), Common Iceplant <i>Mesembryanthemum crystallinum</i>), Denseflower Fumitory (Red) (<i>Fumaria densiflora</i>), Hoary Cress (<i>Cardaria draba</i>), Small flowered fumitory (white) (<i>Fumaria</i> <i>parviflora</i>), Tree Hogweed (<i>Polygonum palulum</i>) White iron weed (Sheepweed, Corn Gromwell) (<i>Buglossoides arvensis</i>)	QLD, NSW, VIC, SA, WA only	1 L/ha	2 to 4 leaf stage but not more than 3 cm in diameter
	5 leaf to fully tillered (from Zadoks 13 to 30)			1.4 L/ha	6 to 8 leaf stage but not more than 10 cm in diameter
		Slender Thistle (<i>Carduus</i> spp.), Spear Thistle (<i>Cirsium vulgare</i>), Variegated Thistle (<i>Silybum</i> <i>marianum</i>), Wall Fumitory (Pink Weed) (<i>Fumaria muralis</i>)	Tas only	1.4 L/ha	4 to 6 leaf stage but not more than 5 cm in diameter. Wall fumitory is severely stunted by 1.4 L/ha
Recreational turf, Established couch (including Queensland Blue), Kikuyu, Paspalum, Fescue, Bent and Rye Lawns		Broadleaf weeds including: Bindy-eye (<i>Calotis hispidula</i>), Capeweed (<i>Arctotheca calendula</i>), Carrot Weed (<i>Cotula australis</i>), Catsear (<i>Ilypochoeris radicata</i>), Caustic Weed (<i>Euphorbia</i> <i>drummondii</i>), Clovers (<i>Frifolium</i> spp.), Common Plantain (<i>Plantago</i> spp.), Creeping Buttercup (<i>Ranunculus repens</i>), Cudweed (<i>Gnaphalium</i> spp.), Dandelion (<i>Taraxacum officinale</i>), Fleabane (<i>Conyza</i> spp.), Funnel Weed (<i>Cotula turbinata</i>), Jo-Jo (Onehunga) (<i>Saliva pterosperma</i>), Thistles, Three- cornered Jack (Doublegee) (<i>Emex australis</i>), Wood Sorrel (<i>Oxalis</i> spp.)	All States	4-5 L in 150 L water/ha	Apply to actively growing weeds. A slight browning in turf, especially Paspalum, may occur under hot conditions. Use the lower rate for weeds up to the 4 leaf stage. DO NOT mow turf for 2 days after treatment. DO NOT apply to Buffalo grass. DO NOT feed turf/lawn clippings from treated areas to poultry or livestock.

**Where GROUP A & B weeds occur together, use highest application rate.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**